

Class RE46

Book .S97

Copyright No.

COPYRIGHT DEPOSIT.

DISEASES OF THE EYE

SWANZY

A HANDBOOK

OF THE

DISEASES OF THE EYE

AND THEIR

TREATMENT

BY

HENRY R. SWANZY, A.M., M.B., F.R.C.S.I.

SURGEON TO THE ROYAL VICTORIA EYE AND EAR HOSPITAL, AND OPHTHALMIC SURGEON TO THE ADELAIDE HOSPITAL, DUBLIN, EX-PRESIDENT OF THE OPHTHALMOLOGICAL SOCIETY OF THE UNITED KINGDOM

EIGHTH EDITION REVISED

WITH 168 ILLUSTRATIONS

AND ZEPHYR CARD OF HOLMGREN'S TESTS

PHILADELPHIA

P. BLAKISTON'S SON & CO.

1012 WALNUT STREET

1903

RE46
.S97

THE LIBRARY OF CONGRESS,
Two Copies Received
AUG 25 1903
Copyright Entry
July 21, 1903
CLASS a XXc. No
64524
COPY B.

Copyright, 1903, by P. Blakiston's Son & Co.

I DEDICATE THIS BOOK

TO

THEODOR LEBER,

PROFESSOR AT THE UNIVERSITY OF HEIDELBERG

AS A MARK OF

MY ADMIRATION FOR HIS EMINENT SERVICES

TO OPHTHALMOLOGY,

AND OF

MY SINCERE REGARD.

TO THE STUDENT.

You should read carefully Chapters I., II., and III., omitting at first the small print, either before or immediately on joining the Ophthalmic Hospital or Department.

H. R. S.

PREFACE TO THE EIGHTH EDITION.

In the preparation of this Edition the book has been again revised throughout. Moreover, some fresh matter has been introduced, and some obsolete matter omitted, in order to bring the work up to date, without materially increasing its size. The following are some of the additions:—In Chapter IV., A Description of Conjunctivitis Petrificans; in Chapter VI., A Description of Grating Keratitis, Guttate Keratitis, Keratitis Aspergillina, Recurrent Abrasion of the Cornea, and of Injuries of the Cornea caused by blows and Burns; in Chapter VII., A More Detailed Account of Herpes Zoster Ophthalmicus, A Description of Lymphangiodes of the Eyelids, and of Pflüger's Method of Tarsorrhaphy; in Chapter VIII., A More Detailed Account of Kuhnt's Method of Extirpation of the Lacrimal Sac; in Chapter X., in view of the vast importance of the subject, A More Detailed Account of Sympathetic Ophthalmitis; in Chapter XIV., A More Detailed Account of the Use of the Magnet for Foreign Bodies in the Eye; in Chapter XVI., A More Detailed Account of Tumors of the Optic Nerve; in Chapter XVII., A Description of the Eye Symptoms attendant upon Myasthenia Gravis; in Chapter XVIII., A Description of Mr. P. W. Maxwell's Operation for Shrunken Socket, and of Professor Krönlein's Temporary Resection of the Outer Wall of the Orbit for Orbital Tumors, etc.; in Chapter XIX., A Description of Amaurotic Family Idiocy.

I am indebted to Dr. Kathleen Lynn for the great care she has given to the preparation of the Index.

23 Merrion Square,
June, 1903.

PREFACE TO THE FOURTH EDITION.

The Third Edition of this book was published in October, 1890, and I am gratified that the work continues to find favor, not only with students, for whom it is mainly intended, but also with practitioners.

The book has now again been revised throughout, and brought up to date.

In an Appendix, Holmgren's Method for Testing the Color Sense has been described in greater detail than before.

Some new illustrations have been added.

The great difficulty of an author in the preparation of a book like this consists in " Saying not all he might, but all he ought." It is his duty to give a succinct and practical account of his subject in its most modern aspect, without weighting his pages with excessive detail and prolonged discussion. This has been my aim. For deeper and wider information, larger handbooks and original monographs must be consulted.

23 Merrion Square,
October, 1892.

CONTENTS.

CHAPTER I.

PAGE

Numbering of Trial-Lenses and Spectacle Glasses—Normal Refraction and Accommodation—The Meter Angle—The Angle Gamma—The Sense of Sight (Light-Sense, Color-Sense, Form-Sense)—The Field of Vision 17

CHAPTER II.

ABNORMAL REFRACTION AND ACCOMMODATION.

Hypermetropia—Correction of H.—Amplitude of Accommodation in H.—Angle Gamma in H.—Cramp of Ciliary Muscle in H.—Accommodative Asthenopia in H.—Internal Strabismus in H.—The Prescribing of Spectacles in H. 36

Myopia—Determination of Degree of M.—Amplitude of Accommodation in M.—Angle Gamma in M.—Complications of Progressive M.—Functional Anomalies Attendant upon M.—Management of M.—The Prescribing of Spectacles in M.—Operative Cure of M. 42

Astigmatism—Symptoms of As.—Estimation of Degree of, and Correction of As.—The Astigmometer—Lental As.—Irregular As. . 52

Anisometropia 64

Anomalies of Accommodation—Presbyopia—Paralysis of Accommodation—Cramp of Accommodation 64

CHAPTER III.

THE OPHTHALMOSCOPE.

Why Necessary—Helmholtz's Ophthalmoscope—Modern Ophthalmoscope—Direct Method—Indirect Method 70

Estimation of the Refraction by Aid of the Ophthalmoscope—Direct Method—Retinoscopy 76

Focal Illumination 87

The Normal Fundus Oculi as Seen with the Ophthalmoscope—The Optic Papilla, or Optic Disc—The Retina—The Macula Lutea—The General Fundus Oculi—The Retinal Vessels 88

CHAPTER IV.

DISEASES OF THE CONJUNCTIVA.

PAGE

Hyperemia—Conjunctivitis—Catarrhal Conjunctivitis—Acute Blennorrhea of the Conjunctiva, or Purulent Ophthalmia—Follicular Conjunctivitis—Spring Catarrh—Hay Fever—Trachoma, Granular Conjunctivitis, or Granular Ophthalmia—Acute Trachoma, or Acute Granular Ophthalmia—Chronic Trachoma, or Chronic Granular Ophthalmia—Lymphoma of the Conjunctiva—Croupous Conjunctivitis—Diphtheritic Conjunctivitis—Conjunctival Complications of Smallpox—Amyloid Degeneration—Tubercular Disease of the Conjunctiva—Lupus—Pemphigus—Xerosis, or Xerophthalmos—Pterygium—Pinguecula—Subconjunctival Ecchymosis—Nevus—Polypus—Dermoid Tumors—Lipoma—Syphilitic Disease of the Conjunctiva—Papilloma, or Papillary Fibroma—Epithelioma—Sarcoma—Simple Cysts—Subconjunctival Cysticercus—Lithiasis—Conjunctivitis Petrificans—Uric Acid Deposits—Injuries of the Conjunctiva 93

CHAPTER V.

PHLYCTENULAR, OR STRUMOUS, CONJUNCTIVITIS AND KERATITIS.

Solitary, or Simple, Phlyctenula of the Conjunctiva—Multiple, or Miliary, Phlyctenula of the Conjunctiva—Modes of Secondary Corneal Affection—Primary Phlyctenular Keratitis—Different Forms of Same—Symptoms of Phlyctenular Keratitis—Causes of Phlyctenular Ophthalmia—Treatment 133

CHAPTER VI.

DISEASES OF THE CORNEA.

Inflammations of the Cornea—(*a*) Ulcerative Inflammations of the Cornea—Simple Ulcer—Deep Ulcer—Serpiginous Ulcer—Rodent Ulcer—Marginal Ring Ulcer—Absorption Ulcer—Neuro-Paralytic Keratitis—Infantile Ulceration of the Cornea with Xerosis of the Conjunctiva—Herpes—Filamentary Keratitis—Bullous Keratitis—Dendriform Keratitis—Keratitis Aspergillina 140
(*b*) Non-Ulcerative Inflammations of the Cornea—Abscess—Diffuse Interstitial, or Parenchymatous, Keratitis—Keratitis Profunda—Grating-like Keratitis—Guttate Keratitis—Keratitis Punctata—Sclerotizing Opacity—Riband-like Keratitis . . . 161
Ectasies of the Cornea—Staphyloma Corneæ (Evisceration and Mules' Operation)—Conical Cornea, or Keratoconus 169
Tumors of the Cornea 177

PAGE

Injuries of the Cornea—Foreign Bodies—Losses of Substance—Recurrent Abrasion, or Disjunction—Blows—Burns—Perforating Injuries 177
Opacities of the Cornea—Nebula, Macula, Leukoma—Arcus Senilis . 183

CHAPTER VII.

DISEASES OF THE EYELIDS.

Eczema—Herpes Zoster Ophthalmicus—Primary Syphilitic Sores—Secondary Syphilitic Sores—Tertiary Syphilitic Affection—Vaccine Vesicles—Rodent Ulcer—Marginal Blepharitis (Ophthalmia Tarsi)—Phtheiriasis Ciliorum—Hordeolum (Stye)—Chalazion (Meibomian Cyst, Tarsal Tumor)—Milium—Molluscum—Nevus—Xanthelasma—Chromidrosis—Epithelioma, Sarcoma, Adenoma and Lupus—Solid Edema, or Elephantiasis Lymphangioides—Clonic Cramp of the Orbicularis Muscle—Blepharospasm—Ptosis—Operations for its Cure—Lagophthalmos—Symblepharon—Blepharophimosis—Canthoplastic Operation—Distichiasis and Trichiasis—Operations for their Cure—Entropium—Operations for its Cure—Spastic Entropium—Senile Entropium—Operations for its Cure—Ectropium—Operations for its Cure—Ankyloblepharon—The Restoration of an Eyelid—Injuries—Ecchymosis—Epicanthus—Congenital Coloboma 187

CHAPTER VIII.

DISEASES OF THE LACRIMAL APPARATUS.

Malposition of the Punctum Lacrimale—Stenosis and Occlusion of the Punctum Lacrimale—Obstruction of the Canaliculus—Stricture of the Nasal Duct—Blennorrhea of the Lacrimal Sac—Obliteration and Extirpation of the Lacrimal Sac—Acute Dacryocystitis—Dacryoadenitis—Hypertrophy of the Lacrimal Gland . . . 233

CHAPTER IX.

DISEASES OF THE SCLEROTIC.

Inflammations of the Sclerotic—Periodic Transient Episcleritis, or Hot Eye—Episcleritis—Deep Scleritis—Injuries of the Sclerotic—Tumors of the Sclerotic—Pigment Spots 244

CHAPTER X.

DISEASES OF THE UVEAL TRACT.

Iritis — Symptoms — Syphilitic — Rheumatic — Gonorrheal — Causes—Prognosis—Treatment 250

PAGE

Cyclitis 259
Chorioiditis—Disseminated—Syphilitic—Central Senile Guttate—Central—Purulent 261
Sympathetic Ophthalmitis, and Sympathetic Irritation—Introductory—Sympathetic Irritation—Sympathetic Ophthalmitis—Diagnosis—Prognosis—Treatment—Prophylaxis—Therapeutic and Optical Operations used in Sympathetic Ophthalmitis—Pathogenesis . 265
Injuries of the Iris—Punctured Wounds—Foreign Bodies—Iridodialysis — Retroflexion — Rupture — Dehiscence—Traumatic Mydriasis 284
Injuries of the Ciliary Body—Punctured Wounds—Foreign Bodies . 287
Injuries of the Chorioid—Foreign Bodies—Incised Wounds—Rupture—Extravasation of Blood 288
New Growths of the Iris—Cysts—Granuloma—Tubercle—Sarcoma—Ophthalmia Nodosa 289
New Growths of the Ciliary Body—Sarcoma—Myosarcoma—Carcinoma 292
New Growths of the Chorioid—Sarcoma—Carcinoma—Tubercle—Sarcoma Carcinomatosum—Myosarcoma—Osteosarcoma . . 292
Other Diseases of the Chorioid—Posterior Staphyloma—Detachment—Central Senile Areolar Atrophy 297
Malformations of the Iris 297
Malformations of the Chorioid 298
Operations on the Iris 300

CHAPTER XI.

MOTIONS OF THE PUPIL IN HEALTH AND DISEASE.

The Size of the Pupil in Health—Contraction of the Pupil—Dilatation of the Pupil 303
The Action of the Mydriatics and Myotics on the Pupil . . . 309
The Size of the Pupil in Disease—Myosis—Mydriasis . . . 309
Tabular Arrangement of Mydriatics, Myotics, and Anesthetics . 316

CHAPTER XII.

GLAUCOMA.

Primary Glaucoma—Chronic, or Non-Inflammatory, Glaucoma—Acute, or Inflammatory, Glaucoma—Glaucoma Fulminans—Subacute Glaucoma—Etiology—Pathology—Treatment . . . 320
Congenital Hydrophthalmos 340

CHAPTER XIII.

DISEASES OF THE CRYSTALLINE LENS.

Complete Cataracts—Senile Cataract—Progress, Pathogenesis, and Etiology—Treatment 342

PAGE

Complete Cataract of Young People—Diabetic Cataract—Complete Congenital Cataract—Black Cataract 349
Partial Cataracts—Central Lental Cataract—Zonular, or Lamellar, Cataract—Anterior Polar, or Pyramidal, Cataract—Fusiform, or Spindle-Shaped, Cataract 350
Secondary Cataract—Posterior Polar Cataract—Total Secondary Cataract 352
Capsular Cataract 353
Traumatic Cataract 353
Operations for Cataract—Extraction of Cataract—Linear Extraction The Modified Peripheral Linear Extraction—The Three Millimeter Flap Operation—Cataract Extraction without Iredectomy . 355
Discission, or Dilaceration—Suction Operation—Secondary Cataract and its Operation—Capsulotomy—Iridotomy 376
Dislocation of the Crystalline Lens—Lenticonus—Aphakia . . . 381

CHAPTER XIV.

DISEASES OF THE VITREOUS HUMOR.

Purulent Inflammation—Other Inflammatory Affections—Opacities—Muscæ Volitantes—Fluidity (Synchysis)—Synchysis Scintillans—Foreign Bodies in the Vitreous Humor and Interior of the Eye in General—Use of Röntgen Rays in Detecting Foreign Bodies—The Sideroscope—Removal of Foreign Bodies—The Magnet—Cysticercus—Blood-Vessels—Persistent Hyaloid Artery—Detachment 384

CHAPTER XV.

DISEASES OF THE RETINA.

Alterations in Vascularity—Hyperemia—Anemia 400
Retinitis — Syphilitic — Hemorrhagic — Albuminuric—Diabetic—Leukemic—Punctata Albescens—Development of Connective Tissue, or Retinitis Proliferans—Retinitis Circinata—Purulent Retinitis . 401
Atrophy of the Retina—Retinitis Pigmentosa—Gyrate Atrophy of the Retina and Chorioid 407
Diseases of the Retinal Vessels—Apoplexy of the Retina—Embolism of the Central Artery—Thrombosis of the Central Artery—Thrombosis of the Central Vein—Aneurysm of the Central Artery—Sclerosis of the Retinal Vessels (Perivasculitis)—Quinin Amaurosis 409
Injury of the Retina by Strong Light—Sunlight—Snow-Blindness—Effect of Electric Light on the Eyes 415
Tumor of the Retina—Glioma 416
Parasitic Disease of the Retina—Cysticercus 417

PAGE

Detachment of the Retina 418
Traumatic Affections of the Retina—Traumatic Anesthesia—Commotio Retinæ, or Traumatic Edema of the Retina 422

CHAPTER XVI.

DISEASES OF THE OPTIC NERVE.

Optic Neuritis (Papillitis), due to: Cerebral Tumors—Tubercular Meningitis—Hydrocephalus—Tumors of the Orbit—Inflammatory Processes in the Orbit—Exposure to Cold—Suppression of Menstruation—Chlorosis—Syphilis—Rheumatism—Lead Poisoning—Peripheral Neuritis—Multiple Sclerosis—Tabes Dorsalis—Acute Myelitis—and to Hereditary and Congenital Predisposition . . 424
Toxic Central Amblyopia, or Central Scotoma—Retrobulbar Neuritis—Optic Neuritis, Associated with Persistent Dropping of Watery Fluid from the Nostril 429
Atrophy of the Optic Nerve, due to: Optic Neuritis—Pressure—Embolism of the Central Artery of the Retina—Syphilitic Retinitis, Retinitis Pigmentosa, Choriodo-Retinitis, and to Disease of the Spinal Cord (Spinal Amaurosis)—Optic Atrophy as a purely Local Disease 433
Tumors of the Optic Nerve—Hyaline, or Colloid, Outgrowths—Injuries of the Optic Nerve 436
Amblyopia due to Hemorrhage from the Stomach, Bowels, or Uterus—Glycosuric Amblyopia 437

CHAPTER XVII.

THE MOTIONS OF THE EYEBALLS, AND THEIR DERANGEMENTS.

Actions of the Orbital Muscles—Inclination of the Vertical Meridian in the Several Principal Positions—Muscles called into Action in the Several Principal Positions—The Field of Fixation—Strabismus 441
Paralyses of the Orbital Muscles—General Symptoms—Paralysis of the External Rectus—Paralysis of the Superior Oblique—Paralsis of the Internal Rectus, Superior Rectus, Inferior Oblique, and Levator Palpebræ—Ophthalmoplegic Migraine—Ophthalmoplegia Externa, or Nuclear Paralysis—Fascicular Paralysis—Myasthenia Gravis—Cerebral Paralysis of Orbital Muscles—The Localizing Value of Paralysis of Orbital Muscles in Cerebral Disease . 447
Convergent Concomitant Strabismus—Causes—Single Vision in—Amblyopia of Squinting Eye—Clinical Varieties of—Measurement of—Mobility of Eye in—Treatment—Orthoptic Treatment—Operative Treatment—Tenotomy—Advancement of External Rectus—Dangers of the Strabismus Operation—Treatment Subsequent to Operation 471

PAGE

Insufficiency of the Internal Recti, and Divergent Concomitant Strabismus—Muscular Asthenopia—Treatment—Operative Treatment 494
Nystagmus 499

CHAPTER XVIII.

DISEASES OF THE ORBIT.

Orbital Cellulitis—Periostitis of the Orbit—Caries of the Orbit—Injuries of the Orbit—Orbital Tumors—Implication of Neighboring Cavities—Pulsating Exophthalmos—Tumors of the Lacrimal Gland—Hernia Cerebri—Exophthalmic Goiter—Enophthalmos . 501

CHAPTER XIX.

PART I.

OCULAR DISEASES AND SYMPTOMS LIABLE TO ACCOMPANY FOCAL DISEASES OF THE BRAIN.

Hemianopsia—Arrangement of the Cortical Visual Centers, their Relations to the Retina, and the Course of the Optic Fibers between these Two Points—Localization of the Lesion in Hemianopsia—Alexia, or Word-Blindness—Visual Aphasia—Dyslexia—Amnestic Color-Blindness—Visual Hallucinations—Mind-Blindness, or Optic Amnesia 522

PART II.

OCULAR DISEASES AND SYMPTOMS LIABLE TO ACCOMPANY DIFFUSE ORGANIC DISEASES OF THE BRAIN.

Disseminated Sclerosis of the Brain and Spinal Cord—Diffuse Sclerosis of the Brain—General Paralysis of the Insane—Amaurotic Family Idiocy—Meningitis—Traumatic Meningitis—Hydrocephalus—Infantile Paralysis—Paralysis Agitans—Encephalopathia Saturnia—Epilepsy—Chorea 536

PART III.

OCULAR DISEASES AND SYMPTOMS LIABLE TO ACCOMPANY DISEASES AND INJURIES OF THE SPINAL CORD.

Tabes Dorsalis—Hereditary Ataxy—Myelitis—Syringomyelia, and Morvan's Disease—Myotonia Congenita—Acute Ascending Paralysis—Injuries of the Spinal Cord 543

PART IV.

NERVOUS AMBLYOPIA, OR NERVOUS ASTHENOPIA.

PAGE

Nervous Amblyopia in Neurasthenia—Nervous Amblyopia in Hysteria—Nervous Amblyopia in Traumatic Neurosis 547

PART V.

VARIOUS FORMS OF AMBLYOPIA.

Transitory Hemianopsia, or Scintillating Scotoma—Congenital Amblyopia—Reflex Amblyopia—Night-Blindness—Uremic Amblyopia Pretended Amaurosis—Erythropsia 552

APPENDIX I.

Holmgren's Method for Testing the Color-Sense 559

APPENDIX II.

Regulations as to Defects of Vision which Disqualify Candidates for Admission into the Civil, Naval, and Military Government Services, the Royal Irish Constabulary, and the Mercantile Marine . 567

INDEX 567

DISEASES OF THE EYE.

CHAPTER I.

THE NUMBERING OF TRIAL-LENSES.

THE lenses in trial cases and in spectacles are numbered according to the metrical system.

The lens of one meter (39 1-2 inches) focal length is called the Dioptric Unit, or the Dioptry (1 D), of the metrical system. 2 D, 3 D, 4 D, etc., indicate the number of meter lenses, or dioptries, contained in each of these lenses. 2 D is therefore twice as powerful a lens (its focal length only half as long) as 1 D.

Convex lenses are indicated by the + sign placed before their number, thus, + 5 D; and concave lenses by the — sign, thus, — 5 D.

If it be required to ascertain the focal length of a given lens, 100 (1 meter = 100 cm.) is to be divided by the dioptric number of the lens, and the answer will give the focal length in centimeters. For example, the focal length of 10 D is $\frac{100}{10}$ = 10 cm.

If the focal length of the lens be known, and it be desired to ascertain its dioptric number, we find it by dividing 100 cm. by the focal length. For example, if the focal length be 33 cm., then $\frac{100}{33}$ = 3 D.

THE POWERS OF NORMAL REFRACTION AND ACCOMMODATION OF THE EYE.

The eye is a dark chamber with a series of convex refracting surfaces—namely, the cornea, and the anterior and posterior surfaces of the crystalline lens; and certain intraocular or dioptric media—namely, the aqueous humor, the substance of the crystalline lens, and the vitreous humor. By aid of this apparatus, which

is called the dioptric system of the eye, distinct inverted images of external objects are formed on the retina.

The refracting media are centered on the optical axis (*O A*, Fig. 1) an imaginary line which, passing through the optical center (*N*) of the eye, meets the retina at a point (*A*) slightly to the inner side of the macula lutea (*M*).

In treating of the eye we have to consider two sets of visual objects—viz., distant objects and near objects. Distant objects




Fig. 1 (*right eye*).

are those at 6 meters and more from the eye; near objects are those closer to the eye than 6 meters. For practical purposes the rays which pass through the pupil, coming from any given point of a distant object, are as good as parallel, their divergence being so very slight when they reach the eye, and we regard them as being parallel.

Refraction.

By the Refraction of the Eye is meant the faculty it has *when at rest* (*i. e.*, without an effort of accommodation) of altering the direction of rays of light which pass into it, making parallel rays convergent, and divergent rays less divergent.

In Normal Refraction, or Emmetropia (*ἔμμετρον, the standard; ὤψ, the eye*), as it is termed, parallel rays (see Fig. 2, in which the object from which the rays come is supposed to be 6 meters or more from the eye) in passing through the dioptric media are given such a convergence that they are brought to a focus on the layer of rods and cones of the retina, and form on it a distinct inverted image of the point or object from which they come. In other words, the retina is placed at the principal focus of the dioptric system of the normal eye, which is thus adapted for parallel rays, and its far point (*vide infra*) is at infinity.

Accommodation.

The eye can see near objects distinctly as well as distant objects, although the rays from any given point (*a*, Fig. 3) of a near object reach the eye with a divergence so considerable that they could not be brought to a focus on the retina by the unaided refraction, but would converge towards a point (their conjugate




Fig. 2.

focus *a'*) behind the retina, and would not form a distinct image on the latter, but merely a blurred image or circle of diffusion (at *b c*). It is obvious, therefore, that an increase of refracting power in the eye is necessary, in order that near objects may be




Fig. 3.

distinctly seen. It is this increase in the refracting power for the purpose of near vision which is called Accommodation.

The Mechanism of Accommodation is as follows: The ciliary muscle (*m*, Fig. 4) contracts, thus drawing forward the chorioid and the ciliary processes, and relaxing the zonula of

Zinn (*z*), which is attached to the latter. The lens (*l*), which was flattened by the tension of the zonula, is now free to assume a more spherical shape, in response to its own elasticity. The posterior surface of the lens scarcely alters in shape, being fixed in the patellary fossa; but the anterior surface becomes more convex, thus increasing its refracting power. Associated with the act of accommodation is a contraction of the pupil. The acompanying figure (Fig. 4) represents the changes which take place in accommodation, the dotted lines indicating the latter state.

Tscherning[1] has shown that the increased curvature of the anterior surface of the lens occurs mainly in the center of that surface—in other words, that in accommodation the anterior surface becomes somewhat conical, and not merely more spherical. He




FIG. 4.—*c*, cornea; *a*, anterior chamber; *l*, lens; *v*, vitreous humor; *i*, iris; *z*, zonula of Zinn; *m*, ciliary muscle.

holds that this is due to a tightening, and not to a relaxation of the zonula. This theory has been vigorously combated by other observers, and its true value remains to be determined.

The Far Point and the Near Point.—It is possible for the eye to see objects accurately at every distance from its Far Point —*i. e.*, its most distant point of distinct vision (Punctum Remotum,—R.), up to a point only a few centimeters from the eye, called the Near Point (Punctum Proximum,—P.). We can find the latter by directing the patient to look at a page printed in small type, and by bringing it slowly closer and closer to his

eye, until a point is reached where he cannot distinguish the words and letters, which become blurred. A point very slightly more removed from the eye than this, where he can read distinctly, is the near point. Between the near point and the eye vision is indistinct, because no effort of the ciliary muscle can produce the amount of convexity of the lens required for so short a distance.

The Amplitude of Accommodation.—This is the amount of accommodative effort of which the eye is capable—*i. e.*, the effort it makes in order to adapt itself from its Far Point (R.) up to its Near Point (P.). The amplitude of accommodation (a), therefore, is equal to the difference between the refracting power of the eye when at rest (r), and when its accommodation is exerted to the utmost (p), as expressed by the formula $a =$




Fig. 5.

$p - r$. It may be represented by that convex lens placed close in front of the eye, which would take the place of the increased convexity of the lens, or, in other words, which would give to rays coming from the nearest point of distinct vision a direction as if they came from the far point. The number of this lens expresses the amplitude of accommodation in a given eye.

For example: if, in an emmetropic eye (E, Fig. 5) the near point be situated at 20 cm., then a convex lens (L) of 20 cm. focal length placed close to the eye (between that point and the eye) would give to rays coming from the near point a direction (*i. e.*, would make them parallel) as though they came from a distant object, and this normally refracting eye would then be enabled by aid of its refraction alone to bring these rays to a focus on the retina. Making use of the above equation, we find in this case—since a focal length of 20 cm. represents a lens of

5 D—that $a = 5—r$, but R. being situated at infinity we designate it by the sign ∞; hence, $r = \frac{1}{R} = \frac{1}{\infty} = 0$; therefore $a = 5 - 0 = 5$ D.*

The amount of amplitude of accommodation (*i. e.*, the number of the lens which would represent it) is the same in every kind of refraction, according to the age of the individual, but in emmetropia alone is $a = p$ as above, because in it alone is $r = 0$.

Under the head of "Anomalies of Accommodation," chap. ii., will be found Professor Donders' diagram representing the amplitude of accommodation at different ages.

Connection between Accommodation and Convergence (Relative Accommodation).—With every degree of convergence of the visual lines a certain effort of accommodation is associated. Thus, if the object be situated 2 meters from the eye the visual lines converge to that point, and a certain effort of accommodation is made. But this connection between accommodation and convergence is somewhat elastic, for the accommodative effort may be increased or decreased, while the object is kept distinctly in view, and the same convergence maintained. That it may be increased is shown by the experiment of placing a weak concave glass before the eye, when it will be found that the object is still distinctly seen; and if a weak convex glass be then held before the eye the object will also be clearly seen, showing that the accommodative effort may be lessened without affecting vision or convergence. This amplitude of accommodation for a given point of convergence of the visual lines, found by the strongest concave and strongest convex glasses with which the object can still be distinctly seen, is called the Relative Amplitude of Accommodation. That part of it which is already in use, and

*It must be observed that R represents the *distance* of the Far Point from the eye, while r represents the *refractive power* which is added to the eye by accommodation or by a lens in order to adapt it for the distance R. Hence it is evident that $r = \frac{1}{R}$, because the strength, or refractive power, of a lens is inversely as its focal length—*e. g.*, a lens of the strength of 4 D will have a focal length of 1-4 that of a lens of 1 D—*i. e.*, $\frac{1m}{4} = \frac{100 \text{ cm.}}{4} = 0.25$ cm. (see above, Numbering of Trial-Lenses). Similarly, $p = \frac{1}{P}$ and $a = \frac{1}{A}$; P representing the *distance* of the Near Point,, and A the focal length of the lens which represents the Accommodation.

is represented by the convex lens, is termed the negative part; while the positive part is represented by the concave lens, and has not been brought into play. For sustained accommodation at any distance it is necessary that the positive part of the relative amplitude of accommodation be considerable in amount.

Moreover, the convergence may be altered, while the same effort of accommodation is maintained, as is shown by the experi-




Fig. 6.




Fig. 7.

ment of placing a weak prism with its base inwards before one eye. In order that the object may then be seen singly, it will be necessary for the eye before which the prism is placed to rotate somewhat outwards; and it will be found that the individual can do this, while at the same time he sees the object with the same distinctness, showing that the same effort of accommodation has been maintained, although the angle of convergence of the visual axis is less than before.

The Meter Angle.

If the visual line (*E* 1, Fig. 6) of an eye (*E*) have to be brought to bear on a point (1, Fig. 6) 1 meter distant from it in the median line (*M* 1), the angle of convergence (*E* 1 *M*) which the visual line thus makes with the median line is called the Meter Angle. It expresses the degree of convergence necessary for binocular vision at that distance, and is employed as the unit for expressing other degrees of convergence. If, for example, an object be situated 1-2 a meter (1-2, Fig. 6) from the eye, the angle of convergence (*E* 1-2 *M*) must be practically twice as large as at 1 meter: C. (Convergence = 2 meter angles. If the object be only 1-3 of a meter distant, 3 meter angles are required: C. = 3 meter angles. If the object be situated 2 meters from the eye, the angle of convergence will be only one-half as great as at 1 meter, and here C. = 1-2 meter angle; while if the eye be directed towards a distant object (*D*) there will be no angle of convergence, and if the visual lines be divergent the meter angle will be negative.

Now the average normal emmetropic eye requires for each distance of binocular vision as many meter angles of convergence as it requires dioptries of accommodation. For a distance of 1 meter an effort of accommodation of 1 dioptry is required, and also 1 meter angle of convergence; at 1-2 meter from the eye 2 D of accommodation is required and 2 meter angles; at 1-3 meter from the eye 3 D of accommodation and 3 meter angles, and so on; while for distant objects neither angle of convergence nor effort of accommodation is required.

The Angle Gamma.

The Optic Axis is an imaginary line (*P' P*, Fig. 7) which passes through the center (*C*) of the cornea and the posterior pole (*P*) of the globe—a point situated between the macula lutea (*M*) and the optic papilla (*D*). *The Visual Line* (*M O*) unites the point of fixation (*O*)—the object looked at—with the macula lutea; it does not coincide with the optic center (*K*) of the eye. *The Line of Fixation* (*R O*) joins the center of rotation (*R*) of the eye with the point of fixation. *The Angle* γ is the angle *O R P'* formed at the center of rotation by the optic axis and the line of fixation.

The line of fixation and the visual line so nearly coincide that in practice we regard them as identical; and hence, *in practice, the angle* γ *is the angle O K P'*. It should not be confounded, as is often the case, with *The Angle Alpha*, which is the angle *O K C'* formed at the nodal point by the visual line and the major axis (*C' K*) of the corneal ellipse. This axis rarely passes through the center of the cornea; but, as it never lies far from the latter, the difference in dimension between the two angles is very slight.

In order to measure the angle γ the eye is placed at the perimeter as for an examination of its field of vision. By means of the corneal reflection of a candle flame, which latter is moved along the arc of the perimeter, the center of the cornea is found. The position of the flame at the perimeter then gives the angle γ. The average size of the angle γ is 5°.

THE SENSE OF SIGHT.

The Sense of Sight consists of three Visual Perceptions or Sub-Senses—namely, the Light-Sense, the Color-Sense, and the Form-Sense.

The Light-Sense is the power the retina, or the visual center, has of perceiving gradations in the intensity of illumination. A convenient clinical method of testing the light-sense is the photometer invented by Messrs. Izard and Chibret. On looking through this instrument towards the sky two equally bright discs are seen. By a simple mechanism one of the discs can be made darker. If the eye does not perceive the difference in illumination between the two discs within 5° its light-sense is abnormal, or we may say its L. D. (Light Difference) is too high. Again, if one disc be made quite dark, and be then gradually lighted, the patient is required to indicate the smallest degree of light, or L. M. (Light Minimum), by which he can observe the disc issuing from the darkness. This should not be more than 1° or 2°.

Another good method is that of Bjerrum, in which the light-sense is tested by gray letters on a white ground, the letters being constructed on the same principle as Snellen's Test Types.

The Color-Sense is the power the eye has of distinguishing light of different wave-lengths. According to the Young-Helmholtz theory, the retina possesses three sets of color-perceiving elements, those for Red, Green, and Blue or Violet. These are termed primary colors, all other colors being compounds of them.

According to Hering's theory, the color-sense and the light-sense depend upon chemical changes in the retina or in the "visual substances" situated in the retina. He suggests the existence of three different visual substances, the white-black, the red-green, and the blue-yellow, by the using up or Dissimilation, and restoration or Assimilation, of which substances the sensations of light and color are produced. In the case of the white-black substance the sensation of white, or of light, corresponds to the process of dissimilation; while the sensation of black, or of darkness, corresponds to the process of assimilation. For the red-green and blue-yellow substances it cannot be said which color-sensation implies assimilation and which dissimilation. The members of the black-white pair can mingle with each other and with those of the other two pairs; but the respective members

of the two-color pairs (being "contrast colors"), *e. g.*, blue and yellow, cannot unite with each other.

In testing the color-sense the spectral colors are the best for exact experiments, but the difficulty of producing them at every moment, and of combining them, renders them of little chemical use.

The clinical method commonly employed for testing the color-sense is that of Professor Holmgren, of Upsala, which is based upon the Young-Helmholtz theory. The test-objects used are colored wools, of which a large number of skeins of every hue are thrown together.

Test I. (*vide* card on inside of end cover) consists in presenting to the individual, in good diffused daylight, a pale but pure green sample, and requiring him to select out of the bundle of wools of all colors before him all of those samples which seem to him to correspond to the test sample. If he does this correctly it is unnecessary to proceed further: the individual has normal color-sense. Amongst the skeins, however, there are some which are termed colors of confusion (grays, buffs, straw-color, etc.); and if he select one, or several, of these he is color-blind.

If, now, we desire to ascertain the kind and degree of his defect we proceed to Test II.*a*. A pink (mixture of blue and red) skein is given to be matched. If this be corectly done, we term the person incompletely color-blind; but if blue and violet, or one of them, be selected, he is red-blind (sees only the blue in the mixture of blue and red); if he select green or gray, or one of them, he is green-blind.

In order to corroborate the investigation we may employ Test II.*b*. A vivid red skein is given. The red-blind chooses, besides red, green and brown shades darker than the red; while the green-blind chooses green and brown shades lighter than the red. But I believe myself, and I think it is now very generally recognized, that red-blindness and green-blindness invariably go together. In violet (or blue) blindness purple, red, and orange will be confused in Test II.*a;* but this is an extremely rare variety of color-blindness. Total color-blindness will be recognized by a confusion of all shades having the same intensity of light, and is also rare. It is impossible by this test for any color-blind person to escape detection, except in the case of a small central color scotoma.

The individual tested should not be allowed to name the colors,

but merely to match them, as above described. The reason for this is twofold. First, because, although the congenitally color-blind person is usually unaware of his defect, yet experience has taught him which of his sensations are called blue, red, etc., by other people; and hence he can often apply the right names to colors which he really does not see as such. He is assisted in this by whatever of color-sight is left to him and by the brightness and saturation of the different colors, but is liable to frequent mistakes. Again, when the color-blind person does happen to know of his defect he is often desirous of concealing it, either because he is ashamed of it or from interested motives.*

Edridge-Green's theory [2] is that light falling upon the retina liberates the visual purple from the rods, and a photograph is formed. The decomposition of the visual purple by light chemically stimulates the ends of the cones, and a visual impulse is set up, which is conveyed through the optic nerve fibers to the brain. He assumes that the visual impulses caused by the different rays of light differ in character just as the rays of light differ in wave length. Then in the impulse itself we have the physiological basis of color. He also assumes that the quality of the impulse is perceived by a special perceptive center within the power of perceiving differences possessed by that center or portions of that center.

Edridge-Green divides the color-blind into two distinct classes, independent of each other, but which may be associated. The first class includes those who are not able to see certain rays of the spectrum; their spectrum is shortened at one or both ends. If an individual have shortening of the red end of the spectrum, he will not be able to see a red light at a distance, though he might be able to pick out all the green wools in Holmgren's Test I. A man of this kind, when shown the red light of Edridge-Green's Lantern Test, declares that there is no light visible, thus at once demonstrating his incapacity. The second class of the color-blind make mistakes, not because they cannot perceive a certain color, but because they are not able to recognize the difference between the colors, which is evident to normal-sighted persons. Normal-sighted persons see six definite colors (points of difference) in the spectrum. The second class of the color-blind see five, four, three, two, or one color, according to the degree of the

* More detailed information and Holmgren's test will be found in Appendix I.

defect; and they confuse the colors of the normal-sighted, which are included in one of their own. If the normal-sighted be designated hexachromic, those who see five colors may be called pentachromic; those who see four, tetrachromic; those who see three, trichromic; those who see two, dichromic; and the totally color-blind, monochromic.

Edridge-Green uses two tests—a Classification Test and a Lantern Test. The Classification Test consists of a number of colored wools, silks, cards, and glasses, with four test-colors—orange, violet, blue-green, and red. The examinee is asked to name the test-colors, and then to pick out from the pile all of similar color. The Lantern Test consists of a lantern and colored glasses, which can be shown alone, or combined or modified by neutral glasses. The examinee is asked to name the color of the light shown. Edridge-Green is of opinion that the use of color names is absolutely necessary, or normal-sighted persons will be rejected, through paying attention to shade rather than to color. It does not matter what name is applied to a color; but ground for rejection is afforded when the examinee calls two of the main colors of the normal-sighted, as, for instance, red and green, by the same name.

A certain proportion of people (3.5 per cent. of men and less than 1 per cent. of women) are congenitally color-blind in greater or less degree, without any diminution in the other visual functions.

Acquired color-blindness is found in toxic amblyopia, in atrophy of the optic nerve, and under some other conditions.

The Form-Sense (Acuteness of Vision) is the faculty the eye possesses of perceiving the shape or form of objects, and in clinical ophthalmology the testing of this function is an important and ever-recurring duty.

In order that an individual may have good use of his eyes it is necessary, not only that the optic nerve, retina, chorioid, and refracting media be healthy, but also that the refraction and accommodation be normal. When applied to by a patient on account of imperfect sight it is our first duty, as a rule, to ascertain accurately the condition of refraction and accommodation of his eyes. Should these be abnormal, and it be found that by aid of the correcting glasses perfect vision is obtained, we may in general conclude that the eye is organically sound, and that the patient's complaints are due to the defect in accommodation or refraction.

If the glasses do not restore perfect vision we must then, by the ophthalmoscope and other methods, decide the nature of the defect.

By Acuteness of Vision (V.) is meant the power which an eye, or rather its macula lutea, has of distinguishing form, any anomaly of its refraction, if such exist, having been first corrected, *i. e.*, while the patient wears the correcting glasses.

Now, in order to measure the acuteness of vision we must have a normal standard for comparison—*i. e.*, we must find what is the size of the smallest retinal image whose form can be distinguished. We cannot, of course, measure this image on the retina directly; but, as its size is proportional to the visual angle—the angle which the object subtends at the eye—it is sufficient to determine the smallest visual angle under which the form of an object can




Fig. 8.

be distinguished. It has been found, experimentally, that the average size of this angle is 5′.

In order practically to ascertain the degree of acuteness of vision, we place our patient with his back to the light, while facing him at a distance of 6 meters, and in good light, are placed Snellen's Test-Types for distance. These types are so designed that at the distance at which they should be seen they each subtend an angle of 5′ at the eye. The largest type should be seen at 60 meters (Fig. 8) by the normal eye, and the types range from this down to a size visible not farther off than 6 meters. If V = Acuteness of Vision, d = the distance from the eye to be tested to the test-types, and D = the distance at which the type should be distinguishable, then $V = \frac{d}{D}$. For example: if d = 6 meters (a distance which most rooms can command), and if the eye see type D = 6, then $V = \frac{6}{6} = 1$, or normal V.; but if at 6 meters the eye see only D = 60, which should be seen at 60 meters, then $V = \frac{6}{60}$, or very imperfect vision.

Should the patient's sight be so bad that he is unable to read any of the letters, it may be tested by finding at what distance he can count the surgeon's fingers; and if he cannot even do that, then his power of perception of light (his P. L.) should be tested. This is done by means of a lamp in a dark room, the eye being alternately covered and uncovered, and the patient being required to say when it is "light" and when "dark." If the flame be gradually lowered the smallest degree of illumination perceptible will be ascertained.

The eyes must be examined separately, that one not under examination being excluded from vision by being shaded with the patient's own hand or other suitable screen; but it must not be at all pressed on, as any pressure would dim its vision when its turn for examination may come.

With the advance of age the acuteness of vision undergoes a slight but steady reduction, owing to certain senile changes in the eye.

THE FIELD OF VISION.

By the Field of Vision (F. V.) is meant the space within which, when one eye is closed, objects can be seen by its fellow, the gaze of the latter being fixed the while on some one object or point. Thus if, standing on a hill, we fix the gaze of one eye on some object on the plain below, the field of vision includes not only that object, but many others also for miles around it. If the fixation object be nearer to us, the area taken in by our field of vision will be proportionately diminished in extent.

The fixation object is seen by central or direct vision, its image being formed on the macula lutea; the other objects in the field of vision correspond with as many different points in the more peripheral parts of the retina, and are seen by eccentric, or indirect, vision. Eccentric vision is of great importance for the guiding of ourselves and avoiding obstacles in our way. Its use may be realized by the experiment of looking through a long small-bore cylinder (*e. g.*, a roll of music) with one eye, thus cutting off its eccentric field, while the other eye is closed.

The Dimensions of the Field of Vision may be measured for clinical purposes by means of an instrument called the perimeter. This is a semicircular metal band, which revolves upon its middle point, being in this way capable of describing a hemisphere in

space. The arc is divided into degrees marked on it, from 0° placed at its middle point, to 90° at either extremity. At the center of the hemisphere is situated the eye under examination, while the fixation point is placed exactly opposite, in the center of the semicircle, at 0°. A small bit of white paper 5 mm. square, the test object, is slowly moved along the inner surface of the arc from the periphery towards the center, until it comes into view, and




Fig. 9.—Chart of F. V. of Right Eye.

this point is observed in various meridians. The horizontal, vertical, and two intermediate meridians, at the least, should be examined by placing the arc of the perimeter in the corresponding planes.

The boundary of the field may be noted on a diagram or chart (Fig. 9), which represents the projection of a sphere on a plane surface. The radii represent different meridians, which may be determined by a dial with pointer on the back of the perimeter,

while the concentric circles correspond with the degrees marked on the arc. A pencil mark is placed on the chart at the spot corresponding to that on the perimeter at which the test object comes into view, and when the different meridians have been examined these marks are united by a continuous line, which then represents the outer boundary of the F. V.

The normal F. V. is not circular, but extends outwards about 95°, upwards about 53°, inwards about 47°, and downwards about 65°, as represented by the strong curve in Fig. 9. The limitation upwards and inwards is partly due to the projection of the supra-




Fig. 10.

orbital margin and the bridge of the nose, but also to the fact that the outer and lower parts of the retina are less practiced in seeing than are the upper and inner parts, and their functions consequently less developed. The acuteness of vision diminishes progressively towards the periphery of the field, two points of a certain size close together being distinguishable from each other only a short distance from the fixation point, while the farther towards the periphery the larger must be the test objects.

Fig. 10 serves to illustrate the projection of the field of vision on the semicircle of the perimeter to its extreme temporal (95°)

and its extreme nasal (47°) boundaries, as well as the portion of the retina (*a* to *b*) which corresponds to this extent of field; and it shows that the sensitive portion of the retina, or rather perhaps the portion of the retina which is most used, extends farther forward on the nasal than on the temporal side. The diagram also




FIG. 11 (*Landolt*).—Chart of F. V. of Left Eye.

explains the remarkable fact that the field extends in the temporal direction more than 90°.

It must be remembered, too, that the fields of vision overlap, as the two visual axes meet at the fixation point. Fig. 12 represents the binocular portion white, P. being the fixation point. The shaded portion on the right belongs to the right eye alone, while that on the left belongs to the left eye alone.

The Blind Spot of Mariotte is a small blind island or scotoma in the F. V., situated about 15° to the outer side of the point of

fixation and just below the horizontal meridian. It is shown as a white spot in Fig. 12. It is due to the optic papilla (optic disc), for at that place the outer layers of the retina are wanting, and hence there is there no power of perception. There are also, occasionally, minute blind spots in the field, due to large retinal vessels, which interfere with the formation of the image upon the layer of rods and cones.

The Perception of Colors in the Periphery of the Field can be examined with the perimeter, by means of bits of colored paper 5 mm. square. It has been in this way ascertained that the boundaries of the power of eccentric perception for the different colors do not seem to correspond with the boundary for white light, nor do the boundaries of the different colors seem to coincide. Ex-




FIG. 12.—Binocular Field of Vision.

amining from the periphery towards the center by ordinary daylight, blue is the color which can be distinguished as such most eccentrically, its field extending nearly as far as the general F. V.; then come yellow, orange, red, and, with the most limited field, green. Blue, red, and green being the most important, their fields are noted in Fig. 11. Although the respective colors are distinguishable within the limits indicated, they are by no means so brilliant in hue as when seen by direct vision. It has, however, been demonstrated that every color is recognizable up to the outer limit of the F. V., if the colored object be of sufficient surface and be sufficiently illuminated; so that there is, in fact, no absolute color-blindness in the peripheral parts of the retina, but merely a diminished sensitiveness to colored light.

The Perception of Form in the Periphery of the Field is very defective, and its examination is not of much practical importance; but this portion of the field is very sensitive to the movement of objects.

References.

[1] "Optique Physiologique," pp. 158-167.
[2] "Color Blindness and Color Perception: International Scientific Series."
[3] "Von Graefe's Archiv," xxxix. 2. p. 71.

CHAPTER II.

ABNORMAL REFRACTION AND ACCOMMODATION.

I HAVE explained what is meant by Normal Refraction, or Emmetropia. We recognize three different forms of Abnormal Refraction, or Ametropia (*α, priv.; μέτρον, standard; ὤψ*). 1. Hypermetropia (*ὕπερ, over; μέτρον, standard; ὤψ*), in which the principal focus of parallel rays of light lies behind the retina. 2. Myopia (*μύειν, to close; ὤψ*), or Short-sight, in which the principal focus of such rays lies in front of the retina. 3. Astigmatism (*α, priv.; στίγμα, a point*), in which the refraction of the eye in its different meridians is different.

HYPERMETROPIA.

In a large proportion of cases this form of Ametropia is due to the eyeball being too short in its antero-posterior axis (Axial H.).




FIG. 13.

It may also depend upon deficient refracting power in the dioptric media (Curvature H.).

Parallel rays of light (*a b*, Fig. 13) falling into the hypermetropic eye (*E*) do not meet on the retina, but converge towards a point (*c*) situated behind it. Consequently these rays do not form on the retina a distinct image of the object looked at, but produce there a circle of diffusion (*d e*), or blurred representation of the object.

Since, therefore, in hypermetropia the retina is in front of the principal focus of the dioptric system, rays passing out of the eye from any point (R, Fig. 14) on this retina will pass out as divergent rays (f g), and will appear to come from a point (R') situ-




Fig. 14.

ated behind the eye, which point is the virtual conjugate focus of the point R.

Now in order to correct the hypermetropia—so that parallel rays (a b, Fig. 15) passing into it may be brought to a focus on the retina—a convex lens (L) must be placed in front of the eye, of sufficient strength to render the parallel rays, before they enter




Fig. 15.

the eye, convergent towards R', so that when they meet the eye they may be brought to a focus on the retina R, which is the conjugate focus of R'. The higher the hypermetropia—*i. e.*, the shorter the antero-posterior axis of the eyeball—the stronger must the correcting glass be. It may be found that, with a lens of some dioptries less power, the eye will see equally well; but this it does by means of an effort of accommodation, which supplements the inadequate refracting power of the lens placed before it. As

we proceed to higher lenses the effort of accommodation is relaxed, until, finally, the strongest lens with which vision is still at its best is reached when, it may for the present be assumed, no further effort of accommodation is made, and *L* represents the whole error of refraction.

In low degrees of hypermetropia the eye can frequently see distant objects distinctly by an effort of accommodation, which completely takes the place of *L*. When such an eye is found to have full vision without a glass, a beginner may fall into the error of regarding it as emmetropic; but if he will take the precaution of placing a low convex lens in front of it, and then finds that the acuteness of vision remains as good as without the glass (because the effort of accommodation is now relaxed), he will avoid this mistake, unless there should be tonic cramp of accommodation, which might partially, or even completely, mask the hypermetropia.

If a glass a single number higher than the exact measure of the defect be placed before the eye, vision again becomes indistinct, because the rays are then brought to a focus in front of the retina, and a circle of diffusion is formed on the latter. The eye, in fact, is put by such a glass in a condition of myopia. Therefore *the strongest convex glass with which a hypermetropic eye can see distant objects (the test-types) most distinctly is the glass which corrects its hypermetropia, and is the measure of the latter.* Very commonly it is only the manifest hypermetropia (*vide infra*) which is ascertained by this method, unless the accommodation has been previously paralyzed by atropin.

This method of determining the refraction by means of the trial-lenses and test-types is not relied on now-a-days by ophthalmic surgeons to the same extent as formerly, the examination of the upright ophthalmoscopic image, or else retinoscopy, the use of which will be explained later on, having largely taken its place. In conjunction with these it is a valuable method.

The degree of the hypermetropia is indicated, as has been said, by the number of the lens which corrects it.* Thus, if the number of the glass (*L*, Fig. 15) required to correct the hypermetropia

* Theoretically the glass which measures the error of refraction should be in contact with the eye, but for practical purposes the distance between the glass and the eye may be neglected, especially if the glasses are worn at the same distance from the eye as they occupied during the testing.

of the eye (E) be 2.0 D, we say this eye is hypermetropic two dioptries, or has a hypermetropia of two dioptries, or we would write it down H = 2.0 D.

Amplitude of Accommodation in Hypermetropia.—When at rest the refraction of the hypermetropic eye is deficient: consequently r must be negative ($-r$), and the amplitude of accommodation must include the power required to adapt the eye to infinity; therefore

$$a = p - (-r) = p + r.$$

For example: if the punctum proximum of a hypermetropic eye of 5 D be at 30 cm., what is the amplitude of accommodation? 5 D ($=r$) is necessary in order to make the eye emmetropic, and to accommodate the emmetropic eye to 30 cm. 3.25 D ($\frac{100}{30} = 3.25$) is required. Hence $a = 3.25 + 5 = 8.25$ D.

The Angle γ in Hypermetropia.—In hypermetropia, as in emmetropia, the cornea is cut to the inside of its axis by the visual line; but in hypermetropia the angle which the visual line forms with the axis of the cornea




Fig. 16 (*right eye*).

is very much greater, owing to the shortness of the eyeball, the effect of which is to increase the distance between the macula lutea (M) and the optic axis (A, Fig 16). Consequently, in extreme cases, when the two visual lines of a hypermetropic individual are directed to an object, the axes of the corneæ may seem to diverge, and thus the appearance of a divergent strabismus will be given (apparent strabismus, see chap. xviii.).

The eyes of animals and of uncivilized nations are hypermetropic; children, too, are hypermetropic at birth, but as they grow older the refraction increases, and they become less hypermetropic, or emmetropic, or even myopic.

The evil effects of the constant and excessive demand upon the accommodation in hypermetropia are chiefly these:

1. **Cramp of the Ciliary Muscle.**— Its persistently maintained contraction frequently gives rise to a tonic cramp of the muscle. This spasm is not, or may be only partially, relaxed when the correcting convex glass is held before the eye, and consequently the

whole or part of the hypermetropia may be masked by the cramp. That part of the hypermetropia which is thus masked is called latent (Hl.), while the part which is revealed by the convex glass is called manifest (Hm.). The entire hypermetropia is made up of the latent and manifest H. (H = Hm.+ Hl.).

If the cramp be excessive, parallel rays may be kept convergent on the retina by it alone, and vision then would be made worse, rather than better, by even a weak convex glass held before the eye, a circumstance which might lead the surgeon to think he had to do with an emmetropic eye. In this case we say that the whole hypermetropia is latent.

Or, in extreme cases of accommodative spasm, parallel rays may be united in front of the retina, and the eye made apparently myopic, the vision being capable of improvement by concave glasses. Serious errors might therefore arise if this cramp were overlooked, as it is very apt to be in the examination with the trial-lenses. When it is present in a high degree, the patient cannot maintain a sustained view of an object at any distance without suffering pain in and about the eyes. It is frequently the reason why perfect acuteness of vision is not obtained by aid of the trial-lenses, and the surgeon must be careful not to be led into an error of diagnosis by it. Examination with the ophthalmoscope, or paralysis of accommodation with atropin, will enable him to avoid mistakes.

In order to relieve this cramp, the ciliary muscle must be paralyzed by a solution of atropin freely instilled; and it will often be necessary to keep the accommodation paralyzed for some days, and to commence the use of the correcting spectacles before the effect of the atropin begins to wear off. In this way a recurrence of the spasm may be often prevented.

As life advances, and the power of accommodation diminishes, the manifest part of the hypermetropia increases, while the latent part decreases, until finally Hm. = H.

2. **Accommodative Asthenopia** (*ἀ, priv.; σθένος, strength; ὤψ*).—In looking at distant objects the accommodation of the emmetropic eye is at perfect rest, and does not come into play until the object is approached close (within 6 m.) to the eye. But even for distant objects the hypermetropic eye must accommodate; and, having for those distances used up part of its accommodative energy, it has for near objects actually less at disposition than the normal eye. Hence we find that hypermetropic people

often complain of inability to sustain accommodative efforts for near objects for any length of time. After reading, sewing, etc., for a short time, sensations of pressure in the eyes and of weight above and around them come on, and the words or stitches become indistinct, and cannot be distinguished. The work must then be interrupted, and after a few minutes' rest it can be resumed, but must soon again be given up. After a Sunday's rest the patient is often able to get on better than on the previous Saturday. These symptoms depend simply upon inability of the ciliary muscle to perform the excessive demands made upon it.

Accommodative Asthenopia often appears suddenly during or after illness, the explanation being that, although hypermetropia had always existed, yet in health the ciliary muscle was equal to the great efforts required of it, but in sickness it shared the debility of the system in general.

3. **Internal, or Convergent Concomitant Strabismus.**—This condition has a certain relation to hypermetropia. It will be treated of in the chapter on the Motions of the Eyeballs and their Derangements (chap xvii.).

The Prescribing of Spectacles in Hypermetropia.—If a person be found to be hypermetropic, but his acuteness of vision without glasses be good, or as good as he desires, and he complain of no asthenopic symptoms, glasses need not, indeed should not, be prescribed for him. No disease in his eye will result from his going without glasses.

If the patient complain of imperfect distant vision due to hypermetropia, then those lenses which correct the Hm. may be prescribed for distant vision, to be worn either constantly or occasionally, as he may desire. Such a patient is almost certain to complain also of accommodative asthenopia; while many patients will be met with who complain of the latter, yet express themselves as perfectly satisfied with their distant vision. For relief of the asthenopia it is usually enough to prescribe spectacles for near work which will correct the Hm., along with 1 D or 2 D of the Hl., if the latter exist.

If there be excessive cramp of accommodation, glasses to correct the whole hypermetropia should be worn while the eye is under atropin; and afterwards as much of the Hl. as possible, along with the Hm., should be corrected by glasses to be worn constantly.

Myopia, or Short-Sight.

This form of ametropia is due, in a vast majority of cases, to the antero-posterior axis of the eyeball being too long (Axial M.), and hence, its refracting media not being proportionately diminished in power, parallel rays of light (*a b*, Fig. 17) are not brought




Fig. 17.

to a focus on the retina, but in front of it (at *f*) and form on the retina circles of diffusion (*c d*).

Myopia may also be caused by abnormally high refracting power in the crystalline lens, as in spasm of the ciliary muscle, and in some cases of commencing cataract, and also by conical cornea (Curvature M.).

Since, in the myopic eye, the retina is beyond the principal focus




Fig. 18.

of the dioptric system, rays emerging from any point (*c*, Fig. 18) of the fundus will pass out convergently, and will unite in front of the eye at the conjugate focus of the retina (*r*).

Conversely, rays diverging from a certain point (*r*, Fig. 18) in front of the eye will be focused on the retina (*c*).

If an object be brought towards the eye, the divergence of those rays which pass from it into the eye increases until, when it has reached the point *r*, their divergence is just sufficient to allow them to be united at the conjugate focus (*c*), which is on the retina. This point *r* is the punctum remotum* of the myopic eye. In order, therefore, that the short-sighted eye may be able to see distant objects, it is necessary that the parallel rays (*a b*, Fig. 19) coming from those objects should be given such a degree of divergence before they pass into the eye as though they came from this punctum remotum. This can readily be effected by placing the suitable concave lens (L) in front of the eye, and the number of this glass will indicate the degree of the myopia—*i. e.*, by how many dioptries the refracting power of the eye is in excess of that of an emmetropic eye. The focal length of the correcting glass corresponds, of course, with the distance of the punctum remotum




Fig. 19.

(*r*) from the eye, provided the glass be held close to the cornea. The focus of the glass and the punctum remotum of the eye are then identical, and, therefore, parallel rays, after passing through the glass, will have a divergence as though they came from the punctum remotum, and will form an exact image of the object from which they come on the retina.

For example: if the punctum remotum (Fig. 19) be situated at 14 cm. from the eye, then the number of the correcting lens

* The punctum remotum is always the conjugate focus of the retina. In an emmetropic eye it is at infinity, since the retina is at the principal focus of the eye, and the rays pass out parallel. In hypermetropia it is behind the eye, and is virtual or negative, because the retina is in front of the principal focus, and the rays pass out divergently, as if coming from a point behind the retina. Lastly, in myopia it is situated at a finite distance in front of the eye, and is real and positive, because the retina is beyond the principal focus, and the rays emerge convergently.

will be 7 D, because the focal distance of this lens is 14 cm. ($\frac{100}{14} = 7$). In practice, however, we cannot hold the glass so close to the cornea, and, therefore, we must subtract the distance between it and the cornea from the focal distance of the required lens. In the above case, suppose the distance from cornea to glass be 4 cm., the required lens in practice will be 10 D ($\frac{100}{10} = 10$).

Determination of the Degree of Myopia.—The degree, or amount, of myopia, as of hypermetropia, may be determined either by the ophthalmoscope, or experimentally by means of the trial-lenses and test-types.

By the latter method, examining each eye separately, we find the correcting glass by placing our patient as directed in the section on Acuteness of Vision (p. 29). A weak concave trial-glass is then held before the eye under examination, and higher numbers are




FIG. 20.

gradually proceeded to until the glass is reached which gives the eye the best distinguishing power for the types. We often find that there are several glasses, with each of which the patient can see equally well. *The weakest of these is the measure of his myopia.* When a higher glass is used the eye may still see well, but it does so only by an effort of accommodation (*i. e.*, the crystalline lens has to be made more convex in order to compensate for the excessive concavity of the glass placed in front of the eye), and the glass employed represents then not merely the myopia present, but also this accommodative effort. No more serious mistake can be made than the prescribing of too strong concave glasses for a myopic individual, as will be seen further on.

The Amplitude of Accommodation in Myopia.—The myopic eye has an excess of refractive power as compared with the emmetropic eye; therefore, in calculating its amplitude of accommodation, this excess must be

subtracted from the positive refractive power (p) which would be required to adapt the emmetropic eye to the same punctum proximum; or, in other words, the myopic eye has need of less accommodative power than the emmetropic eye, because, even at rest, it is adapted for a distance (R., its punctum remotum) for which the emmetropic eye has to accommodate; hence in myopia

$$a = p - r.$$

For example: a myopic person of 10 D who can accommodate up to 8 cm. ($p = \frac{100}{8} = 12$ D) has an amplitude of accommodation of $12 - 10 = 2$ D.

The Angle γ in Myopia.—In myopia, owing to the length of the eyeball, the cornea is cut much closer to its center by the visual line than in emmetropia, or the visual line (M V, Fig. 21) and the optic axis (A O) may coincide, or the cornea may even be cut to the outside of its center by the visual line (*vide* Fig. 20). In any of these cases, but especially in the latter, the effect will be that of an apparent convergent strabismus.

Myopia is rarely congenital. It generally first shows itself from the eighth to the tenth year, and is apt to increase, especially during the early years of puberty. Its progressive increase is encouraged by use of the eye for near work, such as reading, sewing, drawing, etc., and is due to a further elongation of the antero-posterior optic axis. But it is certain that in addition to this exciting cause there must be some predisposing condition or conditions, as but few children become short-sighted, although all are educated in a very similar manner in so far as the use of their eyes is concerned. Moreover, high degrees of myopia are occasionally met with in young children before they have begun to use the eyes much for close work. Stilling [1] and Seggel [2] have found that a low orbit is usually associated with a myopic formation of eyeball, and they are inclined to regard these largely in the light of cause and effect. For with a low orbit, and when, as often happens, the tendon of the superior oblique has an almost transverse direction, the combined pressure of the two obliques upon the plane of the equator during the period of growth would tend to cause elongation of the antero-posterior diameter of the eyeball. Opposed to this view is the fact that in scholars of Esthic nationality, who have broad faces and low orbits, the proportion of myopes is less than in Europeans. There are many other theories as to the causation of short-sight. Certain it is that the tendency to myopia is often hereditary, and is frequently seen in several members of a family; but the whole question of the predisposing causes of myopia must still be regarded as *sub judice*.

In cases of commencing cataract a slight degree of myopia may

sometimes be noticed to come on. This is due to a higher refracting power in the lens, as the result of the changes beginning in it.

Many short-sighted people half close their eyes when endeavoring to distinguish distant objects, in order that the rays may be prevented, so far as possible, from passing through peripheral parts of the crystalline lens, which would increase the circles of diffusion. This habit it is which has given the name of myopia to the condition (*vide* p. 36).

Progressive Myopia frequently becomes complicated with Organic Disease, and to the more serious cases the term **Pernicious Myopia** may be applied. The following are the forms of organic disease met with:

1. *Posterior Staphyloma, or Myopic Crescent.*—This condition is recognized by the ophthalmoscope as a more or less extensive white crescent at the outer side of the optic papilla. Owing to bulging of the eyeball, the chorioid becomes atrophied at this place, and a crescentic portion of the white sclerotic is laid bare. The staphyloma sometimes extends all round the optic papilla, and by stretching of the retina in extreme cases its functions may become deranged, and the blind spot increased in size. But every case in which a small crescent is present is not to be regarded as having a serious bearing. Much here depends on the age of the patient and the degree of the myopia. The younger the patient and the higher the myopia, the more serious is the outlook. As a matter of experience in the majority of cases the short-sight increases by only a very few dioptries, and does not become complicated with the more dangerous forms of organic disease. Infintely more serious than Posterior Staphyloma are the following conditions:

2. *Chorioidal Degeneration in the Neighborhood of the Macula Lutea.*—This should always be carefully looked for, as the region of the yellow spot is very liable to disease in the worst cases of progressive myopia. The disease seems to begin in the chorioid, giving the appearance of small cracks or fissures, which later on develop into a patch of chorioidal atrophy. The retina at the spot becomes disorganized, and very serious disturbance of vision, associated in the early stages with metamorphopsia, is the result, the patient being disabled from reading, although, as the periphery of the fundus is usually sound, he can find his way about freely. Treatment can do little here. Abstention from use for near work and the wearing of dark glasses are to be recommended.

3. *Chorioidal Exudation in the Neighborhood of the Macula Lutea.*—A small gray spot of exudation may appear in the chorioid at this place, accompanied by loss of sight for reading. These cases are often amenable to active mercurial treatment, when sight may be restored. Should the case be neglected or run a bad course, vision will be permanently damaged from secondary chorioidal degeneration.

4. *The Black Spot in Myopia.*—This disease also attacks the chorioid in the region of the yellow spot, and causes a loss of central vision similarly as in the two previous forms of disease. The appearance shown by the ophthalmoscope is that of a black spot, usually quite circular and with a defined margin. In the early stages its size is much smaller than that of the papilla, but later it often attains a dimension of two papilla diameters, or more. The spot is rarely of an equal intensity of blackness all over, but towards its center a faint reddish hue often shines through in places. At a later stage the black spot becomes surrounded by a narrow whitish border, while towards its center it becomes less black, and finally grayish or even white, its margin remaining black. Although small hemorrhages, which often occur in the neighborhood of the black spot, gave rise to the opinion that the black spot itself was the result of hemorrhage, yet this seems not to be so, as the investigations of E. Lehmus[3] have shown. The diseased changes consist in a proliferation of the pigment epithelium, combined with a gelatinous exudation, which in the case examined had attained a thickness, at the center of the black spot, of two-thirds that of the chorioid. The chorioid was but very slightly altered, and the glass membrane was quite normal. At the margin of the proliferating region the pigment epithelium was found to be paler or even quite free of pigment. The black spot very gradually, in the course of years, attains its ultimate dimension, and then very slowly retrogresses, until finally its place is taken by a grayish or bluish-white scar. Treatment is of no avail, and central vision does not becomes restored.

5. *General Chorioidal Atrophy.*—In advanced cases of pernicious myopia, large patches of chorioidal atrophy, other than the crescent, are often present, chiefly in the region of the posterior pole, but very liable to extend also towards the periphery of the fundus. The vitreous humor in these cases usually contains many opacities. Treatment by means of subconjunctival saline injections is occasionally of use in clearing up the vitreous humor,

and thus effecting some improvement of vision. The eyes should not be used for near work, and dark glasses should be worn.

6. *Hemorrhage in the Retina at the Yellow Spot* may occur, and when the hemorrhage becomes absorbed the macula lutea may not recover its function, owing to the delicate retinal tissue having been seriously damaged. Yet we often meet with cases of this kind which do regain their former vision. Rest of the eyes and dark glasses should be prescribed.

7. *Detachment of the Retina.*—This is a frequent and most serious complication of progressive myopia. It will be considered in the chapter on Diseases of the Retina (chap. xv.).

Functional Anomalies attending Progressive Myopia.—

(*a*) *Insufficiency of the Internal Recti Muscles* is almost always associated with progressive myopia, but it can hardly be regarded as a result of the latter. It may more properly be looked upon as a concomitant congenital irregularity, and perhaps as one of the causes of the progressive nature of myopia; yet it is indeed very commonly present in cases of slight myopia, which have no tendency to increase. It will be fully discussed in chap. xvii.

(*b*) *Cramp of Accommodation* is often present in myopic eyes, and will cause the myopia, examined with trial-lenses and test-types, to seem higher than it really is. The surgeon, being aware of this source of error, will guard against it.

The Management of Myopia.—In view of the tendency to become more short-sighted, to which, especially during adolescence, nearly every case of short-sight is liable, and of the fact that in a given case we cannot tell to what extent this increase may go—whether to slight, medium, or high myopia—and, finally, as high degrees of myopia almost invariably lead to disease of the eye, the management of myopia, including the prescribing of glasses for it, is one of the most important matters with which we have to deal.

The Prescribing of Glasses in Myopia.—It is not necessary to prescribe glasses for very slight degrees of myopia (up to 1.0 D or 1.5 D); yet should the patient desire to wear correcting glasses for distant objects, there can be no objection to his doing so. But for cases of myopia of 2.0 D or more, it becomes very desirable to prescribe glasses which fully correct the myopia, to be worn constantly—*i. e.*, for both distant and near objects; and, should the

myopia increase, to increase from time to time the strength of the glasses accordingly.

This has not always been the teaching on this point, and even yet many ophthalmologists do not prescribe fully correcting glasses for their myopic patients, especially for near work. It has been held that any effort of accommodation which might be caused by fully correcting concave glasses would promote the tendency to increase of the short-sight, and to chorioidal disease. In fact, it was regarded as important to save the myopic eye every effort of accommodation in so far as possible. The usual practice was to order a glass 0.5 D, 1 D or 2 D less than the full correction for distant vision; while, for near work, either no glasses were allowed, if the myopia was of low degree; or, if of medium or high degree, a separate pair of spectacles, of such strength as would merely enable the patient to read ordinary print at about 30 cm. from his eye, was given. It was also the practice to restrict the use of the myopic eye for near work to a limited number of hours daily, with intervals of rest, or even to interdict all near work absolutely for a period of months at a time. The main object of the reading-glass was—the patient being enabled thereby to hold near objects farther from his eye—to diminish the angle of convergence of the optic axes, and thus to reduce the pressure of the orbital muscles on the eyeball during convergence, a pressure which it is considered promotes the elongation of the globe.

We now know, on the one hand, that the action of the muscle of accommodation does not produce a pull on the chorioid farther back than the equator of the eyeball, while it is, of course, at the posterior pole that the diseased processes in myopia commence. Nor does the ciliary muscle by raising the tension of the eye, nor in any other way, cause an elongation of the eyeball. Hence, there is no reason to spare the healthy myopic eye any ordinary effort of accommodation. Indeed, it is reasonable to think that if normal efforts be required of the ciliary muscle, its more healthy tone will improve the general healthy nutrition of the uveal tract, and consequently will tend rather to avert morbid changes in it.

On the other hand, the diminution of the angle of convergence at near work is a truly important matter, for the reason above stated; but it is more effectually provided for by full than by partial correction.

Practical experience is here even more valuable than theory, and it shows that in a large majority of those patients whose short-

sight has been fully corrected in youth, and who have worn their spectacles contantly for a number of years, the myopia in many instances has not increased at all, while in a large proportion the increase will have been moderate, and in but a small proportion marked pernicious progress will be noted. In short, the tendency to increase of the myopia and to organic disease is less than in those myopes who have either worn no glasses or only partially correcting glasses.

Well-fitting, properly centered spectacles are much to be preferred to folders, which are difficult to keep correctly centered before the eyes. Any astigmatism present should always be corrected. Patients whose eyes are healthy, and who wear constant full correction, may be permitted, and even encouraged, to use their eyes freely for near work, always keeping the work as far from the eyes as possible, to diminish the angle of convergence. With this latter object in view, too, well-printed books, ample light, and suitable reading- and writing-desks should be provided in all educational establishments, and for home studies.

But in prescribing the full correction for constant wear to young short-sighted persons, we meet with some difficulties. The first of these is due to the range of accommodation, which is imperfect in the myopic eye, and consequently the patients may complain of painful accommodative sensations when first using their fully correcting lenses for near work, and sometimes they decline to persist in the attempt. These complaints are more likely to be made by patients of about twenty years of age or more, whose habit of use of their eyes (relative amplitude of accommodation, and degree of convergence) has become more or less confirmed, and in whom the power of accommodation has naturally diminished to an appreciable degree, rather than by persons of from twelve to seventeen years of age. Patients should be encouraged, in spite of discomfort, to continue for some time longer to read, etc., with the full correction, when, very often, the relative amplitude of accommodation will gradually improve, and the discomfort will cease. Or, a lower number than the full correction may be ordered, and the strength gradually increased, until, in the course of some weeks or longer, the full correction can be worn for near work with ease.

A second obstacle to full correction for constant wear is the vanity of the patients, or of their parents, who for the sake of appearance dislike the spectacles. Again, the rooted and widespread prejudice against "strong glasses," irrespective of the question

whether those glasses correspond to the degree of the myopia of the eyes they are prescribed for or not, will often stand in the way of proper advice on this subject being followed.

Operative Cure of Myopia.— This consists in the removal of the crystalline lens, and was first systematically employed by Fukala.[4] Some surgeons simply extract the clear lens by one of the methods used for cataract, while the majority first perform discission of the lens, and, when it has swollen and become cataractous, they proceed to evacuate it through a linear corneal incision. Discission pure and simple is, I think, preferable to either of these methods, the process of absorption being allowed to go on until the whole lens gradually disappears. The opaque lens matter is not evacuated through a corneal incision, unless increased tension or other complications should make it necessary. I believe that discission pure and simple is not only the safest method as an operative proceeding, but is also less likely to be followed by any intra-ocular complication, such as detachment of the retina, or hemorrhage, than where a sudden reduction in the contents of the eyeball is effected, as by either of the other methods. Moreover, the disfigurement caused by the corneal incision, slight though it be, is a drawback from which the simple discission is free. Nor, again, is astigmatism so apt to be produced. The cure, by simple discission, has the disadvantage of being very slow, but this is not a serious drawback in view of the lifelong advantages to be gained.

It is important, whatever the method used may be, to avoid injury of the vitreous humor, as any such injury may lead to shrinking of the vitreous, followed by detachment of the retina. Whether in these highly myopic eyes their tendency to retinal detachment is increased by the operation, is a question which has been answered by means of statistics, in the sense that the operated cases are neither more nor less liable to detachment than the unoperated cases.

The operative cure of myopia is not to be recommended except for cases of 12.0 D and more; nor should it be performed where there is such serious disease of the fundus or vitreous humor as would render any improved use of the eye on conclusion of the treatment unlikely. Active chorioidal disease is regarded as a contra-indication, but small retinal hemorrhages, even if they be near the macula lutea, need not be so regarded. The best time of life for the cure is in childhood or early youth, but it can be suc-

cessfully undertaken at a much later period. In the myopic eye the nucleus of the lens undergoes sclerosis to a less extent than in hypermetropia or in emmetropia, and hence in it discission is less apt to be followed by high tension or other complication, even when performed in middle age.

The advantages gained by the patient from the operative cure of his myopia are enormous. Not merely does the patient become either only slightly myopic, or else emmetropic or hypermetropic, according to the degree of the original myopia, but his acuteness of vision is usually increased in a remarkable degree. One of my patients, with M. 20 D and V. = Finger-counting at 2 m., obtained after discission and resulting absorption of the lens an emmetropic eye, and V. = $\frac{6}{18}$. The cause of this improved acuteness of vision is not clearly understood, and in some cases it goes on for several months before it reaches its height. Another remarkable point, illustrated, too, by the above case, is that the amount of reduction in the refraction of the eye is much greater in these cases than one would *a priori* expect. A convex lens of about 10.0 D is commonly required to correct an emmetropic eye that has been operated on for cataract, which would imply that the refracting power of the crystalline lens is about + 10.0 D in the emmetropic eye. But, after the operative cure of myopia, it is found that the reduction in the refraction of the eye amounts to from 12.0 D to 20.0 D, and varies in different eyes. Convex spectacles require to be worn for near work after the operation, and sometimes, too, for distant vision where hypermetropia has been produced. Where the original myopia is not more than about 12.0 D it is advisable, in patients who follow certain callings in which the wearing of glasses is objected to, to operate on only one eye, which then subsequently serves for distant vision, while the unoperated eye is used for reading and other near work, and by this means the patient is entirely independent of glasses.

Astigmatism.

This is a compound form of ametropia, due to the cornea being more curved in one meridian than in another, similarly as the back of the bowl of a spoon is more convex from side to side than from heel to point.

In Regular Astigmatism the directions of the greatest and least curvations of the cornea are always at right angles to each other,

and usually fall precisely in the vertical and horizontal meridians, the meridian of greatest curvature being most frequently the vertical. Consequently, we say the astigmatism is "with the rule" in those cases in which the meridian of greatest curvature is the vertical; and, where that meridian is the one of least curvature, we say the astigmatism is "against the rule." The result of this is that a pencil of rays passing into the eye, instead of meeting at a common focus, is irregularly refracted, those rays passing through the vertical meridian of the cornea being brought to a focus much earlier than those which fall through its horizontal meridian; and therefore at the focus of the former the latter rays form a horizontal streak of light. The intermediate, or oblique, meridians will probably be of regularly intermediate refracting power.

The interval between the foci of the two principal meridians is called the Focal Interval, and is a measure of the astigmatism.




Fig. 21.

The accompanying diagram (Fig. 21), after Donders, will assist in the understanding of the course of a pencil of rays after they have passed through an astigmatic cornea, those rays belonging to the horizontal and vertical meridians being chiefly considered.

At A neither vertical ($v\ v'$) nor horizontal ($h\ h'$) rays have yet been united at their foci, but the vertical rays are the nearest to their focus; and therefore the appearance which the pencil of rays would give, if caught here on an intercepting screen, is an oval with its long axis horizontal, as shown by the dotted line. At B the vertical rays have met at their focus, but the horizontal rays not as yet at theirs, and the result is therefore a horizontal straight line. At C the vertical rays are diverging again from their focus, and the horizontal rays have still not come to theirs. At D the same conditions exist, only a little farther on, where the one set of rays is diverging, the other still converging, but each at the same angle; hence the shape of the figure is round. At F the horizontal

rays have met, and the result is a vertical straight line. At G both sets of rays are divergent, and the figure is an oval with the long axis perpendicular.

There are various kinds of regular astigmatism, according to the position of the two principal foci with reference to the retina, as follows:

1. *Simple Hypermetropic Astigmatism.*—When the focus (V, Fig. 22) of the vertical rays is situated on the retina (emmetropia




Fig. 22.




Fig. 23.

in that meridian), while that (H) of the horizontal rays lies behind the retina (hypermetropia in that meridian).

2. *Compound Hypermetropic Astigmatism.*—When the foci of both sets of rays is behind the retina, that (H, Fig. 23) of the horizontal rays is farther back than that (V) of the vertical rays.




Fig. 24.




Fig. 25.

3. *Simple Myopic Astigmatism.*—When the focus (H, Fig. 24) of the horizontal rays is situated on the retina (emmetropia in that meridian), while the focus (V) of the vertical rays is situated in front of the retina.

4. *Compound Myopic Astigmatism.*—When the foci of both sets of rays are situated in front of the retina, but farther forward in the case (V, Fig. 25) of the vertical rays.

5. *Mixed Astigmatism.*—When the focus (H, Fig. 26) of the

horizontal rays falls behind the retina (hypermetropia in that meridian), and the focus (V) of the vertical rays in front of the retina (myopia in that meridian).

Symptoms of Astigmatism.—We may conclude that an individual is astigmatic if he sees horizontal (or vertical) lines, such as the horizontal portions of Roman capital letters, or the horizontal lines in music, or the horizontal rays in Snellen's Sunrise figure (see end of this book) distinctly, while the vertical (or horizontal)




Fig. 26.

lines seem indistinct. We have such a complaint, for example, when the retina is situated at the focus of the parallel rays passing through the vertical meridian of the cornea.

Suppose an eye to be emmetropic in the vertical meridian, and ametropic in the horizontal meridian: we must first consider how a point will be seen by such an eye. The rays of light emitted from the point and passing through the horizontal meridian will




Fig. 27.



Fig. 28.

not be brought to a focus on the retina, but will produce a blurring of the retinal image of the point at each side; while the vertical rays will unite on the retina, and consequently the point will appear distinctly defined above and below.

A line may be regarded as a number of points, and in order to understand how lines will be seen by an astigmatic eye, such as the above, it is only necessary to arrange a number of points in vertical and horizontal lines—as at *a* and *b* in Fig. 27. It is evident

at once from mere inspection that the horizontal line will appear distinct, because the rays which diverge from each point of the latter in a vertical plane—*i. e.*, at right angles to the direction of the line—are brought to a focus on the retina; while those rays diverging in a horizontal plane, although not meeting on the retina, do not render the picture of the line indistinct, because the diffusion images resulting from them exist in the horizontal direction, and consequently cover or overlap each other on the line, and therefore are not seen and do not confuse the sight. At the ends of the line only (*b*, Fig. 28) do the diffusion images cause a fuzziness or make the line seem longer than it is. In this case a vertical line (*a*, Figs. 27 and 28) seems indistinct, because, the horizontal meridian being out of focus, the diffusion images existing in that direction are very apparent, as they do not overlap. On the other hand, in order to see a vertical stripe accurately, it is necessary only that the rays diverging in a horizontal plane should have their focus on the retina; and therefore if an individual can only see vertical lines distinctly at 6 meters we know that his eye is emmetropic in the horizontal meridian (and probably myopic in the vertical meridian). We do not, however, hear this complaint as often as might be expected, because simple astigmatism is not so common as one or other of the compound forms.

Astigmatic people do not generally see very distinctly, either at long or at short distances.

Even in hypermetropic astigmatism the book is very often brought close to the eyes, in order, by increasing the size of the retinal image, to make up for its indistinctness.

Astigmatic individuals frequently suffer much from headache, due to constant effort to see distinctly, and we cure the headache when we correct the astigmatism.

It has been stated that epilepsy, if not capable of being produced by refractive errors, especially astigmatism, in persons with stable brains, may sometimes have such errors as its exciting cause where there is already a predisposition to the disease.

All these signs and symptoms appertain more to the rather high degrees of astigmatism. Slighter degrees may cause no annoyance beyond some indistinctness of vision; and indeed slight degrees of hypermetropic astigmatism often pass unnoticed until late in life, when the accommodation begins to fail.

We are often led to suspect and to seek for astigmatism when, in examining the refraction with spherical glasses, we are able to

bring about some improvement of vision, but cannot obtain normal V. with any glass, while there is no organic disease to account for the defect. Also if, in examining with spherical glasses, we find V. benefited equally by several glasses of considerable difference in power, even perhaps by convex as well as by concave glasses.

The ophthalmoscope affords us an admirable means of diagnosing astigmatism and of determining its amount. Just as the astigmatic eye cannot see horizontal and vertical lines equally well at the same moment, so is an observer unable to see both the vertical and horizontal vessels in the retina of the astigmatic eye simultaneously, but must alter his accommodation to be able to see first one set of vessels and then the other.

A comparison of the shape of the optic papilla, as seen in the upright and in the inverted images, may also give a clew to the presence of astigmatism. Inasmuch as the fundus oculi is very much magnified in the upright image by the dioptric media through which it is seen, and as this enlargement is greater in the direction of the meridian of shortest focus (meridian of highest refraction), which is most commonly the vertical meridian, a circular object, such as the papilla, will seem to be of an oval shape with its long axis vertical. But in the inverted image, in the meridian of highest refraction, the image lies nearer the convex lens than in the meridian of lowest refraction, and hence is much less magnified in the former than in the latter meridian; and here, consequently, the round optic papilla is seen as an oval with its long axis horizontal. Sometimes the papilla is really of an oval shape, and not round, and then the diagnosis is readily made by observing that in one image it is seen as an oval, while in the other image it is circular. Care must be taken in the indirect method not to hold the lens obliquely, as this would be sufficient to make a circular disc appear oval, the long axis of the oval being in the direction of the axis round which the lens is rotated.

The Estimation of the Degree of Astigmatism and its Correction.—It is evident that to correct astigmatism the ordinary spherical lenses would be of little use, for they affect the refraction of the light passing through them equally in every direction. We employ therefore what are termed cylindrical lenses, being sections of cylinders parallel to their axes, which refract light in one direction only—viz., that corresponding to their curvatures and at right angles to their axes. The rays which pass through these lenses in a direction corresponding to their axes are not refracted,

but pass on without deviation, as they would do through a piece of plane glass.

Although astigmatism is nowadays almost universally estimated by means of the ophthalmoscope, or by the astigmometer (see p. 61), yet in order to give the student a clear idea of the matter in the simplest way, I shall here describe a subjective method for its estimation, while its objective estimation by aid of the ophthalmoscope will be treated of in the next chapter.

Simple Astigmatism.—If, now, a case come before us in which we suspect astigmatism, we place Snellen's Sunrise (*vide* diagram at end of book), or some such diagram, at 6 meters from the eye (the other eye being excluded), and inquire of the patient whether there be any line which he sees much more distinctly than the others, and can trace farther towards the central point. If that be so, we know that he is emmetropic in the meridian at right angles to that line, provided his accommodation be at rest, and that he is ametropic in the meridian corresponding to that line.


In case the horizontal line below at each side be the distinct one, the eye is emmetropic in the vertical meridian, and probably hypermetropic in the horizontal meridian, because the latter is generally that of least curvature. Consequently, a convex cylindrical lens, held with its curvature horizontally (axis vertical) before the eye, will correct the defect. The highest convex cylindrical glass which gives the patient the best possible distant vision will be the correcting glass. This would be a case of Simple Hypermetropic Astigmatism (As. H.). If the lens required be + 2 D Cyl., it would be As. H. 2 D; and in prescribing for the optician we should write "+ 2 D Cyl. Ax. Vert."

If the central vertical line be the distinct one, then emmetropia exists in the horizontal meridian, and probably therefore myopia in the vertical meridian; and a concave cylindrical lens held before the eye with its curvature vertical (axis horizontal) will correct the defect. The lowest concave cylindrical lens which gives the patient the best possible distant vision will be the correcting lens. This would be a case of Simple Myopic Astigmatism (As. M.). If the lens be — 2.5 Cyl., it would be As. M. 2.5 D; and for the optician we should write "—2.5 D Cyl. Ax. Horiz."

I advise the reader to make now a few experiments for himself with cylindrical lenses, by means of which he can produce artificial astigmatism in his own eye. Let him place Snellen's Sunrise figure (end of this book) opposite his eye at a distance of about 4

to 6 meters. If he now hold a + 1.0 Cyl. before his eye, with its axis horizontal, it gives a myopia of 1.0 D to the vertical meridian of the eye, while the horizontal meridian remains emmetropic; and consequently he will see the central vertical line of the diagram distinctly, while the horizontal lines will be indistinct. By placing a — 1.0 Cyl. with its axis vertical before the eye, in addition to the + 1.0 Cyl., the artificial astigmatism produced by the latter is corrected, and the whole diagram becomes distinct. Every other kind and degree of astigmatism can be similarly represented by lenses and similarly corrected.

Compound Astigmatism.—If no line be very distinctly seen, then we may commence our examination with Snellen's Distance Test-Types, and test in the ordinary way with spherical lenses until we find that one which gives the best distant vision. This we place in a spectacle-frame before the eye, and proceed, as already explained, to ascertain the meridians of greatest and least curvature of the cornea. If the spherical lens be + 4 D, and with it




the horizontal lines in the Sunrise Diagram be the most distinct, then the vertical meridian is shown to be corrected, and the eye is probably still hypermetropic in the horizontal meridian, and requires a + cylindrical lens with its axis vertical, in addition to the spherical lens, to correct the entire defect. Suppose this cylindrical lens be found to be + 1 D Cyl., then the H. in the horizontal meridian will be shown to be 5 D, and the astigmatism to be 1 D.

The latter noted down would be of little practical value, and therefore we prefer to write in our note-books the factors of the Astigmatism, thus: "H. 4 D + As. H. 1 D Horiz."; or, as for the optician, "+ 4 D Sph. ⊃ + 1 D Cyl. Ax. Vert." * This is Compound Hypermetropic Astigmatism.

In an analogous way we examine for Compound Myopic Astigmatism, in which every meridian is myopic, but the vertical more so than the others.

* The sign ⊃ indicates "combined with."

Mixed Astigmatism.—Lastly we come across cases in which both concave and convex spherical lenses produce a certain amount of improvement, but neither give full vision. Placing, then, one or other before the eye in the spectacle-frame, the examination is proceeded with by aid of Snellen's Sunrise. We ascertain, for example, what is the lowest concave spherical lens which will bring out one horizontal ray distinctly. Let this be — 3 D; we have then myopia of 3 D in the vertical meridian. Now, having removed the — lens, we find what is the highest convex lens which will bring out one vertical line distinctly. Let it be + 5 D; this indicates hypermetropia of that amount in the horizontal meridian. We may correct such a case in either of two ways: (*a*) by a Sph. — 3 D, which will correct the vertical meridian, but will increase the hypermetropia in the horizontal meridian by 3 D, making it 8 D, which can then be corrected by combining a cylindrical lens of + 8 D, axis vertical, with the above spherical lens; (*b*) by a spherical + 5 D, which will correct the horizontal meridian, but



will increase the myopia in the vertical meridian to 8 D, necessitating the combination of a — Cyl. lens of that number with the + 5 D Sph. For reading, writing, etc., an overcorrection of the horizontal meridian with + 8 D Cyl., thus rendering the eye myopic 3 D in every meridian, and enabling the patient to read at, or near, his far point, might be the most suitable arrangement.

As it is necessary, in order to test the degree, etc., of astigmatism accurately, that the accommodation be at rest, it is desirable, before the examination for any of the hypermetropic forms, to instill atropin into the eye.

Measurement of the Degree of Astigmatism by the Astigmometer.— This is one of the most rapid and satisfactory methods of determining both the degree of astigmatism, and the position of the meridians of greatest and least refraction. It is based on the principle of the ophthalmometer, an instrument by which Helmholtz demonstrated the changes in the curvature of the lens during accommodation.

The cornea reflects images of objects in the same manner as a convex mirror, and the smaller the radius of curvature the smaller will the image of any given object be. It is easy to calculate the radius of curvature of the cornea knowing the size of the object, the distance of the object from the cornea, and the size of the




FIG. 29.

cornea, and the size of the corneal image. The only difficulty lies in the measurement of this image; and it has been found that the best method of effecting this is to double the image by looking at it through a double refracting prism, and then to alter the strength of the prism until the two images just come into contact. When this has taken place, a displacement equal to the size of the image

has been produced. The amount of displacement, and hence the size of the image, can easily be calculated.

The astigmometer was first brought into practical use by Javal and Schiotz. The instrument which is in use at the Royal Victoria Eye and Ear Hospital, and which has proved of great service, is a modification of Javal's, made by Kagenaar of Utrecht. In order



Fig. 30.

to measure the degree of astigmatism by it, we do not require to know the radius of curvature of the cornea, but merely to find out the difference in refractive power between the meridians of greatest and least curvature, and this the astigmometer enables us to do in a few seconds without any calculation.

It consists (Fig. 29) of a telescope (*p*) containing a double refracting prism between the object glasses, and two reflectors (*k* and *l*), which are movable on an arc (*m*), which is fixed to the telescope tube. The latter turns on its own axis, and enables the arc to be placed in any meridian, its position being indicated on a graduated circle (*g*). The patient places his chin on the rest *d*, and looks into the tube at *f*, the eye which is not under observation



Fig. 31.

being covered by the disc *e*. The surgeon then looks through the telescope at *u*, turns the arc *m* into a horizontal position, and observes the corneal images of the reflectors, which he gets into focus. He then moves the reflectors until the central images just come into contact; the four images will then occupy the relative positions shown in Fig. 30. The arc is then rotated into the ver-

tical meridian, and if the curvature of the cornea in this meridian be the same as in the horizontal meridian, the central images will still appear to be in contact; but if the radius of curvature in the vertical meridian be smaller, the intervals *a* to *b* and *a'* to *b'* will diminish, and consequently the central images will overlap, as in Fig. 31, each step of *a'* representing a difference of 1 Dioptry. So that in this case (Fig. 31) there would be an astigmatism of 2 D, and the greatest refraction would be in the vertical meridian.

It is generally desirable to begin with the arc in the horizontal meridian. If the axes of the meridians of greatest and least curvature are oblique, then the images will not lie in one line, and the arc must be turned until they do so lie. An index which moves on the circle *g* (Fig. 29) gives the position of the axes. It will be seen from the above description that the astigmometer merely registers the amount of astigmatism, but does not enable us to estimate the general refraction of the eye. Moreover, it is the corneal astigmatism alone which is determined, and it will be found that in the vast majority of cases this is the only astigmatism present.

Lental Astigmatism.—Disturbances of vision due to astigmatism often make their appearance for the first time at middle age or even later, and are then apt to be mistaken for amblyopia. In such cases the cornea has been astigmatic all through life, but the defect has been masked by a compensating astigmatism of the crystalline lens, produced by an unequal accommodative contraction of the ciliary muscle. When, now, as life advances, the amplitude of accommodation diminishes, the power of the ciliary muscle to produce this active compensatory lental astigmatism also diminishes, and finally disappears, and consequently the corneal astigmatism comes to the front; or, in astigmatic individuals the astigmatism may alter in degree at this time of life. Under atropin, too, astigmatism may appear, the existence of which was not previously known. This is termed active, or dynamic, lental astigmatism.

Passive, or static, lental astigmatism is due to irregularity in the shape of the unaccommodated lens, and gives rise to disturbances of vision similar to those caused by corneal astigmatism, or it increases existing corneal astigmatism, or it more or less completely compensates the corneal astigmatism. It has no clinical importance which does not attach to corneal astigmatism.

Irregular Astigmatism.

In irregular astigmatism, the refraction of the eye differs not only in different meridians of the eye, but even in different parts of one and the same meridian. It is frequently due to irregularities on the surface of the cornea, the result of former ulcers, and also sometimes to irregular refracting power in different parts of the crystalline lens. It cannot be corrected. Its presence can be detected by the distortion and irregular movement of the disc when the lens is moved during the indirect method of examining with the ophthalmoscope, and also by the irregular shadow in retinoscopy. In some cases there is a certain amount of regular astigmatism combined with it, correction of which may improve the vision.

Anisometropia.*

By this term is meant a difference in the refraction of the two eyes, one being myopic, hypermetropic, or astigmatic, while the other is emmetropic, or ametropic in a way different from its fellow. So long as the difference in refraction is but slight (say 1 D or 1.5 D), it is generally possible to give the correcting glass to each eye. When the difference is considerable it is often impossible to fully correct each eye, because, binocular vision having never really existed, the patients are unable to tolerate the presence of a clear image on each retina. We must then be content with correction of the least ametropic eye, or of that one which has the best vision; or we may partially correct the most ametropic, and fully correct the least ametropic eye. Each such case must be dealt with as it permits.

ANOMALIES OF ACCOMMODATION.

Presbyopia.

This is a diminution in the amplitude of accommodation, which commences at an early age, and is due solely to natural changes taking place slowly in the crystalline lens. It might not, therefore, strictly speaking, be considered as an anomaly. The power of accommodation commences to diminish in early childhood, the near

* *ἀ, priv.; ἰσος, like; μέτρον, a measure.*

point beginning then to recede from the eye. Donders it was who first discovered this fact, and ascertained the laws which govern the progressive decrease of accommodative power. He designed the accompanying diagram (Fig. 32), which illustrates the decrease from the tenth year of age, and indicates the amplitude of accommodation at different ages.

The numbers along the upper horizontal line refer to the ages, those along the left-hand perpendicular line to the dioptries. The curve *r r* shows the refraction of the eye when in a state of rest. This is unchanged until the fifty-fifth year, when it begins to




FIG. 32.

diminish; the emmetropic eye then becoming hypermetropic, the hypermetropic eye more hypermetropic, and the myopic eye less myopic. The curve *p p* shows the positive refracting power of the eye, corresponding to the punctum proximum, and its gradual diminution as life advances, and how at the age of sixty-five it becomes even less than the minimum refraction in former years. The two curves meet at the age of seventy-three, and then all power of accommodation ceases. The number of dioptries included between the two curves on the vertical line corresponding to any given age represents the amplitude of accommodation at

that age—*e. g.*, at thirty years of age the amplitude is 7 D; at fifty years it is only 2.5 D. The amplitude of accommodation is the same at the same age in all forms of ametropia, as well as in emmetropia.

The cause of presbyopia lies chiefly in a progressive change in the crystalline lens, which becomes less elastic and more homogeneous in its different layers, and refracts light less strongly than before. In more advanced life diminished energy of the ciliary muscle probably becomes a second factor in the production of presbyopia.

The near point gradually recedes from the eye until it reaches a distance beyond that at which the person usually reads, writes, sews, etc. Employments of this kind then become difficult, because the retinal images are too small to be clearly discerned, owing to the increased distance at which the work must be held from the eye; and, in order to make up for this smallness of the images, the individual is often seen to improve their brilliancy by procuring stronger light.

Presbyopia was defined by Donders to be present when the near point lies at more than 22 cm. from the eye, and we correct it by giving such a convex glass for reading, etc., as will bring the near point back to 22 cm. Now in order to see at that distance a positive refracting power (p) of ($\frac{100}{22} =$) 4.5 D is necessary, and if the eye have not so much positive refraction, a convex glass must be given to it of such power as will bring p up to 4.5 D; and this lens is the measure of the presbyopia. At the age of forty (*vide* Donders' diagram, Fig. 32) the eye possesses a positive refraction of just 4.5 D; and therefore from this age presbyopia (πρέσβυς, an old man; ὤψ,) is said to commence in emmetropic eyes. The presbyopia, then, is equal to the difference between the positive refracting power possessed by the eye and 4.5 D, and the number thus found is the correcting glass for the presbyopia. The distance of 22 cm. is rather close to the eyes for the comfort of most people, and 33 cm. is now commonly taken as the reading distance. Presbyopia on this assumption is thus postponed for two or three years.

It is important that in prescribing glasses for presbyopia, if there be any hypermetropic astigmatism present, it should be corrected by the suitable + cylinder lens added to the spherical glasses. It is also important that the glasses should be carefully centered for the reading distance—*i. e.*, that the visual lines, when they are con-

verged to the distance at which the work is held, should pass through the optical centers of the glasses. Or, if there be any insufficiency of the internal recti, it will be for the patient's comfort to decenter the lenses slightly inwards.

The following table indicates the presbyopia of the emmetropic eye:

Age.	⌀. required.	⌀. existing.	Presbyopia.
40	4.5	4.5	0.
45	4.5	3.5	1.0
50	4.5	2.5	2.0
55	4.5	1.5	3.0
60	4.5	0.5	4.0
65	4.5	0.25	4.25
70	4.5	−1.0	5.5
75	4.5	−1.75	6.25
80	4.5	−2.5	7.0

It is hardly necessary to point out that presbyopia comes on at a much earlier age in hypermetropes than in emmetropes; while in myopes its advent is postponed; or, in the higher degrees of myopia, it may not come on at all. The hypermetrope of 3 D would be presbyopic at the age of twenty-seven; because, in order to arrive at the 4.5 D of positive refraction required, he must have an amplitude of accommodation of (3 D + 4.5 D) 7.5 D, and this he has only up to that age (*vide* Fig. 32).

The myope of 4.5 D can get along until something over sixty years of age without any glass for reading (*vide* above table). At sixty-five, if he were emmetropic, he would have presbyopia of 4.25; consequently he will now require a + glass of only 0.25 D.

Presbyopia must not be mistaken for slight paralysis of accommodation. They are distinguished by the fact that in the former the amplitude of accommodation corresponds to the age of the patient as given in Donders' table, and the difficulty of near vision comes on gradually.

Paralysis of Accommodation.

This may be partial or complete, and one or both eyes may be affected. It is usually combined with paralysis of the sphincter iridis (mydriasis), and the condition is then called ophthalmoplegia interna; but it is also seen without paralysis of the sphinc-

ter, and either alone or with paralysis of some of the orbital muscles supplied by the third pair—rarely with paralysis of the external rectus.

The Symptoms are similar to those of presbyopia, but come on rather suddenly. They give inconvenience to the patient according to the state of his refraction. If he be emmetropic, his distant vision continues good, while his vision for near work is much impeded. If he be hypermetropic, as he requires his accommodation for distant objects, vision for distance is interfered with, and still more so vision for near objects. If he be myopic, vision is less affected than in either of the other forms of refraction; indeed, if he be very near-sighted, being able to see near objects at his far point, he may suffer little or no inconvenience.

Micropsia is a common symptom in cases of partial paralysis of accommodation, and is due to the fact that, while the retinal image is unaltered in size, the great effort of the defective accommodation gives the sensation of the object being much nearer to the eye than it really is.

Causes.—The most common cause of paralysis of accommodation is the action of atropin; but it is also the result of, or is attendant upon, various diseases. It is one of the symptoms of paralysis of the third nerve; it may be due to exposure to cold; or it may depend upon syphilis, syphilitic periostitis at the sphenoidal fissure, syphilitic gumma, or syphilitic inflammation of the nerve itself.

In cases of double paralysis of accommodation a central cause must often be looked for. Paralysis of accommodation and mydriasis are sometimes forerunners by many years of serious mental derangement.

Diphtheria is a frequent cause of paralysis of accommodation, usually without, but sometimes with, mydriasis. The onset occurs most commonly some weeks after the throat affection, which need not have been of a severe character. Indeed, the faucial attack may have had no apparent diphtheritic character, and may have been so slight as almost to have escaped the notice of the patient. The lesion in these cases is probably a nuclear one, and the evidence points to miliary extravasations of blood in the floor of the fourth ventricle; but there are those who hold that the paralysis is due to a poison, that it is a toxic paralysis.

In influenza paralysis of accommodation is seen, occurring sometimes in the acute stage and sometimes during convalescence.

One recorded case went on to bulbar paralysis, and ended fatally; but complete recovery is usual.

Paralysis of accommodation in middle life may be due to diabetes, and should make us suspicious of the presence of this disease.

Blows on the eye are apt to cause paralysis of accommodation, usually with mydriasis.

The Treatment depends, of course, upon the cause of the paralysis. The instillation of a 1 per cent. solution of sulphate of eserin or of muriate of pilocarpin may be employed in all cases, and will at least produce temporary improvement of sight; but it can hardly be said to assist in the cure, except perhaps in slight diphtherial cases. Iodid of potassium and mercury are indicated in syphilitic cases, and iodid of potassium and salicylate of sodium in rheumatic cases. The prognosis in these cases must be very guarded, as it often happens that recovery does not take place. Where cure does not result the patient may be enabled to make better use of his eye or eyes by means of a convex glass or spectacles; but in this matter each case must be dealt with for itself—no general rule can be laid down.

In diphtheritic cases a general tonic treatment, especially iron, is indicated; and here the prognosis is invariably favorable.

Accommodative Asthenopia

has been already treated of under the head of Hypermetropia (p. 40).

Spasm of Accommodation.

Spasm, or cramp, of accommodation in connection with hypermetropia and myopia has already been referred to. A few cases of acute spasm of accommodation have been reported.[4] Occurring in an emmetropic or slightly hypermetropic eye, such a spasm produces apparent myopia. In some of the cases there was no assignable cause for the spasm, in some it was due to overwork, and in one to trauma of the cornea. The treatment is a lengthened course of atropin locally.

References.

[1] "Trans. Internat. Ophth. Congress," 1888, p. 97.
[2] "Von Graefe's Archiv," xxxvi. 2, p. 1.
[3] "Zeitschrift für Augenheilkunde," vol. v. p. 171.
[4] "Von Graefe's Archiv," xxxvi. p. 232.

CHAPTER III.

THE OPHTHALMOSCOPE.

ALTHOUGH the dioptric media of an eye may be perfectly clear and normal, yet no detail of its fundus can be discerned by the unaided eye of an observer who looks through the pupil, the latter being for him merely a dark opening. The reason of this is that these dioptric media are composed of a system of convex lenses. To explain: Suppose the inside of a small box (*vide* Fig. 33) to be blackened, and on its floor some printed letters fastened, and a hole cut in the lid, which is then replaced—it will be found that, by aid of a lighted candle and with a little experimentation, the letters may be read through the aperture. The rays passing from the light (*L*) into the box through the aperture illuminate the




FIG. 33.

opposite surface, and from this surface the rays *a*, *b*, and others pass out again through the opening, and some of them fall into the observer's eye at *E*.

But if, in order to make this box represent an eye more accurately, we place a convex lens immediately within the aperture, the course of the rays is altered. All the rays passing into the box (Fig. 34) from *L* are brought to a focus on its opposite side at *m* by the convex lens *n*, and according to the optical law of conjugate foci, all the rays passing out from the box meet again at the source of light (*L*), and hence none of them can be received by the eye (*a*) of the observer; nor can this eye be placed in any

position where it could catch any of these rays, for if it be placed anywhere between the aperture and *L*, it would cut off the light passing from *L* into the box.

Helmholtz's Ophthalmoscope.—If the eye of the observer could itself be made the source of light, the difficulty would be solved;



Fig. 34.

and, practically, this is what Helmholtz accomplished with his ophthalmoscope in the year 1851. The instrument he invented was composed of a number of small plates of glass (*O*, Fig. 35), from which light from *L* was reflected into the eye (*E*), and thus the fundus of the latter illuminated. From *m* rays pass back again by the same course to the ophthalmoscope, some being reflected back to *L*; but some, passing through the ophthalmoscope, and falling into the observer's eye placed close behind the instrument at *a*, form in it an image of *m*.

Modern Ophthalmoscope.—For the original ophthalmoscope of Helmholtz a concave mirror of 20 cm. focal length with a central




Fig. 35.

opening has been substituted. This mirror (*O*, Fig. 36) throws convergent rays into the eye (*E*); and these, being made more convergent by the refracting media, cross in the vitreous humor, and light up part (*a b*) of the fundus. From every point of this illuminated surface rays are reflected back again out of the eye.

If the latter be emmetropic the rays from any one point become parallel on leaving it; and some of these parallel rays, passing through the aperture (*c*) of the ophthalmoscope, fall into the observer's eye, and, if it be emmetropic, are brought to a focus on its retina; the rays from *m* at *m'*, those from *x* at *x'*, and those from *y* at *y'*—and thus an image of the part *x m y* is formed on the observer's retina.

The foregoing method of examining with the ophthalmoscope is called the **Direct Method,** or the **Examination of the Upright Image.** By it the various parts of the fundus are seen in their natural positions, but much enlarged (about 15 diameters in the emmetropic eye); and it is consequently very valuable for examining minute details.

It is necessary for this method that the surgeon should approach




FIG. 36.

his eye as close as possible to the eye under examination, in order to receive as much of the light coming out of it as possible.

It is also necessary for this method that the accommodation both of the surgeon's and of the patient's eye be at rest, as otherwise the rays coming from the latter cannot form an image on the retina of the former, at least if both be emmetropic.

If the patient exert his accommodation, the rays will, on leaving his eye, become convergent instead of parallel, and, falling into the surgeon's eye, will be brought to a focus in front of his retina. If the surgeon exert his accommodation, the parallel rays from the patient's eye will likewise, on falling into his (the surgeon's) eye, be brought to a focus in front of his retina. And if both patient and surgeon accommodate, the focus of the rays from the patient's fundus oculi will, of course, lie still farther in front of the surgeon's retina. The patient's accommodation can be relaxed by

making him gaze at the black wall behind the surgeon's head, or his accommodation may be paralyzed with atropin. But atropin should never be used unless absolutely necessary, owing to the inconvenience it causes the patient.

Voluntary relaxation of the accommodation on the part of the surgeon is often a matter of much difficulty to beginners. The ciliary muscle not being a voluntary muscle is not under our direct control, and can be influenced only in a secondary way through the convergence of the optic axes; for this convergence is regulated by voluntary muscles (the internal and external recti), and is intimately associated with the effort of accommodation. With parallel optic axes our accommodation is relaxed; therefore, when we want to relax our accommodation, we produce parallelism of our optic axes. This sounds easy enough; yet, when the beginner approaches his eye close up to that of his patient, the knowledge




Fig. 37.

that he is so close to the object he wishes to see renders the accomplishment of this parallelism and relaxation of accommodation very difficult to many.

It is not easy to teach another person how to relax his accommodation, but the following hint may be of use. Take a printed page, and hold it at the ordinary reading distance, so that the type may be clearly seen; then gaze vacantly at it, so that the type may become indistinct. The accommodation is now relaxed, and the act is accompanied by a peculiar sensation in the eyes. When examining in the erect image, cause this same sensation to take place; and it may be assisted if, with the eye which is not in use, the black wall behind the patient's head be gazed at.

The **Indirect Method**, or the **Examination of the Inverted Image,** is employed in order to obtain a more general view of the fundus than the direct method admits of.

In addition to the ophthalmoscope a convex glass (*l*, Fig. 37) of about 14 D is here used. The latter is held about 10 cm. from the eye (*E*) under examination, while the observer throws the light through it into the eye. In passing through the lens the rays are made convergent, and this convergence is increased by the refracting media, so that the rays cross in the vitreous humor, and light up a portion of the fundus oculi. From any points (*a* and *b*) of this illuminated place pencils of rays pass out again from the eye, and, becoming parallel, pass through the lens, and are united by it at *a' b'*; and thus a real inverted image is formed of the part *a b*, which image may be seen by the observer whose eye is placed behind *O*. The stronger the lens (*l*) the more convergent must rays from the examined eye be made; and consequently the closer must *a' b'* be to each other, and the smaller and brighter must be




Fig. 38.

the image formed. The weaker the lens (*l*) the larger and less brilliant is the image, and the less annoying to the surgeon are the reflexes from the surfaces of the lens.

In examining by the indirect method the observer first places the upper edge of the ophthalmoscope to his right supra-orbital margin, and, taking care that he is looking through the central opening of the mirror, he reflects the light of the lamp into the patient's eye at a distance of about 50 cm. A red glare from the fundus will then be seen in the pupil. Keeping the pupil illuminated, the convex 14 D, held between the forefinger and thumb of the surgeon's left hand, is brought up in front of the patient's eye, and kept there in the perpendicular position, the surgeon steadying this hand with the tip of the little finger on the patient's forehead. The convex glass is now removed just far enough from the patient's eye to cause the margin of the pupil to disappear out of the

surgeon's field of vision. The observer then ceases to look into the eye, and fixes his gaze on the convex glass, when the inverted image of the fundus should at once become visible—and will seem to be situated in the convex lens, although it really is in the air somewhat this side of the lens.

The diagram (Fig. 38) serves to illustrate the effect of inversion of the image.

The left eye is seen in the upright image in the left-hand picture, while the same is seen in the inverted image in the right-hand picture. In the diagram the two images are of the same size for the sake of convenience; although, of course, in reality the upright image is much larger than the inverted image. Moreover, it should not be supposed that nearly the whole fundus oculi, as here represented, can be taken in at one view with the ophthalmoscope. The portion visible with the ophthalmoscope at one moment, even in the inverted image, is small; so that it is necessary to examine the different regions in detail in order to become acquainted with their condition.

The reflex from the surface of the cornea gives a good deal of annoyance to every beginner. It cannot be done away with; but as it moves in the opposite direction to a motion of the object lens it is possible to see past it. The reflections from the convex object-lens are also extremely annoying, but may be removed to a great extent from the line of sight by a slight rotation of the lens on its axis. If a very high convex lens (say + 20 D) be used, the reflections from it are more disturbing than from a lower number (say + 14 D).

To examine the *Optic Nerve* the surgeon sits in front of the patient, and directs him to turn his eye somewhat to the nasal side, and slightly upwards; because the papilla, or disc, is situated about 15° to the inner side of the posterior pole of the eye, and about 3° above it. For instance, if the left eye be examined the patient is to direct his gaze, without turning his head, to the right and a little upwards, say towards the surgeon's left ear. It is well always to seek out the optic papilla in the first instance, not only because it is so important a part of the fundus oculi, but also because, examining from it towards the periphery, we are the better able to determine the locality of any pathological alteration.

Should the patient not direct his gaze in such a way as to enable the surgeon to see the optic disc or other desired region, it may be brought into view either by a motion of the surgeon's head in

the opposite direction, or by a motion of the convex lens in the same direction, or by a combination of both these maneuvers.

The *Macula Lutea* should then be examined. It may be seen by directing the patient to look straight at the hole of the ophthalmoscopic mirror, for it will then correspond with the macula lutea of the observer's eye. It is more readily seen in the inverted than in the upright image; but its examination is often very difficult, owing to contraction of the pupil produced by the strong light falling on so sensitive a portion of the retina, and by the reflections from the surfaces of the cornea and crystalline lens, which fill the area of this contracted pupil. It is therefore a better plan to direct the patient to look somewhat to the side of the eye under examination—*e. g.*, to the right side of the observer's forehead, if the left eye be under examination, and then by motions of the convex lens to bring the macula lutea into view.

After this the *Periphery of the Fundus* in every direction is to be examined by making the patient look upwards, downwards, to the right, to the left, etc.

Estimation of the Refraction by Aid of the Ophthalmoscope.

From what has been said with reference to the Direct Method of ophthalmoscope examination, it will have become evident that this method affords a means for determining the refraction of the eye.

At a little distance from the observed eye into which light from the ophthalmoscope mirror is thrown, the surgeon will be able to make the *qualitative* diagnosis of the refraction. If he can see some of the details of the fundus, it is either myopia or hypermetropia; but if it be emmetropia, he will be unable to see any detail. The explanation of this is that in myopia the rays coming out of the eye form an inverted image at the far point of the eye in the air, and this image can be seen by the observer who accommodates his eye for that point. In hypermetropia the rays coming out divergently from the eye pass into the observer's eye, and by an effort of accommodation on his part he will see an upright image of the portion of the patient's fundus oculi from which they come. But in emmetropia, inasmuch as the rays come out parallel, those from any two points (*m n*, Fig. 39) at a short distance from each other in the fundus on emerging from the eye diverge quickly from

each other, and the observer a little way off (at *A*) receives none of them into his eyes, or obtains only an indistinct image or red glare. If he go very close to the eye he can see details.

If on the observer moving his head from side to side the vessels, etc., of the observed fundus move with him, the case is one of hypermetropia, because the image is an erect one, which is situated behind the plane of the pupil to which it is referred. If the vessels, etc., move in the opposite direction to that of the observer's head, the observed eye is myopic, because there the image is inverted and in front of the pupil.

For the *quantitative* determination of ametropia a refraction ophthalmoscope is required. This instrument provides a number of convex and concave lenses capable of being brought into posi-




Fig. 39.

tion behind the sight-hole in rapid succession by a simple mechanism, and the direct method (p. 72) is employed.

It is necessary, in the first instances, that the surgeon be acquainted with the nature of his own refraction.

If the Surgeon be Emmetropic he can see the fundus oculi of an emmetrope in the upright image without any lens, provided he go close enough, as the parallel rays coming from the examined eye will be focused on his retina, because his eye is adapted for parallel rays.

In order to see the fundus oculi of a hypermetrope without any effort of accommodation he must place such a convex lens behind his ophthalmoscope as will render the divergent rays coming from the patient's eye parallel before they pass into his eye. This lens is the measure of the patient's hypermetropia, because it shows how many dioptries the eye wants of being emmetropic; or, in

other words, so that the rays coming from it may be made parallel. The lens which makes the divergent rays coming from the patient's retina parallel would also give to parallel rays passing into the eye such convergence that they would meet on the retina—*i. e.*, it would correct the hypermetropia.

The emmetropic surgeon can of course see the fundus oculi of a hypermetrope by the direct method without the correcting glass if he use his accommodation to overcome the divergence of the rays, and this is usually the case in the lower degrees of hypermetropia. The surgeon generally relaxes his accommodation according as he substitutes convex lenses for it, until he reaches the *strongest* lens with which he can distinctly see the fundus. This is the correcting lens.

To see the fundus oculi of a myope the emmetropic surgeon must place a concave glass behind his ophthalmoscope, in order that the convergent rays coming from the observed eye may be made parallel before they pass into his eye; and the *lowest* concave lens which enables him to see the fundus oculi is the measure of the myopia, as showing by how many dioptries it is in excess of emmetropia.

The emmetropic surgeon cannot possibly see the fundus oculi of a myope without the correcting glass, as the rays are brought to a focus in front of his retina, and if he use his accommodation he merely makes them still more convergent. But by means of an effort of his accommodation he can see the myopic fundus with a lens which overcorrects the myopia, and hence the importance of selecting the *weakest* concave glass with which the fundus is distinctly seen.

If the surgeon be ametropic, he may either correct his ametropia by wearing the suitable lens, and then proceed as though he were emmetropic, or else, and which is perhaps the better plan, he may add or subtract the amount of his ametropia from that of his patient's. For example:

The *Hypermetropic Surgeon* of say 3 D requires a + lens of 3 D in order to see an emmetropic fundus oculi, this lens going altogether to correct his own defect. If in order to examine the fundus of another eye he require a + lens of 6 D, the examined eye must be hypermetropic 3 D, the other 3 D going to correct the surgeon's H. If he be able to see the fundus oculi under observation without any lens, it shows that the eye has an excess of refraction corresponding to the want of refraction in his own eye—

that is to say, it is myopic 3 D. If he require a concave 2 D, his want of refraction—his hypermetropia—is not enough by that number of dioptries, and he has to do with an eye which is myopic 5 D (3 D + 2 D). Again, if he can see the fundus distinctly with a + lens, say + 1.0, which is less than his own correcting glass, this shows that the eye he is examining is myopic, but myopic to a lesser degree—in this instance by 1 D—than he himself is hypermetropic, and the examined eye here would be M. 2.0 D (*i. e.*, 3.0 — 1.0).

If the Surgeon be Myopic say 2 D, he requires a — 2 D to see the fundus of an emmetropic eye, this lens going wholly to correct his own ametropia. If he see the fundus with a — 7 D, the examined eye has M. 5 D, because 2 D has been used in correcting the surgeon's M. If he be able to see a fundus without any lens, the patient has H. 2 D, the want of refraction in the latter's eye compensating exactly for the excess of refraction in the surgeon's eye. If he finds it necessary to use a + lens of 7 D, it will indicate that his excess of refraction is not able to make up for the defect of refraction in his patient's eye, and that the latter has H. = 9 D (2 D + 7 D). If he have to use a — lens, say — 1.0 D, which is less than his own correcting glass, this shows that the eye he is examining is hypermetropic to a lesser degree—in this instance by 1.0 D—than he himself is myopic, and the hypermetropia here would be 1.0 D (*i. e.*, 2.0 — 1.0).

The Existence and Degree of Astigmatism may be Determined with the Ophthalmoscope.—We know that astigmatism is present if in the upright image we see the upper and lower margins of the disc and the horizontal vessels well defined, while the lateral margins and the vertical vessels are blurred, or *vice versa.* Again, we know that astigmatism is present if in comparing the shape of the optic disc in the upright and inverted images we find it to be an oval with its long axis perpendicular in the former, and with its long axis horizontal in the latter, showing that the refracting media are more powerful in the vertical than in the horizontal meridian.

We may ascertain the kind and degree of astigmatism as follows:

If in the upright image with relaxed accommodation we can see the retinal vessels in one meridian distinctly, while in order to see those in the opposite meridian a concave or convex lens behind the ophthalmoscope is required, we know that the case is one of

simple myopic or hypermetropic astigmatism; the emmetropic meridian being that at right angles to the vessels* seen without any lens, and the number of the lens indicating the amount of ametropia in the other meridian.

If in the two principal meridians two concave lenses or two convex lenses of different strength be required, we have to deal with a case of compound astigmatism, myopic or hypermetropic; the greatest error of refraction being in the meridian at right angles to that one, the vessels of which are made distinct by the strongest lens.

If a concave lens be required to bring into distinct view the vessels in one meridian, while a convex lens is required for the opposite meridian, the case is one of mixed astigmatism. Myopia exists in the meridian at right angles to that in which the vessels are brought into view by the concave lens, and hypermetropia exists in the opposite meridian.

I would again impress upon the reader the absolute necessity of thoroughly relaxing his accommodation in all examinations in the upright image.

Retinoscopy.

Another and most useful method for determining the refraction by the ophthalmoscope is termed the Shadow Test, or Retinoscopy. The appearances upon which this method depends are due to the play of light reflected from the mirror on the fundus oculi. Either a concave or a plane ophthalmoscopic mirror may be employed.

Retinoscopy with the Concave Mirror.—If the rays from a light (*L*, Fig. 40) be reflected from the concave mirror (*m*) of an ophthalmoscope, they cross at a certain point (*A*), and form there an inverted image of the flame, and then diverge again. If these diverging rays be made to pass through a convex lens (*B*) placed at such a distance in front of a screen (*E*) that the rays meet at a focus on the latter, a very small and brilliant upright image (*O*) of the flame is there formed, surrounded by a deep shadow. If the screen be moved slightly towards the lens (to *H*), so that the focus of the rays would lie behind it, or if it be removed slightly away from the lens (to *M*), so that the focus come to lie in front

* The vessels may be regarded as lines, and the explanation given on pp. 55 and 56 applies to them also.

of it, the brilliancy of the image on the screen and the intensity of the surrounding shadow are reduced; because in each instance a circle of diffusion, and not an accurate image, is formed on the screen, and the farther the focus of the pencil of rays is situated from the screen in either direction the weaker does the image become and the more ill-defined the shadow.

If the mirror be rotated in various directions the illuminated part* and the shadow are seen (care being taken to look at the screen directly, and not through the lens) to move on the screen in the opposite direction to the motion of the mirror. For example, if the position *m* (Fig. 40) be given to the mirror, the path of




Fig. 40.

the rays reflected from it is shown by the dotted lines, and the image of *O* is moved to *O'*. This will also be the case if the screen be at *H* or at *M*. Now these three positions of the screen may be

* "The area of light," "the image," "the illuminated area, or part of the fundus," and "the illumination" are different terms for one and the same thing. "The shadow" or "shade" refers merely to the margin of the illuminated area—*i. e.,* where the illumination ceases and darkness begins; it does not mean that the shadow of any object is thrown on the fundus oculi. When we speak of the motion of the shadow we mean that the margin of the illuminated area, or boundary-line between illuminated and non-illuminated area, moves along with the illuminated area in response to the motion of the mirror. It is easier to see how the illuminated area moves by watching the margin of the shadow (which comes across the pupil from behind the iris like a revolving shutter across a shop window), and hence it is that we have come to talk always of the motion of the shadow and not of the motion of the illuminated part.

supposed to represent emmetropia (E), hypermetropia (H), and myopia (M). Fig. 40 more particularly illustrates the motion of the light and shade in E and H only, while Fig. 41 demonstrates that in M.

In the eye, in like manner, the area of light and shade in the pupil moves against the motion of the mirror. Now we cannot, of course, see the real motion on the retina directly, but only through the dioptric media, and they will influence the apparent motion according to the condition of the refraction.

In emmetropia and in hypermetropia the rays coming out of the observed eye are parallel and divergent respectively; and, conse-




FIG. 41.

quently, an upright image being formed by them in the observer's eye, the true motion given by the mirror is perceived.

In myopia, at least in all cases of more than 1 D, the observer does not see an upright image of the flame on the fundus of the observed eye, but a real inverted aërial image formed between his mirror and the observed eye. The reason of this is that the rays coming out of the patient's eye are convergent, and meet at a focus, which is the far point of the eye, and form there an inverted image of the object from which they come, and which, in this instance, is an upright image of the flame (the illuminated area). When, therefore, the upright image on the fundus moves against the mirror, the inverted image (which the observer sees) moves in the opposite direction—*i. e.*, with the mirror. For example, if in Fig. 41 we suppose a to be the position of the image on the

fundus of a myopic eye, and a^2 the position of its real inverted aërial image, a motion of the mirror to m' (the rays reflected from m' are omitted in order to avoid confusion in the diagram) throws the image of a to a', as already explained, but the inverted aërial




Fig. 42.

image of a' is formed at a''—*i. e.*, it seems to have moved with the mirror.

In myopia alone, then, does the image move with the mirror; while in emmetropia and hypermetropia it moves against the mirror. In low myopia (1 D and less), as will just now be seen, the image also moves against the mirror.

From what has been said it is evident that the higher the ametropia (the farther from the screen, in Fig. 40, the focus of the rays) the larger and feebler the illumination becomes (*i. e.*, the




Fig. 43.

greater the circles of diffusion), and the more crescentic the margin of the shadow, because it is the margin of a circle of diffusion.

Again, the extent of the motion of the image and the rate of this motion are in inverse proportion to the degree of the ame-

tropia. Thus, if Fig. 42 represents a myopic eye, whose far point is situated at a^2, a motion of the mirror to m' may be supposed to throw the illuminated part to a', and then a^2 will move to a'. But if the myopia be of less degree, so that the far point is at a^3, the same motion of the mirror will throw a^3 to a', and the distance between these two latter points is evidently much greater than that between a^2 and $a^{2'}$. In a hypermetropic eye (Fig. 43), the image may be supposed to be formed at a, and a motion of the mirror to m' will throw it to a'; while in a lower degree of hypermetropia it would be formed at b, and the same motion of the mirror would throw it to b'. The distance between b and b' is much greater than that between a and a'.

In practicing retinoscopy with the concave mirror the surgeon sits 1.20 m. in front of the patient. The eye to be examined is shaded from the direct rays of the lamp, if the latter be placed beside the patient; but a better plan is to have the light above his head. The focus of the mirror should be 22 cm., and any error of refraction of the surgeon is to be corrected. The light is then thrown into the eye at an angle of about 15° with its axis of vision, so that if the pupil be not under the influence of atropin the macula lutea may be avoided. When now the ophthalmoscope is rotated in different directions, motions of the light and shade on the fundus oculi are seen in the pupillary area. The surgeon directs his attention to the *edge of the shadow* rather than to the illuminated part, for its motion is more easily appreciated. If the edge of the shadow move *with* the motion of the mirror, *myopia* is present; if it move *against* the mirror, emmetropia, hypermetropia, or myopia of only 1 D or less is present.

The reason why the motion is against the mirror in cases of M. 1 D and less is that the surgeon being seated only 1.20 m. from the eye he is examining, if that eye have a myopia of 1 D, its far point is so close to his eye that he cannot clearly observe the image there formed; but if the myopia be of even slighter degree, the image will be formed behind the surgeon's head, and he gets a shadow moving against the motion of his mirror, because the image he then sees is the upright one of the patient's fundus oculi, and not the inverted aërial image.

We proceed as follows:

A trial spectacle-frame is put on the patient's face. If the shadow move with the mirror, we know at once the eye is myopic. To find the degree of myopia the surgeon puts a low concave-glass

(say — 1 D) into the frame; and if the shadow still move with the mirror, he puts in a higher number (say — 1.5 D), and so on until he comes to a glass which makes the image move against the mirror. If this be — 3 D, the myopia is 3 D. It might be supposed, as the shadow now moves against the mirror, that this glass overcorrects the myopia; but this is not so, because, as already explained, when the myopia is very low the image is formed close to the surgeon's eye, or behind his head, and he consequently gets a shadow moving against the mirror, although low myopia, and not emmetropia, is present. Consequently — 0.5 D, or — 1 D, has to be added on to the lens, which gives the effect of no distinct shadow; or rather, by the above plan, it is not deducted from the lowest lens, which makes the shadow move against the mirror.

If the shadow move against the mirror, we have to determine whether the eye is emmetropic, hypermetropic, or slightly myopic. Should the illumination be bright, and the shadow well defined, the eye is emmetropic, or not far removed from it; and if the shadow be ill defined and crescentic, we may feel sure the eye is highly hypermetropic. We first put on + 1 D, and if the motion be still against the mirror, the case is one of hypermetropia, and higher numbers are at once proceeded with until that one is reached which causes the shadow to move with the mirror. The measure of the hypermetropia is 1 D less than the glass so found, for it has evidently overcorrected the defect.

If, however, on putting on + 1 D we find the shadow to move with the mirror, we change it for + 0.5 D; and if still the motion be with the mirror, the eye is, beyond doubt, slightly myopic, — 0.5 D or so. But if with + 1 D the shadow move with the mirror, while with + 0.5 it continue to move against it, the eye is emmetropic.

It may be found that in two opposite meridians there is a difference in the motion of the shadow, and this leads us to diagnose the presence of astigmatism. When the difference is one merely of rapidity of motion, or of intensity of illumination and shadow, we know that we have to do with either simple or compound astigmatism. But if in the two meridians there be a difference in the direction of the motion, then it is a case of mixed astigmatism. The best method for ascertaining the degree of astigmatism and its correcting glass is to correct each of the principal meridians separately with spherical lenses. In compound

astigmatism the difference between the two lenses found indicates the degree of astigmatism and also the cylindrical lens which, combined with the correcting spherical lens for the least ametropic meridian, is required to neutralize the defect. In mixed astigmatism the addition of the two numbers gives the cylindrical lens, while one or other of them, usually the + D, is used as the spherical lens.

Retinoscopy with the Plane Mirror.— With the plane mirror the source of illumination of the observed eye is not a real inverted image of the light, as in the case of the concave mirror, but a virtual upright image behind the mirror; and as this image moves in the opposite direction to the motion of the mirror, the motion of its illumination on the fundus of the patient's eye must be *with* the mirror in all cases, and not against it, as in using the concave mirror.

With the plane mirror, therefore, the shadow is seen to move *with* the motion of the mirror in H. and E.; but in M. it *seems* to move *against* the motion of the mirror, for what we here see is an inverted image of the fundus situated at the far point of the eye. If the myopia be high, this inverted image will be close to the eye; if low, it will be far away from it. In using the plane mirror it is important to remember this point, because, if the observer go nearer to a myopic eye than its far point, he will not obtain a myopic motion, but one which is the same as that in E. or H. Consequently, in using the plane mirror, the rule is to go as far from the eye under examination as possible. If at the beginning the surgeon retire a little more than 2 meters from the eye, and there obtain a with-motion, he at once knows that the eye is not myopic 0.5 D; or if he stand a little more than 4 meters away, and obtain the same motion, he knows there is not a myopia of even 0.25 D present. If the myopia be high, he will be able to begin close to the patient, but must gradually retire from the eye as he increases the number of the concave glass put up—for the far point is thereby moved farther off—in order that he may not think he has corrected the myopia before he really has done so. Again, if at every distance the motion be with the mirror, the surgeon has to decide whether this indicates E. or H. He does this by putting a low + lens (say + 0.25) before the patient's eye, and if then, standing at a distance of 4 meters, the motion be altered by this glass to one against the mirror, he knows that the eye has not a hypermetropia of 0.25 D, consequently that it is

emmetropic. But if this lens does not at that distance cause a change in the motion of the shadow as originally obtained, the eye must be hypermetropic to at least the extent of 0.25 D; and, in order to ascertain how much more of H. than this may be present, it is now only necessary to go on increasing the strength of the lens in front of the patient's eye until one is reached which at 4 meters from the eye produces the myopic motion. The observer knows that he has now slightly overcorrected the hypermetropia of the eye, and that the next lens lower is its measure.

With some practice it is possible, unless the pupil be small, to obtain sufficient light from the fundus with the plane mirror at a distance of 4 meters.

I find this method much more easily worked than that with the concave mirror. It has the advantage, too, of not requiring any wearisome addition to, or subtraction from, the data obtained.

The pleasantest plane mirror is one of 4 cm. diameter, and of which the sight-hole is 4 mm. in diameter.

FOCAL OR OBLIQUE ILLUMINATION.

is employed for the *examination of the cornea, iris, and anterior part of the lens.* With a high + lens (16 to 18 D) the light of the gas flame is concentrated on the part to be examined with an oblique, not a perpendicular, incidence of the concentrated rays. Small foreign bodies in the iris, cornea, or lens, or opacities in either of the latter can be thus detected. Extremely delicate opacities in the cornea are not seen best with the strongest illumination which can in this way be produced, but rather by the half-light which is obtainable at the edge of the cone of light passing from the lens. In examining the center of the crystalline lens the incidence of the light must necessarily be more perpendicular.

But opacities in the refracting media can be best observed by examination with strong convex lenses in the upright image. The farther forward the opacity lies the more hypermetropic (so to speak) it is, and the stronger the lens required. Very minute opacities of the cornea can be seen in this way with a + 18 D or + 20 D lens in the ophthalmoscope.

THE NORMAL FUNDUS OCULI AS SEEN WITH THE OPHTHALMOSCOPE.

Reference has been made to the enlargement of the image of the fundus oculi seen with the ophthalmoscope. The cause of this enlargement is that the fundus is observed through a dioptric system at or close to the principal focus of which it is situated, and which consequently magnifies it to our view. The enlargement of the inverted image is not so great as that of the upright image, and it is smaller the shorter the focal length of the convex lens employed. The inverted image of a hypermetropic eye is larger than that of an emmetropic eye, and the latter larger than that of a myopic eye. It is possible to determine mathematically the degree of enlargement of the image; but into this it is not necessary to enter.

The Optic Papilla, or Optic Disc.—This is the first object to be sought for by the observer. It presents the appearance of a pale pink disc, somewhat oval in shape, its long axis being vertical. Occasionally the long axis lies horizontally, and sometimes the papilla is circular. The papilla is generally surrounded by a white ring, more or less complete, called the sclerotic ring, and often, outside this again, by a more or less complete black line, the chorioidal ring. The sclerotic ring is due to the chorioidal margin not coming quite up to the margin of the papilla, the foramen in the chorioid for the passage of the optic nerve fibers being somewhat larger than that in the sclerotic, and, consequently, a narrow edging of the white sclerotic is exposed. The chorioidal ring is the result of a hyperdevelopment of pigment at the margin of the chorioidal foramen. The complexion of the optic disc results from the pink hue derived from its fine capillary vessels, combined with the whiteness of the lamina cribrosa and the bluish shade of the nerve fibers. It is frequently not equal all over, but is paler on the outer side, where the margin is more defined and where the nerve fibers are often fewer than on the inner side. The apparent color of the papilla depends also upon the complexion of the rest of the fundus. If the latter be highly pigmented, the papilla appears pale in contrast; while, if there be but little pigment in the chorioid, the papilla may appear very pink. The complexion of every normal papilla is not identical, and care must be taken not to make the diagnosis "Hyperemia of the papilla" where merely

a high physiological complexion is present. The upper and lower margins of the papilla are often, especially in young people, a little indistinct, and show a delicate striation by the direct method of examination. This may be greatly exaggerated in hypermetropes, and has in them been sometimes erroneously taken for optic neuritis.

A physiological excavation of the optic papilla is often met with. It is always on the temporal side of the papilla, and can be recognized from the parallax* which may be produced, and from the paleness of this portion of the papilla. When the excavation is very deep, one may sometimes observe the lamina cribrosa in the form of gray spots (the nerve fibers) surrounded by white lines (the fibrous tissue of the lamina).

A physiological excavation differs from a pathological excavation by the fact that it does not reach the margin of the papilla all round. It is caused by the crowding over of the nerve fibers to the inner side of the papilla. Yet sometimes a healthy optic papilla will be met with in which the excavation apparently reaches the margin all round. Doubtless in such cases the thickness of the translucent nerve-fiber layer alone it is which is interposed between the sclerotic margin and the margin of the cup all round.

The Normal Retina is so translucent that it cannot be seen, or at most a shimmering reflection or shot-silk appearance is obtained from it, particularly about the region of the yellow spot and along the vessels, but also towards the equator of the eye, and especially in dark eyes, and in young people.

A peculiar, but physiological, appearance known as "opaque nerve fibers" is occasionally seen. It is produced by some of the nerve fibers forming the internal layer of the retina regaining the medullary sheath on the distal aspect of the lamina cribrosa, or near the margin of the papilla, which they had lost in the optic nerve just before entering the lamina cribrosa; the rule being that the nerve fibers lose their medullary sheath at the latter place definitely, and enter the retina as axis cylinders only, and hence are quite translucent. Instead of that, in these cases their fibers reflect the light strongly, giving the effect of an intensely white spot, commencing at the disc, extending more or less into the surrounding retina, and terminating in a brushlike extremity. This appearance is constant in the rabbit's eye.

The Macula Lutea is generally seen as a bright oval ring

* For explanation of the parallax see chap. xii.

with its long axis horizontal, this ring being probably a reflex from the surface of the retina. It is remarkable that this halo is not visible with the direct method of examination—a fact due probably to the illumination being much weaker than with the indirect method. The area inside the ring is of a deeper red than the rest of the fundus, and at its very center there is an intensely red point, the fovea centralis. This ring is not seen in old people.

The General Fundus Oculi surrounding the optic papilla and macula lutea varies a good deal in appearance according to the




Fig. 44—(*Graefe and Saemisch*).

a. n. s., Art. nas. sup.; *a. n. i.*, Art. nas. inf.; *a. t. s.*, *a. t. i.*, A. temp. sup. and inf.; *v. n. s.*, *v. n. i.*, Ven. nas. sup. and inf.; *v. t. s.*, *v. t. i.*, Ven. temp. sup. and inf.; *a. m. e.*, *v. m. e.*, Art. and ven. median; *a. m.*, *v. m.*, Art. and ven. macularis.

amount of pigment contained in the chorioid and in the pigment-epithelium layer of the retina. 1. If there be an abundant supply of pigment in each of these positions, the chorioidal vessels are greatly hidden from view, and the effect is that of a very dark red fundus. 2. If there be but little pigment in the pigment-epithelium layer, the larger chorioidal vessels may be visible, and the

fundus may appear to be divided up into dark islands surrounded by red lines. 3. If the individual be a blonde, there is little pigment either in the pigment-epithelium layer or in the chorioid, and the fundus is seen of a very bright red color, the chorioidal vessels down to their fine ramifications being discernible. In albinos even the chorioidal capillaries may be seen.

The Retinal Vessels.—The arteries are recognized as thin bright red lines running a rather straight course, in the center of each of which is a light-streak. As to the cause of this light-streak there is considerable divergence of opinion. Some attribute it to reflection from the coats of the vessel, or from the surface of the blood column; while others believe that the light is reflected from the fundus through the vessel, which then acts as a very strong cylindrical lens. This light-streak divides the vessel into two red lines. The veins are darker, wider, and more tortuous in their course than the arteries, and, their coats not being so tense, the light-streak is very much fainter.

On reaching the level of the nerve-fiber layer of the retina the central artery and vein divide into a principal upper and lower branch. This first branching often takes place earlier in the vein than in the artery, and the former may even branch before appearing on the papilla, as in Fig. 44. The second branching may take place in the nerve itself; and when this occurs it will appear as though four arteries and four veins sprang from the optic papilla; but more usually this branching occurs on the papilla, as in Fig. 44. The vessels produced by this second branching pass respectively towards the median and temporal side of the retina, and are termed by Magnus the Art. and Ven. nasalis and temporalis sup. and inf. (*vide* Fig. 44). The temporal branches run in a radial direction towards the anterior part of the retina. A small horizontal branch, the Art. and Ven. mediana of Magnus, from the first principal branches is found passing towards the nasal side of the retina. The temporal branches do not run in a horizontal direction, but make a *détour* round the macula lutea, sending fine branches towards the latter. Two or three minute vessels from principal branches run directly from the papilla towards the macula lutea, and around the macula lutea a circle of very fine capillary vessels is formed which cannot be distinguished with the ophthalmoscope; but no vessels run to, or cross over, the fovea centralis itself. The retinal arteries do not anastomose, nor do the larger retinal veins. The small retinal veins have some slight

anastomoses near the ora serrata. Occasionally a vessel emerges near the margin of the disc, usually at the temporal side. It arises from the ciliary vessels, and is hence called a cilio-retinal vessel.

No pulsation of the arteries is observable in the normal eye. In the larger veins near or on the optic papilla, or more usually just at their point of exit, a pulsation may sometimes be seen. This venous pulsation is due to the following sequence of events: Systole of the heart; diastole of, and high tension in, the retinal arteries; consequent increased pressure in the vitreous humor; communication of this to the outside of the walls of the retinal veins, impeding the flow of blood through them, especially in their larger trunks, which offer little resistance, or at their exit from the eye, where they offer the least resistance; and in this way the veins are emptied—the blood gradually coming on from the capillaries overcomes the resistance, and the veins are for a moment refilled. The phenomenon can be most readily observed if the normal tension of the globe be slightly increased by pressure of a finger.

CHAPTER IV.

DISEASES OF THE CONJUNCTIVA.

The Conjunctiva consists of three portions: the *palpebral,* lining the inside of the eyelids; the *bulbar,* covering the sclerotic; and a loose folded portion, uniting these two, which forms the *sulcus* or *fornix,* upper and lower. When the conjunctiva reaches the margin of the cornea it overlaps the latter slightly, and this overlapping portion is known as the *limbus* conjunctivæ, or cornea.

Hyperemia of the Conjunctiva.—In this condition the blood-vessels of the palpebral conjunctiva are especially engaged. Slight chemosis sometimes appears, small vesicles may form, and there may be some swelling of the papillæ and development of lymph follicles. Yet there is not any abnormal discharge from the conjunctiva, and herein lies the chief point of difference between this affection and simple conjunctivitis.

Causes.—Foreign bodies. Foul air, or air loaded with tobacco-smoke. Alcoholic excesses. Accommodative asthenopia. Stenosis lacrimalis, and other forms of lacrimal obstruction. The use of unsuitable spectacles, or the use of the eyes for near work without spectacles when the condition of the accommodation (*e. g.,* hypermetropia, presbyopia) requires them.

Symptoms.—The eyes are irritable. There is lacrimation and photophobia, with hot, burning sensation, and sensations as of a foreign body in the eye, and the eyelids feel heavy. All these symptoms are aggravated in artificial light.

Treatment.—In addition to the removal of the cause, the instillation of a drop of tincture of opium and distilled water in equal parts morning and evening will be found beneficial. The eyes should be protected from the glare of light by dark glasses, and out-of-door exercise is to be recommended.

Conjunctivitis in general.—In addition to hyperemia there is here abnormal secretion. There are several forms of conjunctivitis, the discharge from each being more or less infectious. The secretion from any given form will not, however, always re-

produce that form, but may give rise to another of greater or less severity. Infection takes place by the direct application of the secretion, or also—it is very generally thought—through the air, in which float particles of the infecting substance. This latter mode is especially liable to exist in an ill-ventilated room, where a number of people affected with conjunctival diseases are lodged with others who possess healthy eyes—*e. g.*, in crowded charity-schools. It must be stated, however, that some authorities dispute the possibility of air-borne infection, upon the ground that most of the organisms which cause the various forms of infectious conjunctivitis are non-sporing, and are readily killed by drying. The palpebral conjunctiva is often affected when the bulbar portion remains normal, and the conjunctiva of the lower lid is more frequently attacked than that of the upper lid.

Catarrhal, or **Simple Acute,** or **Muco-purulent Conjunctivitis.**—In mild cases the affection is confined to the palpebral conjunctiva, often even to the conjunctiva of the lower lid; but in the severer cases it extends to the bulbar conjunctiva. Lymph follicles and enlarged papillæ are frequently present, but not necessarily so. There is a sticky, thin, mucous, or muco-purulent secretion, which causes the eyelids to be fastened together on awaking in the morning, and sometimes produces ulceration of the intermarginal portion of the eyelids (intermarginal blepharitis). In some of the very mildest cases this "stickiness" or "gumming" on awaking in the morning is a valuable diagnostic sign, for it is in such cases difficult or impossible to recognize the very slight variation from the healthy appearance of the conjunctiva.

In the severer cases the papillæ are markedly swollen, and may even conceal the Meibomian glands from view. Also one often sees small ecchymoses in the bulbar conjunctiva, especially in certain epidemics; but these have no serious import.

Minute gray infiltrations sometimes form at the margin of the cornea. When there are many of them they may become confluent and form a small gray crescent, which ulcerates, and thus a crescentic marginal ulcer is formed, and very occasionally such an ulcer is followed by iritis.

The catarrh may become chronic, and then the papillæ are more developed, while the blepharitis is liable to extend over to the cutis, causing eversion of the lower punctum lacrimale with resulting lacrimation, and this, in its turn, aggravates the conjunctival affection.

The Koch-Weeks bacillus is the specific bacillus of acute conjunctivitis. It can be cultivated on human serum only, and that with difficulty.[1] A very mild form of acute conjunctivitis is produced by the diplobacillus of Morax,[2] and the pneumococcus is also capable of setting up a mild conjunctivitis.

The Symptoms are those of a severe case of hyperemia (sensations of sand in the eye; hot, burning sensations; weight of the eyelid), with the addition of the annoyance consequent on the secretion, which, by coming across the cornea, may cause momentary clouding of sight. Photophobia is not generally severe unless there be some corneal complication.

Causes.—Draughts of cold air. Contagion. Foul atmosphere. As an epidemic. Foreign bodies. As a sequel of or attendant on, scarlatina, measles, and smallpox.

Diagnosis.—The presence of the gummy secretion distinguishes this affection from mere hyperemia of the conjunctiva. A common mistake amongst those not familiar with eye diseases is to regard a case of iritis as one of simple acute conjunctivitis, the redness of the white of the eye in the former affection being taken for conjunctival hyperemia, etc., and, moreover, a slight secondary conjunctivitis does undoubtedly attend many cases of iritis.

The circumcorneal subconjunctival vessels, which are the episcleral branches of the anterior ciliary vessels, are those which become engorged in iritis, and their engorgement gives rise to a pink or pale violet zone around the cornea, of which the separate vessels cannot be distinctly seen. The conjunctival vessels may be distinguished from the subconjunctival or ciliary vessels by the possibility of moving the former along with the membrane in which they are, by manipulations which can be made with the lower lid of the patient, while these manipulations do not affect the ciliary vessels. The separate conjunctival vessels, too, can be easily distinguished, and they are of a bright red color. The appearance of the iris itself, however, is that upon which the diagnosis finally depends. (See *Iris,* chap. x.)

The Prognosis is good if there be no reason to suspect that the mild form is but the commencement of a more severe inflammation. The infiltrations, and even the ulcers, which sometimes form at the margin of the cornea are not often of serious import, and usually heal according as the treatment restores the conjunctiva to health.

Treatment.—Cold or iced compresses, with the use of a 1 in

5000 solution of sublimate as a lotion, should be used frequently at the first onset, and in mild cases will alone bring about a cure. But the habit, which some patients so readily acquire, of bathing the eyes frequently with cold water should not be permitted, for it is deleterious to the conjunctival affection. When in a day or two the irritation and swelling have somewhat subsided—or from the very commencement, if there be no such irritation—a solution of nitrate of silver, of from 5 to 10 grains to ℥j, should be applied by the surgeon to the palpebral conjunctiva with a camel's-hair pencil, the lid being well everted, and this then should be thoroughly neutralized with a saturated solution of common salt, the whole being finally washed off with plain water. The neutralization with salt water is very important to check prolonged action of the nitrate of silver, as also to obviate conjunctival staining when the treatment is a lengthened one. The application is to be repeated after twenty-four hours, by which time the slight loss of epithelium, the result of the superficial slough, will have been repaired. Immediately after such an application cold sponging or iced compresses are useful, and grateful to the patient. The greatest care is required in the use of nitrate of silver in conjunctival affections for any prolonged period, lest it cause that brownish staining of the membrane called Argyrosis (*ἄργυρος, silver*); thorough neutralization and washing, as above recommended, being the best safeguards. I am opposed to the use even of weak solutions of nitrate of silver as eye-drops to be used at home by the patient, for staining is very apt to be caused in this way.

The application of a 20 per cent. solution of protargol to the conjunctiva with a camel's-hair brush is effective in many cases. It also is liable to cause argyrosis. Ichthargan in a 2 per cent. solution can be similarly used.

Should the surgeon be unable to see the patient daily, the following simple eye-drops are capable of effecting a rapid cure in most cases: ℞ Acid Boracici, gr. v; Zinci Sulph. gr. ii; Tinct. Opii, ℥j; Aq. destill. ad ℥j; one drop in the eye morning and evening, or only once a day in mild cases. Solutions of alum (gr. iv to ℥j of water) and of tannic acid (gr. v to viij to ℥j of water) are often prescribed, but are not so effectual as the foregoing.

A weak boracic acid ointment, to be applied along the margins of the lids at bedtime, is to be ordered. It prevents the gummi-

ness in the morning, which is not only unpleasant to the patient, but is also injurious, by fastening the eyelids together, and thus preventing free drainage of the secretion during the night.

Acute Blennorrhea of the Conjunctiva, or **Purulent Ophthalmia.**—We most commonly find this very dangerous affection either as gonorrheal ophthalmia or as blennorrhea neonatorum.

Etiology.—In gonorrheal ophthalmia the etiological moment is the introduction of some of the specific discharge from the urethra or vagina into the conjunctival sac; while in the latter the infection is believed to take place, either during or just after the passage of the head through the vagina, by an abnormal secretion from the latter finding its way into the infant's eyes. A few instances have been observed of infants born with the disease. Inoculation may also occur a few days after birth by pus conveyed by the fingers of the mother or nurse, or by towels, etc., used for washing the child's face. It is never due to exposure to strong light or to cold, as is popularly supposed.

The more severe cases of blennorrhea neonatorum are caused by a vaginal discharge, which is always gonorrheal. Neisser, who first observed the presence of a peculiar micrococcus in the gonorrheal discharge, also found the gonococcus in the pus from the conjunctiva in cases of gonorrheal ophthalmia, and the same micrococcus has been found in the conjunctival discharge in cases of blennorrhea neonatorum. But the slight cases of the latter affection, which amount to little more than a catarrh of the conjunctiva, may be caused by a vaginal discharge, which is not of the specific gonorrheal nature.

If the infection take place during or immediately after birth, the disease appears from the second to the fifth day, according to the virulence of the secretion. If the inflammation come on later than the fifth day, it may be concluded that the infection was produced by the vaginal discharge being introduced into the eye by the fingers of the mother or nurse, etc. Acute conjunctival blennorrhea also comes about without any assignable cause; but in all such cases it may be regarded as certain that the introduction of some infective pus into the eye has taken place, although without the knowledge of the patient.

Symptoms and Progress.—In mild cases the bulbar conjunctiva may be but little, or not at all, affected, the palpebral conjunctiva alone becoming velvety and discharging a small amount of pus, while there may be no swelling or edema of the eyelids. Such mild

cases are not uncommon in ophthalmia neonatorum, and indeed some of these mild cases cannot be classed as purulent ophthalmia, owing to absence of the gonococcus and presence of the Koch-Weeks and other bacilli. In severe cases of blennorrhea of the conjunctiva there is, soon after the onset, serous infiltration of the palpebral mucous membrane—which consequently becomes tense and shiny—serous chemosis (*χαίνω, to gape open**) of the bulbar conjunctiva, serous discharge, dusky redness and swelling of the eyelids—which make it difficult to evert them—pain in the eyelids, often of a shooting kind, burning sensations in the eye, and photophobia. This first stage lasts from forty-eight hours to four or five days.

Then begins the second stage, in which, owing to swelling of the papillæ, the palpebral conjunctiva becomes less shiny and more velvety; while the discharge alters from serous to the characteristic purulent form, the chemosis, however, remaining unaltered, or becoming more firm and fleshy. The swelling of the lids continues, the upper lid often becoming pendulous and hanging down over the under lid; while, at the same time, it becomes less tense and more easily everted. Gradually the chemosis and swelling of the conjunctiva and eyelids subside, and the discharge lessens, the mucous membrane finally being left in a normal state, unless in a small percentage of cases in which chronic blennorrhea remains. A moderately severe attack of conjunctival blennorrhea lasts from four to six weeks.

Complications with corneal affections form the great source of danger from this affection. They are found chiefly in four different forms: (1) Small epithelial losses of substance on any part of the cornea. If these occur at the height of the inflammation, they are apt to go on to form deep perforating purulent ulcers. (2) The whole cornea becomes opaque (diffusely infiltrated), and towards its center some grayish spots form, which are interstitial abscesses or purulent infiltrations. (3) The infiltration may form at the margin of the cornea, and extend a considerable distance around its circumference, giving rise to a marginal ring ulcer, and, later on, to sloughing of the whole cornea. (4) A clean-cut ulcer may form at the margin of the cornea without any purulent infiltration of the corneal tissue, and may also extend a long way round the cornea. Such ulcers are particularly apt to

* From the appearance produced when the conjunctiva in this condition is much elevated round the margin of the cornea.

occur where there is much chemosis which overlaps the margin of the cornea; and, being hidden in this way, these ulcers are easily overlooked. The chemosis should be pushed aside with a probe, and these peculiar ulcers looked for. They are very liable to perforate.

All the foregoing forms of corneal complications occur both in ophthalmia neonatorum and in gonorrheal ophthalmia. They may appear at any period of the affection, but the earlier they occur the more likely are they to result seriously.

The danger of these ulcers consists in the perforation of the cornea they are apt to produce, of which more later on.

The severer the case, especially the more the bulbar conjunctiva is involved in the process, the more likely is it that corneal complications will arise. For the corneal process is to be regarded as the result of infection by the conjunctival secretion; and this infection is all the more apt to occur where the nutrition of the cornea is impeded by a dense chemotic swelling of the bulbar conjunctiva. Severe chemosis is less common in the blennorrhea of the new-born than in gonorrheal ophthalmia, and this may be the reason for the fact that the latter is much the more dangerous affection of the two.

The Prophylaxis of purulent ophthalmia is a most important matter. It should form part of the routine of lying-in practice. Careful disinfection of the vagina before and during birth, and the most minute care in cleansing the face and eyes of the infant immediately after birth with a non-irritating disinfectant (*e. g.*, a solution of corrosive sublimate 1 in 5000), are to be recommended. The method of the late Dr. Credé has found very general acceptance, and is an admirable one. It is as follows: When, after division of the umbilical cord, the child is in the bath, the eyes are carefully washed with water from a separate vessel, the lids being scrupulously freed, by means of absorbent wool, of all blood, slime, or smeary substance; and then, before the child is dressed, a few drops of a 2 per cent. solution of nitrate of silver are instilled into the eye. Many obstetricians employ this method now as a matter of routine in their lying-in hospitals for all the infants, whether or not it be suspected that there is danger of infection. By its aid Credé reduced the percentage of his cases of ophthalmia neonatorum from 8 or 9 per cent. to 0.5 per cent.

In all cases of gonorrhea it is the duty of the surgeon to explain to his patients what is the danger of their carrying any of the ure-

thral discharge to their eyes; and to charge them to exercise punctilious cleanliness as regards their hands and finger-nails, and care in the use of towels, handkerchiefs, etc.

In respect of *Local Treatment* when the disease has once broken out: In the very commencement of the affection the only local applications admissible are antiseptic lotions (Permanganate of Potash Solution, 1 in 10,000; Sublimate, 1 in 5000) and iced compresses, or Leiter's tubes. With the former the conjunctival sac should be freely washed or irrigated—not syringed—out. In syringing out the conjunctival sac a morsel of the corneal epithelium may be removed, and through this the cornea become infected, and, therefore, this method is objectionable. The iced compresses, or Leiter's tubes, should be kept to the eye for an hour at a time, with a pause of an hour, and so on, or even continuously. In this and in the next stage the chemosis should be freely, and daily, incised with scissors. If the swelling of the lids be great, the external canthus should be divided with a scalpel from without, leaving the conjunctiva uninjured, in order to reduce the tension of the eyelids on the globe, and, by bleeding from the small vessels, to deplete the conjunctiva. Depletion alone can be obtained by leeching at the external canthus, and in many cases is of great benefit at the very commencement. If in adults the chemosis, palpebral swelling, and rapidity of the onset indicate that the inflammation is severe, it is well, in my opinion, to place the patient quickly under the influence of mercury by means of inunctions or small doses of calomel, as by so doing the chemosis is often rapidly brought down, and one source of danger to the cornea removed.

In the second stage (*i. e.*, when the conjunctiva has become velvety, and the discharge purulent) caustic applications are the most trustworthy, and in this respect iodoform and other lauded means cannot compete with them. The application employed may be a solution of nitrate of silver of 15 to 20 grains in ℥j of water, which should be applied by the surgeon to the conjunctiva of the everted lids, and then neutralized with a solution of common salt, as described when discussing the treatment of simple catarrhal conjunctivitis; or the solid mitigated nitrate of silver (one part nitrate of silver, two parts nitrate of potash) may be used, the first application being lightly made in order to test its effect, while careful neutralization with salt water and subsequent washing with fresh water are most important. No remedy is of greater value in puru-

lent ophthalmia than mitigated lapis, when the proper indications for its use are present, and when it is applied with care and intelligence.

The immediate effect of a caustic application to the conjunctiva is the production of a slight slough, under which a serous infiltration takes place. This latter increases and finally throws off the slough, and then the epithelium begins to be re-formed. From the time the slough separates until the epithelium has been regenerated a diminution in the secretion may be noted; but the discharge again increases as soon as the regenerative period is ended, and this now is the moment for a new application of the caustic. From one caustic application of ordinary severity until the end of the regenerative period about twenty-four hours usually elapse. Immediately after a caustic application iced compresses should be used for thirty minutes or longer. Between the caustic applications the pus should be frequently washed away from the eyelids and from between the eyelids with a 4 per cent. solution of boric acid, or with a 1 in 5000 solution of corrosive sublimate, and boric acid ointment should be smeared along the palpebral margins, to prevent them from adhering, and thus retaining the pus.

No corneal complication contra-indicates the active treatment of the conjunctiva by the method just described. Iodoform, finely pulverized, has been much praised as a local application in the second stage of acute blennorrhea of the conjunctiva. It is to be dusted freely on the conjunctiva once or twice a day. For my part I should trust to it in mild cases only.

Permanganate of potash in solution (1 in 10,000) is strongly recommended by Kalt and by Leber[3] as a substitute for nitrate of silver even in severe cases. They state that it can be used from the beginning, and with greater benefit. It is used as a wash four times a day.

When but one eye is affected it is important to protect its fellow from infection by means of a hermetic bandage. This may be made by applying to the eye a piece of lint covered with boracic acid ointment, and over this a pad of borated cotton-wool. Across this, from forehead to cheek and from nose to temporal region, are laid strips of lint soaked in collodion in layers over each other; or a piece of tissue putta-percha may take the place of the lint and collodion, its margins being fastened to the skin by collodion. The shields invented by Maurel and by Buller are serviceable for this purpose.

Treatment of Corneal Complications.—Many surgeons, I understand, use solution of the sulphate of eserin (gr. ij ad aq. ℥j), dropped into the eye as soon as any corneal complication arises, and as long as it continues, on the ground that this drug is believed to have the effect of reducing the intra-ocular tension (a circumstance to be desired in these instances), and also to act as an antiseptic. Its power to reduce the normal intra-ocular tension is not great, and its antiseptic action, if it exist, must be very insignificant, while, in my opinion, it has a decided tendency to promote iritis in these cases, where the iris is so liable to become inflamed secondarily to the corneal process. I therefore do not recommend its use in these cases. I employ atropin here with the object of diminishing the tendency to iritis. Only if a marginal ulcer should perforate, with prolapse or danger of prolapse, into the opening, is eserin indicated, and then simply for the purpose of drawing the iris out of, or away from, the perforation by the contraction of its sphincter.

On the first appearance of an ulcer or infiltration of the cornea, besides the use of atropin, nothing can be done further than the steady continuance of the conjunctival treatment, no remission or relaxation of which is indicated or, indeed, admissible. Greater care is now required in everting the lids, lest pressure on the globe might cause rupture of the ulcer; and it should be remembered that when a case of acute blennorrhea first presents itself, the surgeon, not knowing the condition of the cornea, must use the utmost caution in making his examination, and yet must never fail to get a view of the cornea for the purposes both of prognosis and of treatment. At each visit the cornea must be examined, and it may be found that, as the conjunctival process subsides, any existing corneal affection also progresses towards cure, infiltrations becoming absorbed and ulcers filled up. But even though the conjunctiva be improving, and still more so if it be not, the corneal process may progress, the infiltration becoming an ulcer, and the ulcer becoming gradually deeper, until, finally, it perforates.

Should a corneal ulcer become deep, and seem to threaten to perforate, paracentesis of the floor of the ulcer must be resorted to without delay. By thus forestalling nature a short linear opening is substituted for the circular loss of substance, which would have resulted in the ordinary course of events. Through this small linear opening no prolapse of the iris, or else a relatively

small one, takes place; and, consequently, the ultimate state of the eye is usually a better one than it otherwise would have been. The reduction of the intra-ocular tension after the paracentesis promotes healing of the ulcer. It is often desirable to evacuate the aqueous humor, by opening the little incision in the floor of the ulcer with a blunt probe, on each of the two days after the operation.

If an ulcer perforate spontaneously, the aqueous humor is evacuated, and, unless the ulcer be opposite the pupil and at the same time small in size, the iris must come to be applied to the loss of substance. Should the latter be very small, the iris will simply be stretched over it and pass but little into its lumen, and when healing takes place will be caught in the cicatrix, which is but slightly, or not at all, raised over the surface of the cornea, and the resulting condition is called Anterior Synechia.

If the perforation be larger, a true prolapse of a portion of the iris into the lumen of the ulcer takes place. This prolapse may either act as a plug, filling up the loss of substance and keeping back the contents of the globe, but not protruding over the level of the cornea, or it may bulge out over the corneal surface as a black globular swelling, and may then play the part of a distensor of the opening, causing fresh infiltration of its margins. In either case cicatrization will eventually occur; and if the scar be fairly flat, it is called an Adherent Leukoma, but if it be bulged out, the term Partial Staphyloma of the Cornea is used.

If the perforation be very large, involving the greater part of the cornea with prolapse of the whole iris and closure of the pupil by exudation, the result is a Total Staphyloma of the Cornea. The lens may lie in this staphyloma, or it may retain its normal position, but become shrunken.

The question of the treatment of a recent prolapse of the iris in cases of blennorrheic conjunctivitis is an important one. It has been, and is still largely, the practice to abscise small iris-protrusions down to a level with the cornea, or if large to cut a small bit off their summits, with the object of obtaining flat cicatrices. Horner [4] pointed out that in cases of blennorrhea this proceeding opens a way for purulent infection of the deep parts of the eye, and that serious consequences are not rare. He confined interference with the iris in these eyes to incision of the prolapse, when it seems to be acting as a distensor of the opening, causing fresh infiltration of the cornea. Under other circumstances he restricted

his treatment of the prolapse to the instillation of eserin, which has a marked effect in diminishing the size of the protrusion.

It may occur that on the surgeon's visit to a case of blennorrhea of the conjunctiva he will find the margins of the eyelids gummed together by sero-purulent secretion, while the eyelids are bulged out by the pent-up fluid behind them. The attempt to open the eye should then be very cautiously made, lest some of the retained pus spurt into the surgeon's eye. The surgeon should also be most careful to thoroughly wash and disinfect his hands and nails at the conclusion of his visit.

In cases of blennorrhea neonatorum, when the ulcer has been small, on perforation taking place, the lens, or rather its anterior capsule, comes to be applied to the posterior aspect of the cornea. The pupillary area is soon filled with fibrinous secretion. The opening in the cornea ultimately becoming closed, the iris and lens are pushed back into their places by the aqueous humor which has again collected. Adherent to the anterior capsule on the spot which lay against the cornea is a morsel of fibrin, which gradually becomes absorbed by the aqueous humor. In the meantime changes have been produced by this exudation on the corresponding intracapsular cells, which result in a small, permanent, central opacity at that place, where there is also a slight elevation of pyramidal shape over the level of the capsular surface. This condition is called central capsular cataract, or pyramidal cataract, and rarely results from corneal perforation in adults.

In cases of blennorrhea neonatorum an inflammatory swelling of the joints, so-called gonorrheal arthritis, is very occasionally seen. Deutschmann [5] found the gonococcus in the fluid removed from the joints in two such cases, while other observers found in their cases only the usual pyogenic cocci. Even more rarely do peri- and endocarditis, pleuritis, and meningitis occur.

Follicular Conjunctivitis.—This is catarrhal conjunctivitis, to which is added the presence in the conjunctiva of small round pinkish bodies the size of a pin's head, which disappear completely as the process passes off, leaving the mucous membrane as healthy as they found it. These little bodies are situated chiefly in the lower fornix of the conjunctiva, and may be discovered by eversion of the lower lid, when they will be seen arranged in rows parallel to the margin of the lid. Whether they are easily discovered or not depends on their size and number and on the amount of coexisting hyperemia or chemosis of the conjunc-

tiva. The structure of these bodies shows them to be lymph follicles.

Follicular conjunctivitis is a very tedious affection, lasting often for months. According to Saemisch it is more apt to give rise to marginal ulceration of the cornea than the simple catarrhal form; but I have not observed this to be so. I agree with those who hold that the disease has no clinical relation to trachoma, although some authors regard it as an early stage of the latter.

The Symptoms are much the same as those of catarrhal conjunctivitis. Frequently there is little or no injection of the bulbar conjunctiva, and the chief symptom is asthenopia—an inability to continue near work for any length of time—and much distress in artificial light. Boys and girls from five to fifteen years of age are those most liable to this affection.

Causes.—These are much the same as in simple catarrhal conjunctivitis. The long-continued use either of atropin or of eserin is liable to bring on the disease.

Treatment.—The most useful remedy in this troublesome affection is an ointment of sulphate of copper of from gr. ss to gr. ij in ʒj of vaselin. The weaker ointments should be used at first, and later on the stronger ones if it be found that the eye can bear them. The size of half a pea of the ointment is inserted into the conjunctival sac with a camel's-hair pencil once a day. Eye-drops of equal parts of tincture of opium and distilled water are of use in some cases. Abundance of fresh air, with change from a damp climate or neighborhood to a dry one, is of importance. If the use of a solution of atropin have induced the disease, it should be discontinued; and if a mydriatic be still required, a solution of extract of belladonna (gr. viij ad ʒj) may be employed in its stead.

Spring Catarrh.—The tarsal conjunctiva is invaded by hard flattened bodies of a pinkish color arranged close together, and known as the tesselated or pavement granulations. They are slightly pedunculated. The conjunctiva assumes a milky-like opacity. The bulbar conjunctiva becomes injected, slightly edematous, and at the limbus somewhat elevated with grayish swellings. All these appearances may be present in the same case, or any one (the bulbar appearances, or the pavement granulations, or the milky-like opacity) or two of them may be absent. The margin of the cornea itself is apt to be invaded with minute infiltrations. Very occasionally the cornea becomes seriously

complicated by the growth on the limbus extending over a great portion, or even the entire cornea. There is some mucous or muco-purulent secretion, and the patient complains of the eyelids being stuck together in the morning, of difficulty of using the eyes for near work, of itching and burning sensations, and all these symptoms are increased by exposure to heat. The eyelids droop slightly, giving the patient a sleepy look. The swelling at the limbus is of a fibromatous structure.

The affection makes its appearance with the advent of warm weather in the late spring or early summer, and generally disappears, or is much modified in the cool seasons, to return again with the next warm season, and this is liable to go on for a long series of years.

The Treatment of the majority of these cases yields very unsatisfactory results. Protection glasses should be worn. So far as possible all exposure to great heat of sun or artificial heat should be avoided. Weak astringent collyria, or ointments, may be used; or iodoform ointment (1 in 15), a little put into the eye once a day, or massage twice daily in conjunction with yellow oxid ointment. De Schweinitz recommends boroglycerid locally, and arsenic internally. Antipyrin and quinin internally have proved of use in some cases. For the tarsal granulations Theobald has found marked benefit from use of the roller forceps.

Hay Fever.—This is not uncommon among the better classes in these countries, although it is rarely seen in our hospital outpatient departments. The symptoms, in those liable to it, appear in the early summer each year, and disappear again in the course of six weeks or two months. They consist in catarrh of the nostrils, accompanied by great itching of them and frequent sneezing; while the conjunctiva, especially in the lower fornix, becomes somewhat hyperemic, and there is lacrimation. There is excessive itching of the eyes, which renders the patient most wretched, and forces him to rub his eyes violently. There is photophobia. The respiratory tract may become involved, with some bronchitis and asthma, and general malaise and elevation of temperature are present. Sometimes the eyes alone are affected. There is no tendency to corneal complications.

Treatment is of no avail in preventing the annual recurrence of the affection, nor is it of much use in alleviating the attack. No strong local application should be employed. Weak collyria, or ointments, of sulphate of zinc, or copper, boric acid, or sublimate,

etc., may be tried. Cocain or holocain eye drops (2 per cent.) afford the best relief. Dark glasses should be worn.

Trachoma (*τραχυς, rough*), **Granular Conjunctivitis,** or **Granular Ophthalmia** (also called Egyptian Ophthalmia and Military Ophthalmia).—In this disease, in addition to the usual appearances of simple conjunctivitis, there are developed grayish or pinkish gray bodies about the size of the head of a pin, situated in and close to the fornix conjunctivæ, chiefly of the upper lid, but also disseminated over other parts of the membrane, except that they do not form on the bulbar conjunctiva. These bodies are the trachoma bodies or granulations, or so-called "sago" grains, and in the acute form of the disease they somewhat resemble the follicles of follicular conjunctivitis, but are paler, not so apt to occur in rows, and are more isolated. Microscopically, the trachoma bodies have no capsule, but seem to grow from, or in, the stroma of the conjunctiva. The trachoma bodies consist at first of lymph corpuscles alone; but at a later period this is true of them only towards their surface, while at their bases they are formed chiefly of connective tissue. They are to be regarded as new growths in the conjunctiva.

In addition to the true trachoma bodies, the luxuriant papillary growth, which so often springs up in the chronic cases, is crammed with lymph corpuscles, and these papillæ are termed papillary trachomata, in contradistinction to the true granular trachomata.

No specific micro-organism has as yet been discovered as the cause of trachoma. Leber [6] describes the presence of certain large cells containing peculiarly formed bodies in trachoma, and very probably these have to do with the pathogenesis of the affection.

The disease comes under our notice in two forms—the acute and the chronic. The latter may result from the former, but it is much more common to find it as the primary condition, without any appreciable acute stage having gone before.

Causes.—Both forms are contagious, and probably the infection occurs only by transference of the secretion from one eye to the other by means of fingers, towels, handkerchiefs, etc. Hence the more slovenly in their personal habits, and the more crowded in their dwellings, families, schools, or barracks, nations may be, the more likely is this disease to spread from one individual to another when it once gains a foothold. A great deal, however, remains to be learned as to the manner in which contagion takes

place. For instance, inoculation with discharge from an acute case may give rise only to catarrhal or purulent conjunctivitis, which may recover completely. Again, the infectiousness of chronic cases cannot be very great, for nurses and doctors rarely, if ever, become infected by their patients. Neither do we see trachoma patients infecting other patients in the hospitals in this country, where the disease is so prevalent. Were the infectiousness of the disease very great, even the precautions taken in a well-ordered hospital against contagion would hardly be sufficient to prevent such an occurrence occasionally.

It has been stated that the acute form is often epidemic in places where the hygienic conditions are bad; but in this country I have never seen it as an epidemic, and sporadically not often, although the chronic form is extremely common in Ireland.

Amongst the better classes, both here and elsewhere, the disease is very uncommon. High, dry, mountainous countries are almost free from it, so that, probably, the atmospheric conditions play some part in the etiology.

Some hold that the affection is dependent on constitutional disease, such as scrofula, tuberculosis, syphilis, etc.; but I cannot indorse this view. No doubt many of these patients are anemic and out of health, but this is due to the moping habits they contract, and the little open-air exercise they take in consequence of their semi-blindness.

Acute Trachoma or **Acute Granular Ophthalmia.**—As already stated, this is an affection rarely seen in this country. An attack commences with swelling of the upper lid, great injection of the whole of the bulbar and palpebral conjunctiva, and swelling of the papillæ, with development of the characteristic trachoma bodies. There may be but little discharge; but there is generally much lacrimation, with photophobia, and great pain in the brow and eye. Superficial marginal ulcers of the cornea may form.

The inflammation and papillary swelling increase for a week or so to such a degree that the granulations are hidden from view; and then, taking on a blennorrheic form, the process gradually subsides, until, in the course of two or three weeks longer, the blennorrhea disappears, having brought about absorption of the trachoma bodies, and ultimately the mucous membrane is left in a healthy state.

If, however, in the blennorrheic stage, the inflammation be excessive, the eye may run all the dangers of an attack of acute puru-

lent conjunctivitis; or if, on the other hand, the inflammation be very slight, it may not be sufficient to effect absorption of the granulations, and the process may run into the chronic form.

Egyptian ophthalmia, which is an acute form of trachoma, seems to be a combination of trachomatous disease with purulent ophthalmia, as the gonococcus can be found in the discharge.[7]

Treatment.—It is desirable to abstain from active measures in the commencement of the affection, owing to the tendency to natural cure which is often present, and especially astringents and caustics should be avoided. At the utmost an antiseptic lotion of boric or salicylic acid, and cold applications for relief of the pain and heat are admissible. Dark protection-glasses are agreeable, and, wearing them, the patient should be encouraged to take open-air exercise. But if it be evident that the inflammatory reaction is not active enough, poultices or warm fomentations should be employed to promote it. Once the blennorrheic stage has been reached, great care is required to control it, and if it threaten to exceed safe bounds it must be restrained by means of suitable applications, such as acetate of lead, nitrate of silver, or sulphate of copper in solutions of medium strength; or it may be necessary to use them in strong solutions, or to employ the solid mitigated nitrate of silver.

Chronic Trachoma or **Chronic Granular Ophthalmia.**—The first onset of this disease is often without inflammation, and is then unattended by any distressing symptoms, except that the eye may be more easily irritated by exposure to cold winds, foreign bodies, etc., or more easily wearied by reading and other near work. At this period the conjunctiva will be found free from injection or swelling; but grayish-white semi-transparent trachoma bodies, of the size of a rape-seed and less, will be seen disseminated over the conjunctival surface and protruding from it. Gradually these trachoma bodies or granulations give rise to a more or less active vascular reaction, attended with swelling of the papillæ and purulent discharge—in short, slight blennorrhea. The patients then begin to be more inconvenienced, owing to the discharge which obscures their vision, to sensations of weight in the lids and of foreign bodies in the eye, and to partial ptosis, which is often observable; and this, consequently, is generally the earliest stage at which we see the disease. The enlarged papillæ often grow to a great size, completely hiding the granulations. In this stage the granulations may become absorbed, and the disease undergo cure;

but more commonly it makes further progress. Fresh granulations appear, while the old ones increase in size and undergo a peculiar gelatinous change. They then often become confluent, leaving only here and there an island of vascular mucous membrane. Sometimes the trachoma bodies are very small, and pre-




FIG. 45—(*Saemisch*).

a, Muscle; *b b*, Tarsus having undergone fatty degeneration; *c*, Atrophied Meibomian Gland; *d d*, Hypertrophied Papilla; *e*, Cicatricial Tissue in the conjunctiva; *f*, Tarsus.

sent the appearances of minute white dots, and in this form they are not always easily found.

These chronic granulations consist of lymph cells towards their surface, but towards their bases are formed chiefly of connective tissue. Gradually the cellular elements are transformed into con-

nective tissue, or they may previously undergo the gelatinous change mentioned above, and in this way cicatricial degeneration of the conjunctiva is brought about at each spot where a granulation was seated.

As the disease advances the submucous tissue becomes implicated in the connective tissue alterations, while the tarsus undergoes fatty degeneration and becomes hypertrophied. The granulations disappear, having reduced the conjunctiva more or less to a cicatrix. Contraction of the diseased conjunctiva on the inner surface of the lid causes entropion and distortion of the bulbs of




Fig. 46.

the eyelashes, followed by irregular growth of the latter, with resulting trichiasis and distichiasis. These changes are represented in Fig. 45.

The great danger of granular ophthalmia lies in the complications which may attend it or which follow in its wake; the former consist in pannus, ulcers of the cornea, and severe purulent conjunctivitis, while the latter are the distortions of the lids and eyelashes just referred to.

Pannus (*Lat.*, a cloth rag) presents the appearance (Fig. 46) of a superficial vascularization of the cornea, with more or less diffuse opacity, and often small infiltrations. It invariably commences in the upper portion of the cornea, extending generally over the upper half, and frequently remains confined to this region. But in many cases, at a later stage, it extends to the whole surface of the cornea; and this latter occurrence often takes place almost

suddenly; and the vascularization and opacity sometimes become so intense as to present quite a fleshy appearance, completely hiding the corresponding part of the iris from view. Histologically pannus consists of a new growth, which is extremely rich in cells, and which closely resembles the conjunctiva when occupied with confluent granulations. It is situated between the corneal epithelium and Bowman's layer, and is permeated by vessels derived from the conjunctival vessels. After a length of time Bowman's layer becomes destroyed in places, and then the cellular infiltration gains access to the true cornea, and gives rise to permanent changes in its transparency and curvature. In some bad cases of old-standing pannus the latter undergoes a connective-tissue change. It then becomes smooth on the surface, and the vessels almost disappear, so that the cornea is covered with a thin layer of connective tissue, which obstructs the passage of light and is not capable of cure.

Another result of pannus, sometimes, is a bulging or staphylomatous condition of the cornea, the tissues of which have become so altered that they give way before the normal intra-ocular tension.

A pannus in which as yet there is no connective tissue alteration, and where there is no staphylomatous bulging, is capable of undergoing cure without leaving any opacity behind, except that which may be due to ulcers that have been present.

Pannus is usually a painless affection, but is sometimes accompanied by photophobia and ciliary neuralgia. It may come on at any stage of the disease, and causes defective vision, in proportion to the degree and extent of the opacity. Severe pannus is liable to induce iritis.

The connection between pannus and the condition of the conjunctiva is not altogether evident. It was for long held that the corneal affection is due to mechanical irritation, caused by the rough palpebral conjunctiva; but this view is obviously incorrect, for severe pannus is often seen with a comparatively smooth conjunctiva, while with a truly rough conjunctiva the cornea is frequently perfectly clear. There can now be little doubt that pannus is analogous to the granular disease in the conjunctiva. It is, in fact, the same disease modified by reason of the different tissue in which it is situated, this different tissue being itself a modification of the conjunctiva; and it would seem probable that the cornea becomes diseased by direct inoculation from the conjunctiva of the upper lid. Yet it is remarkable that the bulbar conjunctiva,

lying between the upper margin of the cornea and the fornix of the upper lid, never becomes apparently diseased.

Prognosis.—At any period prior to cicatrization of the conjunctiva an attack of purulent blennorrhea is liable to come on. If not too severe, this may result in a cure by absorption of the trachoma bodies, and should not be checked. If, however, the attack be very severe, the eye runs dangers similar to those of an ordinary attack of purulent conjunctivitis. These are less the more complete and the more intense the pannus.

On the whole, if the disease come under care at an early period, and if treatment be carried out strictly, vision will be retained in a majority of cases, although a radical cure may be difficult or impossible. These cases require to be under constant treatment for long periods, and the very lengthened time, and steady continuous treatment needed for a cure are probably the main obstacles to that cure. In most cases of chronic granular ophthalmia, attendance three times a week for a year will be required, to effect anything that can be called a cure. The common experience is that patients attend for some weeks, and then, being very considerably relieved of their distressing symptoms, and finding their sight vastly improved, they cease attendance long before the disease has been eliminated, to return after a brief interval with a condition of things as bad, if not worse, than before. It is therefore desirable at the very outset of treatment to explain the tedious and dangerous nature of the ailment to each patient.

Treatment.—The aim of this is to bring about absorption of the trachoma bodies with the greatest possible dispatch, in order to prevent the destruction of the mucous membrane to which they tend. No caustic application should be made with the object of directly destroying the trachoma bodies, for this can only be done at the expense of the mucous membrane around them. For chronic cases, with little swelling of the papillæ (blennorrhea), and with little or no cicatrization, the best application is the solid sulphate of copper lightly applied to the conjunctiva, especially at its fornix; but when there is considerable papillary swelling, a 10-grain solution of nitrate of silver, properly neutralized after its application with a solution of salt, or a light application of mitigated lapis, similarly neutralized, is to be preferred. Should there be ulcers on the cornea, or much inflammatory irritation of the eye, sulphate of copper should not be applied to the conjunctiva. An interval of twenty-four hours at least should be

allowed to elapse between each application, whether of sulphate of copper or nitrate of silver, and cold sponging for fifteen minutes should be employed immediately after the application. A change of treatment will be occasionally required even if the remedy first used answer well in the beginning, and one or other of the following can be adopted: Pure carbolic acid liquefied has been used with good result, but I have no experience of it. It is applied with a camel's-hair pencil, and the excess washed off with plain water. Solution of Sublimate, 1 in 2000 to 1 in 1000, applied with lint or cotton-wool to the everted conjunctiva with some pressure and rubbing. Liq. plumb. acetatis dil., never to be used except with everted lids, and washed off with plain water by the surgeon; and not even in this way if there be ulcers of the cornea, as the corneal tissue forming the floor of the ulcer is liable to become impregnated with a white deposit, probably the albuminate of lead, which is by no means easy to remove by




FIG. 47.

operation subsequently. Tannin ointment: Tannin gr. j, to vaselin ʒj, the size of half a pea, to be put into the eye once a day. Sulphate of copper ointment: same strength as the last, and to be used in the same way. Solution of alum: Gr. x to ʒj of distilled water; one drop in the eye once a day. Where an active pannus is present, a drop of solution of atropin should be instilled into the eye once a day as a precaution against iritis.

Some employ scarifications of the conjunctiva when it is much swollen and the papillæ too exuberant; but I have never adopted them, fearing the resulting cicatrices. Brushing of the conjunctiva with a small stiff horsehair brush, or special metallic brush, is a form of scarification used by others, and is sometimes combined with applications of solution of sublimate; and scraping of the conjunctiva with a sharp spoon, with subsequent rubbing in of corrosive sublimate solution, has been recommended.

Again, it has been proposed to excise or abscise the trachoma

bodies; and this may perhaps be allowable if they are isolated and protrude much over the surface of the conjunctiva.

Squeezing out the granulations between the thumb-nails used to be practiced by the late Sir William Wilde, of Dublin, and has recently again come into use. But the proceeding of "expression" is nowadays performed by means of an instrument instead of by the finger-nails. The best instrument for the purpose is Grady's trachoma forceps (Fig. 47). The retro-tarsal fold of the lower or upper lid is grasped as far back as possible by the instrument, compressed and drawn upon, and in this way the trachomatous tissue is squeezed out without laceration of the conjunctiva. The instrument has to be re-inserted and a neighboring part of the conjunctiva treated in the same way, and so on, until the whole conjunctiva of each affected eyelid has been operated on. The four eyelids may be manipulated at one sitting, and the evacuation should be so complete that a repetition of the proceeding will not be required. Particular care should be taken to reach the part of the conjunctiva which is hidden under the commissures. Some cases are immediately and permanently cured by this operation; while others, although greatly benefited, will still require a further routine treatment with local remedies. Expression is indicated only where trachomatous substance can be pressed out. My experience with this method leads me to regard it as a useful one for the acceleration of the cure of some cases of granular ophthalmia before the cicatricial stage has come on.

Excision of the fornix conjunctivæ has been proposed by Schneller,[8] and largely practiced by him and other surgeons. It is claimed for this method that it shortens the treatment of all forms of the disease; that, after it, existing corneal processes undergo rapid cure; that the granular disease in the palpebral conjunctiva, although not directly included in the operation, disappears quickly; that recurrences of the disease are rarer than by other plans of treatment; and that the resulting linear cicatrix has no serious consequence, and is as nothing when compared with the extensive cicatricial degeneration of the whole mucous membrane which the operation is calculated to prevent. Supplemental treatment with the customary local applications is employed until the cure is obtained. I find that this is a useful procedure in some cases.

Infusion of Jequirity (*Arbus precatorius,* Paternoster Bean),

long used in the Brazils, was introduced to the notice of European surgeons by de Wecker. The infusion is made by macerating 154 grains of the decorticized jequirity seeds in 16 oz. of cold water (a 3 per cent. infusion) for twenty-four hours. Twice a day for three days the lids are everted, and the infusion thoroughly rubbed into the conjunctiva with a sponge or bit of lint. The result is a severe conjunctivitis of a somewhat croupous tendency (even the cornea being often hidden by the false membrane), accompanied by great swelling of the eyelids, much pain, and considerable constitutional disturbance, rapid pulse, and temperature of 100°, or more. In the course of eight or ten days the inflammation subsides, and the cornea in many cases will then be found to be free from pannus, or almost so, while complete cure of the granular ophthalmia itself is rarer. Iced compresses to the eyelids should be used during the inflammation. A fresh infusion (not more than seven days old) must be employed in order to secure the best reaction. The majority of surgeons, amongst them myself, find the remedy harmless, if not always successful; but a good many cases are on record where violent diphtheritic conjunctivitis, followed by blennorrhea of the conjunctiva, and by more or less extensive ulceration of the cornea, and even complete loss of the eye, were produced. I have, two or three times, seen a small superficial ulcer form on the lower third of the cornea without further injury. De Wecker regards the presence of a purulent discharge from the conjunctiva as a contra-indication for the remedy, which he finds is then liable to increase the intensity of the blennorrhea in a dangerous degree. Cases where there is little or no papillary swelling, but nearly dry trachoma with pannus, are the most suitable for its use, and I cannot recommend it too highly in these cases. It is marvelous to see the rapid and beautiful cures of the severest pannus by this remedy in properly selected cases. But the presence of well-marked pannus of the cornea without ulceration is, I think, the only thing that can render the employment of jequirity justifiable, and in addition to this the conjunctiva should be free from blennorrhea.

Freshly prepared finely pulverized jequirity beans, from which all oily matter has been removed, may be substituted for the infusion. It is flicked into the eye from a camel's-hair pencil, and is preferred by some to the infusion as being more easily managed in respect of dosage.

Abrin, the active principle of the jequirity bean, has also been

proposed as a local application for trachoma,[9] but it has not come into use owing to its highly toxic properties.

The occurrence of acute dacryocystitis sometimes forms an unpleasant complication of the jequirity treatment even in cases in which the sac was previously quite normal; but I have never myself seen it to occur.

After the subsidence of the jequirity inflammation some of the local remedies above referred to should be regularly applied for the purpose of completing the cure of the conjunctival condition.

Where pannus is present, an occasional drop of atropin should be instilled, in order to control the tendency to iritis.

Besides local remedies, it is of great importance that the hygienic surroundings of patients suffering from granular ophthalmia be seen to, and that they be obliged to spend a considerable time daily in the open air.

If the upper lid be tightly pressed on the globe, as it sometimes is, the physiological pressure varying in different individuals, an impediment is offered to the cure by any method, and pannus is promoted. It is then necessary to relieve the pressure by a canthoplastic operation. (See chap. vii.)

Peritomy.—This procedure is adopted for the cure of pannus by destruction of the vessels which supply it, and is as follows: About 5 mm. from the margin of the cornea an incision is made in the conjunctiva with scissors, and carried at this distance all the way round the cornea. This ring of conjunctival tissue is then separated up from the sclerotic, and cut off at the corneal margin, and the underlying connective tissue is dissected off the corresponding portion of the sclerotic, which is thus laid quite bare. The proceeding is not always satisfactory, and of late years I have practiced it but little.

Lymphoma of the Conjunctiva.—Under this heading cases have been described which present the appearances, at first sight, of acute granular ophthalmia; but the "granulations," which are enormous in size, attack both lids, and are associated with enlarged lymphomatous masses in the neck, which do not lead to ulceration or scarring. The conjunctival affection runs a rapid and favorable course, without any cicatricial contraction.

Croupous Conjunctivitis.—This is a disease of early childhood, and is not common. The palpebral conjunctiva is a good deal swollen, and is covered with a false membrane, that may be peeled off, leaving a mucous surface underneath, which bleeds little or

not at all. The disease is not a severe one, and does not cause secondary corneal affections, unless when the bulbar conjunctiva, as it very rarely does, participates in the attack. It must not be mistaken for diphtheritic conjunctivitis, from which it is readily distinguished by the ease with which in it the false membrane can be removed, and by the vascular condition of the underlying mucous membrane.

This is usually regarded as nothing more than a severe form of catarrhal conjunctivitis, in which the secretion happens to be rich in fibrin, and hence possessed of a marked tendency to coagulate on the surface of the conjunctiva. But the presence of the Klebs-Löffler bacillus has been demonstrated in cases of croupous conjunctivitis, which ran a favorable course, and hence there seems to be a bacteriological relationship between this form of conjunctivitis and diphtheritic conjunctivitis, which are so different from each other clinically.

Causes.—Contagion, Epidemic.

Treatment.—Iced compresses or Leiter's tubes to the eyelids during the croupous stage, with antiseptic cleansing of the conjunctival sac (Sol. Hydrarg. Perchlor. 1 in 5000, or Sol. Acid. Borac. 4 per cent.). No caustic should be used in this stage, as it is apt to produce corneal changes. Sulphate of quinin sprinkled on the conjunctiva is praised by some surgeons as a useful application at this period. When the false membrane ceases to be formed a slight blennorrhea comes on; and this is to be treated with nitrate of silver applications in the usual way.

Diphtheritic Conjunctivitis.—There is no more serious ocular disease than this, for it may destroy the eye in twenty-four hours; while in severe cases treatment is almost powerless. Forutnately it is almost unknown in these countries, while in Berlin it used to be so frequent that von Graefe set apart two wards for it in his hospital, which were under my care as his assistant. It is now a much less common disease there, owing probably to the improved hygiene of the city.

The subjective symptoms of its initial stage are similar, although severer, especially in the matter of pain, to those of blennorheic conjunctivitis. The objective symptoms differ from those of blennorrhea, in that the lids are excessively stiff, owing to plastic infiltration of the subepithelial and deeper layers of the conjunctiva, while the surface of the mucous membrane is smooth, and of a grayish or pale bluff color. If an attempt be made to peel off some

of the superficial exudation the surface underneath will be found of the same gray color, not red and vascular, as in croupous conjunctivitis. This stage of infiltration lasts from six to ten days, and constitutes the period of greatest peril to the eye; for while it lasts the nutrition of the cornea must suffer, and sloughing of that organ is extremely apt to take place. Towards the close of the first stage the fibrinous infiltration is eliminated from the eyelids, and the conjunctiva gradually assumes a red and succulent appearance, and at the same time a purulent discharge is established. This constitutes the second or blennorrheic stage. A third stage is formed by cicatricial alterations in the mucous membrane, which often lead to symblepharon, or to xerophthalmos; so that, even if the eye escape corneal dangers in the first and second stages, others almost as serious may await it in the final stage.

Corneal complications are most likely to occur in the first stage, and are then also most likely to prove destructive to the eye. The earlier they appear the more dangerous are they. If the blennorrheic stage come on before corneal complications appear, or even before an ulcer contracted in the first stage has advanced far, they are more easily controlled.

Causes.—It is difficult to assign a cause for this disease, which chiefly attacks children. It is frequently epidemic, is extremely infectious, and, although similar in its nature, is rarely, if ever, found in connection with an attack of diphtheritis of the fauces. The Klebs-Löffler bacillus is always present in the conjunctival discharge.

Treatment.—Injections of the anti-toxin serum are the sovereign remedy, but local applications are also of use. In the first stage frequent warm fomentations, with antiseptic cleansing, are the only local measures admissible. No caustic or astringent application should be used. Internally, the patient should be treated with iron and quinin and generous diet. In the second or blenorrheic stage careful caustic applications are to be used. Corneal ulcers must be dealt with, whenever they arise, in the same way as though the case were one of blennorrheic conjunctivitis. When the purulent discharge ceases, solutions of soda, milk, or glycerin may be prescribed as lotions for the conjunctiva, to arrest, if possible, the xerophthalmos.

Conjunctival Complication of Smallpox.—Of this I have, fortunately, too little experience to enable me to speak authoritatively.

The following embodies the views of the late Professor Horner,[4] who studied the subject during an epidemic in 1871. A good deal of uncertainty prevailed previously, for the initial stages of the eye affection were not carefully observed by physicians, owing to the swelling of the eyelids, while the ophthalmologist saw only the results of the process in the period of convalescence.

Smallpox pustules on the cornea are, Horner believed, extremely rare; indeed, he saw but one such case. The most frequent and most serious mode of attack consists in a grayish-yellow infiltration in the conjunctiva close to the lower margin of the cornea, not extending to the fornix conjunctivæ, nor far along the inner or outer margin of the cornea. It occurs in the eruptive stage, and is to be regarded clinically as a variola pustule. This infiltration or pustule gives rise to a corneal affection, as does a solitary marginal phlyctenula, either in the form of a marginal ulcer or as a deep purulent infiltration, ulcerating, perforating, leading to staphyloma, purulent irido-chorioiditis and panophthalmitis—results which are often first observed long after the primary conjunctival affection has disappeared.

Horner believed that the germ of the conjunctival infiltration makes its way between the eyelids, and that the constancy of the position of the infiltration below the cornea is accounted for by this theory, that part of the conjunctiva, with closed eyelids and eyeball consequently rotated upwards, being the most exposed to particles entering.

Treatment.—On this ground he recommended the prophylactic use of boracic acid ointment on lint applied over the eyelids. If a conjunctival pustule have already formed without any, or only commencing, corneal affection, he would destroy the pustule with fresh chlorin water, or with mitigated lapis carefully neutralized. Corneal complications are treated as in blennorrhea of the conjunctiva or diphtheritis.

The frequency with which the eyes become affected varies in different epidemics.

As true post-variolous eye-affections, Horner recognized diffuse keratitis, iritis, and iridocyclitis, with opacities in the vitreous humor, and glaucoma; in the hemorrhagic form of the disease, hemorrhages in the conjunctiva and retina; and, where pyemic poisoning comes on, septic affections of the chorioid and of the retina take place.

Amyloid Degeneration.—This rare disease attacks chiefly the

palpebral conjunctiva, but is also seen in the bulbar portion. It causes great tumefaction of the affected lid, without any inflammatory symptoms. The eyelid can be but partially elevated, and is often so stiff and hard that it can be everted only with difficulty. The conjunctiva has the appearance of white wax. The disease ultimately extends to the tarsus, but is a strictly localized process, and not associated with amyloid disease in any other part of the system. It sometimes seems to be developed from granular ophthalmia, but occurs also as a primary disease. The positive diagnosis can be made by submitting a small portion of the diseased conjunctiva to the iodin test.

Hyaline Degeneration of the conjunctiva has also been observed. It cannot clinically be distinguished from Amyloid Degeneration, and is really an early state of the latter condition.

Treatment consists in the removal of the diseased parts, by the knife and scraping, so far as may be possible.

Tubercular Disease of the Conjunctiva.—This is an extremely rare disease. It usually commences in the palpebral conjunctiva of the upper lid, and very rarely in the bulbar conjunctiva, in the form of small round yellowish-gray nodules, which soon ulcerate. The margins of these ulcers are well defined, and their floors either of a yellowish lardaceous appearance, or covered with grayish-red granulations. The surrounding conjunctiva is swollen, and if the palpebral conjunctiva be much involved the lid becomes enlarged in every dimension, and the ulcerative process may soon destroy part of the lid. It may also extend to the bulbar conjunctiva, and the cornea may become covered with pannus. The preauricular and submaxillary glands usually become enlarged. The positive diagnosis of the nature of the disease should be made by the examination of portions of the floor of the ulcer for the characteristic tubercle bacillus, which will distinguish this from secondary syphilitic ulceration of the conjunctiva, between which and the tubercular ulceration there is sometimes a resemblance. The appearance may also be suggestive of trachoma, or even of a malignant growth. Tubercular conjunctival disease is usually unattended by pain, or there is only a slight burning sensation; but, again, when the ulceration is extensive, severe pain may set in.

The disease is very chronic, its progress sometimes extending over many years. It is rarely met with except in youth. Some of those whose eyes are attacked are already the subjects of tuber-

culosis in other organs, but very many of them are perfectly healthy in that respect. In fact, we have reason to believe that tuberculosis of the conjunctiva is much more often a primary disease, the result of an ectogenic infection, even in cases where already tuberculosis exists elsewhere, than of infection occurring through the blood. Tubercle bacilli introduced into the normal conjunctival sac have, it is true, been found to be harmless, for the intact epithelium offers an insuperable obstacle to their entrance into the tissue. But a superficial loss of substance of the conjunctiva is sufficient to allow of its inoculation with the bacilli, and then the disease becomes established. The frequent lodgment of foreign bodies under the upper lid explains why this is the most common place for the disease to begin in. But although conjunctival tubercular disease is not often secondary to tubercular disease in other parts of the system, yet it is itself liable to be the starting-point of general tuberculosis.

Treatment.—The fact last mentioned makes it most important, in cases of primary tubercular disease of the conjunctiva, to thoroughly eradicate the diseased focus so as to avert an infection of other organs, and this can often be effected. If the ulcers be not already too extensive they must be scraped, and the actual cautery freely applied to them; and, where the disease has already spread to the cornea, sclerotic, iris, or chorioid, enucleation of the eyeball is imperatively called for.

Lupus of the conjunctiva usually occurs as an extension of the disease from the surrounding skin, or rarely from the lacrimal sac, as in a case of Dr. Louis Werner's, where the disease extended from the mucous membrane of the nose, through both lacrimal sacs, to the inferior palpebral conjunctiva. It is seen as a patch or patches of ulceration, covered with small dark-red protuberances or granulations, chiefly on the palpebral conjunctiva, which bleed easily on being touched.

Like lupus of the skin, these ulcerations undergo spontaneous healing and cicatrization in one place (unlike tubercular ulceration in that respect), while they are still creeping over the surface in another direction. But we now know that lupus, wherever it occurs, is really a tubercular disease, and that the two forms differ only in their clinical aspect.

The Treatment is scraping with a sharp spoon and the application of the actual cautery.

Pemphigus of the Conjunctiva.—This is another rare disease.

It has been seen in connection with pemphigus vulgaris of other parts of the body, but it also occurs as an independent disease. It is attended by attacks of much pain, photophobia, and lacrimation; and the conjunctiva, at each place where subconjunctival exudation of serum has been situated, undergoes degeneration and cicatricial contraction. Such attacks succeed each other at shorter or longer intervals, for weeks, months, or years, until, finally, the entire conjunctiva of each eye may have become destroyed, and the eyelids are adherent to the eyeball. The cornea gradually becomes completely opaque, or, having ulcerated, becomes staphylomatous. In the course of the disease the eyelashes are apt to become turned in on the eyeball, or even entropion may form; and these conditions aggravate the suffering of the patient.

The foregoing is a description of a severe case. In less severe cases the conjunctiva may not be completely destroyed, and the cornea may not be affected.

The formation of a true bulla hardly ever occurs, for the conjunctival epithelium is so delicate that it cannot be disturbed in this way by the serous exudation beneath it, but rather breaks down at once. Consequently, the conjunctival surface is found in these cases to be covered by what looks like a membranous deposit, upon removal of which a raw surface is exposed; and these appearances have led to the mistaken diagnosis of croupous and of diphtheritic conjunctivitis.

Treatment is helpless in respect of arresting the progress of the disease, or of restoring sight when lost in consequence of it. The most one can do is to relieve the distressing symptoms by emollients to the conjunctiva, and by the use of closely fitting goggles, to protect from wind, dust, and sun. Internally, arsenic is indicated.

Xerosis (*ξηρός, dry*), or **Xerophthalmos** is a dry, lusterless condition of the conjunctiva, associated in the severer forms with shrinking of the membrane. There are two forms of the affection —the parenchymatous and the epithelial.

In Parenchymatous Xerophthalmos there is a more or less extensive cicatricial degeneration of the conjunctiva, dependent upon changes in its deeper layers, while its surface and that of the cornea become dry, and the latter becomes opaque, and the eye consequently sightless. The conjunctiva shrinks so completely in many of these cases that both lids are found adherent in their whole extent to the eyeball, which is exposed merely at the

palpebral fissure, where the opaque and lustreless cornea is to be seen. From what remains of the conjunctiva, scales, composed of dry epithelium, fat, etc., peel away. The motions of the eyeball are restricted in proportion to the extent of the conjunctival degeneration. There is no cure for this condition.

Fig. 48 represents a case of xerophthalmos, the result of pemphigus, which was under my care at the Royal Victoria Eye and Ear




Fig. 48.

Hospital. Here the eyelids were not wholly adherent to the eyeball, and the cornea remained clear.

The Causes of parenchymatous xerosis of the conjunctiva are granular ophthalmia, diphtheritic ophthalmia, pemphigus, and the condition is said to be very occasionally seen as a primary disease, described as essential shrinking of the conjunctiva. Many observers altogether deny the existence of the primary affection, and maintain that the cases described as being of that nature are merely the result of pemphigus, and I am inclined to agree with this view.

Treatment.—As cure is impossible in this form of xerophthalmos, the only indication is to afford relief, so far as it can be done,

from the distressing sensations of dryness of the eyes which are complained of. The best applications are milk, glycerin, olive oil, and weak alkaline solutions, and the eyes should be protected from all irritating influences by protection goggles.

Epithelial Xerosis of the conjunctiva is confined to the epithelium of that part of the conjunctiva which covers the exposed portion of the sclerotic in the palpebral opening. It there becomes dry and dull and covered with small white spots; while the whole bulbar conjunctiva is loose, and easily thrown into folds by motions of the eyeball, and there may be a good deal of secretion. This form of xerophthalmos often occurs in epidemics, but also sporadically, accompanied, oddly enough, by night-blindness (the light-sense unimpaired) and contraction of the field of vision. The combined condition has been noticed chiefly in persons of debilitated constitution, who have been exposed to strong glares of light, and is said to have appeared in epidemics, under these conditions, in foreign prisons and barracks.

Treatment by rest, protection from glare of light, nutritious diet, and tonics invariably restore the eyes to their normal functions.

Again, epithelial xerosis occurs in very young children in connection with a destructive ulceration of the cornea (see Infantile Ulceration of the Cornea with Xerosis of the Conjunctiva, chap. vi.).

Pterygium (*πτέρυξ, a wing*).—This is a vascularized thickening of the conjunctiva, triangular in shape, situated most usually to the inside of the cornea, sometimes to its outer side, and rarely either above or below it. The apex of the triangle, the head of the pterygium, is on the cornea; and its base, the body, at the semilunar fold. The neck of the pterygium is that part of it at the margin of the cornea. There is frequently, but not always, a tendency of the growth to advance into the cornea, of which it seldom reaches the center, and still more rarely extends quite across it.

In its early growth the pterygium is somewhat thick and succulent-looking, and very vascular. But finally it ceases to grow, and then becomes thin and pale, and this is its retrogressive stage; yet it never entirely disappears. Sight is not affected unless the pterygium extends over the pupillary region of the cornea. A limitation of the motion of the eye to the other side, and consequent diplopia, is sometimes caused by a pterygium; but, for the

most part, the disfigurement alone it is which brings these cases to the surgeon.

Causes.—The starting-point of a pterygium is often an ulcer at the margin of the cornea, which in healing catches a morsel of the limbus conjunctivæ and draws it towards the cicatrix, throwing the mucous membrane into a triangular fold. The ulcer then forms anew in the cornea immediately inside the cicatrix, and, in healing, the point of conjunctiva is drawn into it again, and is carried a little farther into the cornea, and so on. The hollow lying between a pinguecula (see below) and the margin of the cornea is apt to lodge small foreign bodies, which cause shallow marginal ulcers, and these, in healing, draw the pinguecula over on the cornea. A marginal ulcer in phlyctenular keratitis, or in acute blennorrhea, may serve the same end. The only objection to this theory of the causation of pterygium is that an ulcer is not always to be found at the head of the growth.

Fuchs [10] believes that pterygium develops from the pinguecula, and that the latter causes nutritive changes in the cornea, loosening the superficial lamellæ, and allowing the connective tissue of the limbus to grow in on the cornea.

Pterygium is a rare affection in this country, but is more common in countries or localities where the air is filled with fine sand or other minute particles.

Treatment.—Unless the pterygium be very thick, and have invaded the cornea to some extent, or be progressing over the cornea, it is well to let it alone; the more so as by removing it a quite normal appearance is not given to the eye, for a mark is necessarily left both on cornea and conjunctiva. If it be progressive or very disfiguring, it should be removed, other proposed modes of dealing with it being futile. This may be effected either by ligature or excision.

In the method by ligature a strong silk suture is passed through two needles. The pterygium being raised with a forceps close to the cornea, one needle is passed under it here and the other needle in the same way close to its base, the ligature being drawn halfway through. The thread is cut close behind each needle, thus forming three ligatures, which are respectively tied tight. In four or five days the pterygium comes away.

For excision the apex is seized with a forceps and dissected off either with a scissors or fine scalpel, care being taken not to injure the true cornea; or a good plan is to pass a strabismus hook under

the pterygium when raised up from the sclerotic, and to forcibly separate the corneal portion by drawing the hook under it. The dissection is continued towards the base of the pterygium, where it is finished with two convergent incisions meeting at the base. The mucous membrane in the neighborhood of the base is separated up somewhat from the sclerotic, and the margins of the conjunctival wound are then carefully brought together with sutures. Skin grafts, according to Thiersch's method, have been used with success to cover the defect.

Pinguecula (pinguis, *fat*) is the name given to a small yellowish elevation in the conjunctiva near the margin of the cornea, usually at its inner side, more rarely at its temporal margin, but sometimes in each place. It contains, notwithstanding its name, no fat, but is composed of connective tissue and elastic fibers. It is supposed to be due to the irritation caused by small foreign bodies. It rarely grows to a large size, and requires no treatment unless it becomes very disfiguring, when it may be removed with forceps and scissors.

Subconjunctival Ecchymosis.—The rupture of a small subconjunctival vessel in the bulbar conjunctiva, without conjunctivitis, is of frequent occurrence. It suddenly gives a more or less extensive purple hue to the "white of the eye," causing the patient much concern. It is common enough in old people, but may occur in the young, and even in children, from severe straining, as in whooping-cough, vomiting, or raising heavy weights. It is occasionally significant of diabetes. It also occurs sometimes during epileptic fits, and profuse subconjunctival hemorrhage is occasionally found in cases of fracture of the base of the skull, having made its way along the floor of the orbit. It is of no importance so far as the integrity of the eye is concerned.

Treatment.—None is required, the extravasated blood gradually becoming absorbed.

Nevus of the conjunctiva may occur along with the same condition of the lids, but it also occurs separately, especially on the plica or caruncle.

Treatment.—Electrolysis or ligature. Good results have been obtained by Snell [11] with ethylate of sodium carefully painted on.

Polypus of the conjunctiva, for which it is difficult to assign a cause, is sometimes seen. It is generally small, in connection with the semilunar fold or caruncle, and can readily be removed

with the scissors. Granulations occurring after tenotomy for strabismus are sometimes, and incorrectly, called polypi.

Dermoid Tumors.—These are pale yellow in color, and in size from that of a split pea to that of a cherry. They are smooth on the surface, and sometimes have fine hairs, and sit usually at the outer and lower margin of the cornea; but Fig. 49 was drawn from a case on which I operated, where the dermoid was situated on the inner side of the cornea, extending over somewhat on the latter, and not at the most usual seat. In structure they resemble that of the skin. They are congenital tumors, supposed to be due to an arrest in development, but they often have a tendency to extend over the cornea. If this tendency be present, the tumor




Fig. 49.

must be removed by dissecting it off the cornea, care being taken not to go into the deep layers of the latter.

Lipoma occurs as a congenital subconjunctival formation of fat, usually situated between the superior and external recti muscles.

Syphilitic Disease of the Conjunctiva occurs both as primary and as secondary disease. It will be treated of in chap. vi. on Diseases of the Eyelids.

Papilloma, or **Papillary Fibroma.**—This is a non-malignant growth, which may spring from any part of the conjunctival sac. It appears in the beginning as a small round red knob. The papillomata growing from the tarsal conjunctiva and from the semilunar fold frequently take on a cauliflower appearance; while on the bulbar conjunctiva and in the fornix the growths are liable to be pedunculated, with a papillary surface. The limbus of the conjunctiva is a favorite seat for a papilloma, and in the early stage

it may be impossible to distinguish it from an epithelioma. But if the case come under observation at a later stage, when the growth has overlapped the cornea, this difficulty does not arise; for the papilloma merely lies on the cornea, and can be lifted freely off it with a probe, while epithelioma infiltrates the corneal tissue.

Treatment.—Thorough removal with knife or scissors, and actual cautery, as otherwise the growth is liable to recur.

Epithelioma is not common as a primary disease of the conjunctiva. When it is so found it is seen as a little non-pigmented tumor growing from the limbus of the conjunctiva, surrounded by vascularization, and may in this stage be mistaken for a phlyctenula—of which, however, the margins are not so steep—or for a papilloma (*vide supra*). As the tumor increases in size it becomes lobulated, and ulcerates, and soon attacks the cornea, giving rise on the latter to an appearance very like pannus. The neighboring lymphatic glands become enlarged.

Sarcoma, too, is rare. It likewise most commonly takes its origin in the limbus conjunctivæ, and is then usually a pigmented tumor, a melanosarcoma. That these sarcomata are pigmented is explained by the fact that the limbus contains pigment, although so slight in amount as not to be visible to the naked eye. The growth spreads backwards over the sclerotic, with which it becomes loosely adherent. It does not attack the cornea so readily as the epitheliomatous growths, although it often overlaps the surface of the cornea. In some cases it burrows under Bowman's layer, and gradually into the deep layers of the cornea, and by this path reaches the interior of the eyeball. In its later stages this tumor grows to an enormous size.

But conjunctival sarcoma also starts from other parts of the conjunctiva, and in a case of my own I four times removed sarcomatous tumors from different parts of the fornix, an interval of some months elapsing between the appearance of each small tumor.

Treatment.—Both epithelioma and sarcoma of the conjunctiva demand prompt operative removal in order to prevent an extension of the growth to the rest of the eye, if the case be seen early, as well as to avert metastases to other organs. The knife and actual cautery may save the eye, and the life in the early stages, but later on removal of the whole eye is often called for.

Simple Cysts of the conjunctiva are very rare. They appear

as clearly spherical protuberances of about the size of a pea, seated usually on the bulbar conjunctiva. The walls of the cysts contain but few vessels, are thin, and almost transparent; while for contents they have a clear limpid fluid. These cysts cannot as a rule be moved from their position, because they are adherent to the conjunctiva, which indeed takes part in the formation of their walls. They are, very probably, dilated lymphatic vessels. Small beadlike strings of dilated lymphatics are very frequently seen on the bulbar conjunctiva.

These simple cysts are most commonly congenital, but they may begin to be developed during life.

Treatment.—The cyst may be dissected out, or it may be sufficient to abscise its anterior wall.

Subconjunctival Cysticercus is a little more common than simple cyst of the conjunctiva, and yet only about fifty examples of it have been recorded.

Cysticerus is distinguished from simple cyst by its free mobility under the conjunctiva, to which it is not attached; by its thicker and more vascular walls; and, above all, by the presence of a round, white, opaque spot on the anterior surface, first pointed out by Sichel, and looked on by him as pathognomonic of a cysticercus. This spot indicates the position of the receptaculum; and occasionally, when this comes to be placed on the posterior surface of the cyst, it may be difficult, or impossible, to make the diagnosis with certainty.

Treatment.—The cyst may be pushed to one side under the conjunctiva, an incision made in the latter, the cyst then pushed back again, and out through the opening.

Lithiasis consists in the calcification of the secretion of the Meibomian glands, which are seen as little brilliantly white spots not larger than a pin's head in the conjunctiva. There may be one only or very many. These concretions often give rise to much conjunctival irritation, and if they protrude over the surface of the conjunctiva may injure the cornea. Each one—the eye having been cocainized—must be separately removed by a needle, with which first an incision has been made into the conjunctiva over the concretion.

Conjunctivitis Petrificans.—Under this title Leber [12] has described a rare and remarkable disease of the conjunctiva. In the course of a brief period, and accompanied by some slight inflammatory reaction, a stony hard, white, chalky substance is deposited,

in more or less extensive patches, in the previously healthy conjunctiva, the deposit being scarcely raised over the conjunctival surface. The disease attacks a part of the bulbar or palpebral conjunctiva, and may extend to the intermarginal portion of the eyelid. One or both eyes may be attacked. After a time, which varies from a week to several months, the deposit is thrown off or absorbed, and the affected part suffers either no detriment or there may be slight thickening and shrinking. There is no great tendency to corneal complications, but slight marginal ulcerations, which heal readily, occasionally occur. In one case severe diffuse opacity of the cornea seriously affected the sight. Frequent relapses are liable to take place in the same or in different parts of the conjunctiva, and the whole course of the affection may extend over several years, and then end in complete cure.

No cause has as yet been assigned for this disease, although Leber suspects it to be an ectogenic microbic infection. The only treatment which seems to be of use is warm fomentations, and the careful operative removal from time to time of the chalky scales as they become loosened from the main mass.

Uric Acid Deposits have been seen in the palpebral conjunctiva in gouty cases.

Injuries of the Conjunctiva.—Foreign bodies frequently make their way into the conjunctival sac, and cause much pain, especially if they get under the upper lid, by reason, chiefly, of their coming in contact with the corneal surface during motions of the lid and of the eye. If the foreign body be under the lower lid it will be easily found on drawing down the latter, and, provided it be not actually embedded in the mucous membrane, it is easily removed with a camel's-hair pencil or with the corner of a soft pocket-handkerchief; but if the foreign body be under the upper lid it is necessary to evert the latter before it is reached. Should the foreign body be embedded in the conjunctiva it must be pricked out of its position with the point of a needle or other suitable instrument.

The conjunctiva is frequently injured in severe wounds of the eyelids or eyeball. The interest and treatment are centered here chiefly on the other more important parts, which have been injured. A tear or wound of the conjunctiva (usually of the bulbar portion), when it occasionally occurs without injury to other parts, is in general of very slight moment. If the wound be extensive its edges should be drawn together with a few points of suture;

but otherwise healing will take place with the aid simply of a bandage to keep the eye closed for a few days.

A common form of injury, which may involve the conjunctiva alone, is a burn by acid or lime. In the case of a strong acid getting into the eye, if the patient be seen immediately after the occurrence, the whole conjunctival sac should be well washed out with an alkaline solution (1 per cent. soda solution). In the case of lime, after all the particles have been most carefully removed from the eye with forceps, a weak solution of a mineral acid may be used for washing out the conjunctival sac; or, as is recommended by some, a solution of sugar as thick as syrup may be poured into the eye. Later, olive or castor oil, or even butter, may be applied, the subsequent treatment being continued with weak sublimate ointment. Cocain may be employed to relieve the pain.

In the case of a severe burn of the conjunctiva, the resulting cicatrix is liable to produce a more or less extensive union of the eyelid to the eyeball (Symblepharon), which often interferes with the motion of the latter, or even with vision, if the cornea be obscured. No measures taken during the healing process can prevent symblepharon if the degree of the burn be such as to bring it about. The relief of symblepharon by operation will be dealt with in chap. vi., on Diseases of the Eyelids.

References.

[1] "Von Graefe's Archiv," xlviii. p. 639.
[2] "Annales d'Oculistique," Janvier, 1897.
[3] "Bericht der ophthalmologischen Gesellschaft." Heidelberg, 1897, p. 249.
[4] Gerhardt's "Handbuch der Kinderkrankheiten," V. Part 2.
[5] "Archiv für Ophthalmologie," xxxvi. 1, p. 109.
[6] "Bericht der ophthalmologischen Gesellschaft." Heidelberg, 1896, p. 156.
[7] "Annales d'Oculistique," 1894, p. 19.
[8] "Von Graefe's Archiv," xxx. 4. p. 131; and xxxiii. p. 113.
[9] *Ibidem*, lii. i. p. 72.
[10] *Ibidem*, xxxviii. 2. p. 1.
[11] "Trans. Ophthal. Society of the United Kingdom," xiii. p. 39.
[12] "Bericht der ophthalmologischen Gesellschaft." Heidelberg, 1895, p. 46; and "Von Graefe's Archiv," li. i. p. 1.

CHAPTER V.

PHLYCTENULAR, OR STRUMOUS, CONJUNCTIVITIS KERATITIS.*

Both from a clinical and nosological point of view it would be incorrect to divide this affection into two, under the heads of Diseases of the Conjunctiva and Diseases of the Cornea; and therefore I treat of it here as one disease, and, being a very important disease, I devote a special chapter to it. It is important, because it is excessively common, and because it is capable of causing considerable damage to sight. Moreover, even when it occurs on the cornea, it might, strictly speaking, be regarded as a conjunctival disease, for that corneal layer, which it primarily attacks, is the epithelium, and this—and probably also Bowman's membrane and the anterior layers of the true cornea—as we know from the fetal development of the membrane, is a continuation of the conjunctiva in a modified form over the cornea.†

The disease is characterized by the eruption of phlyctenulæ (*φλύκταινα*, a *vesicle,* or *pustule*) on the conjunctiva bulbi, on the conjunctival limbus, or on the cornea, and, if the upper lid be everted, small phlyctenular ulcers will not rarely be found on the tarsal conjunctiva. It is chiefly a disease of children up to the eighth or tenth year of age.

Notwithstanding the derivation of the word, a phlyctenula, or phlyctene, is originally neither a vesicle nor a pustule. It is a formation *sui generis,* and, when on the conjunctiva, is a solid elevation consisting of a collection of lymph cells, and is of a grayish color. In a late stage the phlyctenula, especially on the cornea, may become a pustule by infection. On the conjunctiva two types of the disease can be recognized:

1 **The Solitary,** or **Simple, Phlyctenula.**—Of this there

* *κέρας, a horn.*

† The posterior epithelium—or, according to some, this along with the membrane of Descemet and the posterior layers of the true cornea—is to be reckoned to the uveal tract; while the true cornea is a modification of the sclerotic.

may be one or several, varying in size from 1 mm. to 4 mm. in diameter. The vascular injection is immediately around the phlyctenula, and is not diffused over the conjunctiva. At first there may be shooting pains and lacrimation, but these soon pass away. If the phlyctenulæ be not seated close to the cornea the affection is not serious; and the length of time required for its cure depends on the size of the phlyctenulæ, varying from seven to fourteen days, as a rule.

2. **Multiple,** or **Miliary, Phlyctenulæ.**—These are very minute, like grains of fine sand, and are always situated on the limbus of the conjunctiva, which is swelled. The general injection and swelling of the conjunctiva are considerable; and, occurring as it does almost exclusively in young children, the affection may be called Eczematous Conjunctival Catarrh of Children (Horner). The irritation and so-called photophobia and lacrimation are often considerable, and there is a good deal of conjunctival discharge. This form is very apt to appear after measles and scarlatina.

Both forms are liable to extend to the cornea, and then only does the disease become serious. This event may come about in the following different ways:

The Solitary Phlyctenula may be seated partly on the limbus conjunctivæ and partly on the margin of the cornea, and may undergo resolution.

Or it may give rise to a deep ulcer, which either heals, leaving a scar, or perforates, causing prolapse of the iris, etc.

Or it may form the starting-point of a progressive riband-like keratitis (Fascicular Keratitis), the pustule becoming an ulcer, at the margin of which the corneal epithelium is raised and infiltrated in crescentic shape. This now steadily advances for many weeks towards the center of the cornea, followed by a leash of vessels which has its termination in the concavity of the crescent. The process is accompanied by much irritation of the terminal branches of the fifth nerve in the cornea, and the consequent reflex blepharospasm. A permanent mark indicates the track of the ulcer.

The Multiple Miliary Phlyctenulæ on the limbus conjunctivæ may cause some slight superficial infiltration and vascularization of the cornea in their immediate neighborhood, which pass off when the phlyctenulæ disappear.

Or they may be accompanied by deeper marginal infiltrations of the cornea, which become confluent and result in an ulcer that extends along the margin of the cornea for some distance, and is

termed a Ring Ulcer. It is a serious form of ulcer; for if it extends far round it may destroy the cornea in a few days by cutting off its nutrition.

Primary Phlyctenular Keratitis occurs principally in three different forms: (1) Very small gray subepithelial infiltrations, which are apt to turn into small ulcers, and then heal, leaving a slight mark. This mark may ultimately quite disappear, especially in the case of children, and when situated peripherically. (2) Somewhat larger and deeper infiltrations, resulting in ulcers of corresponding size, which heal by aid of vascularization from the margin of the cornea. The opacity left after these ulcers is




FIG. 50.—*E*, Epithelium; *B*, Ant. elastic Lamina; *C*, True Cornea; *N*, Nerve Filament, with Lymph Cells on its course; *D*, Phlyctenula.

rather intense, and clears up but little, especially if the situation be central. (3) Large and deep-seated pustules, often at the center of the cornea, giving rise to large and deep ulcers, which may be accompanied by hypopyon and even by iritis, and which frequently go on to perforation.

Photophobia is usually a prominent symptom in phlyctenular keratitis. The term photophobia, however, is not altogether correct, for it is the fifth nerve (from the cornea) which is mainly the afferent nerve here rather than the optic nerve. This is evident from the fact that in the dark the patient does not get complete relief. The explanation of this reflex blepharospasm has been given by Iwanoff,[1] who showed that the round cells, in making their way from the margin of the cornea to their position under

the epithelium, follow the course of the nerve filaments, which they irritate in their progress. The accompanying Figs. 50 and 51 are from his original paper.

Eczema of the eyelids, face, and external ear, and catarrh of the Schneiderian mucous membrane, frequently accompany phlyctenular conjunctivitis and keratitis.

In these cases, in children of three or four years of age, temporary amaurosis has sometimes been observed after a severe and long-continued blepharospasm has passed away. The patient is found to be unable to see even large objects, or to find his way, although the pupil reflex is active, and a strong light may still be distressing. There are no ophthalmoscopic appearances. This blindness passes away completely, usually in from two to four




Fig. 51.

weeks, although the interval before recovery of sight may be several months. A certain mental dullness, which also ultimately disappears, is noticed in some cases. This temporary loss of sight has been regarded as a reflex phenomenon, and again it has been held to be due to disturbance of the intra-ocular circulation from pressure of the eyelids on the eyeball. But the view (Leber, Uhthoff) which represents it as having a cerebral origin of a functional nature is probably the correct one. It is likely at this tender age, when the psycho-physical processes are not as yet firmly established, that the desire not to see, and the active withdrawal from the act of vision, leads in a short time to a functional paralysis of the visual centers in the brain, and that these take some time to recover or to re-learn their functions when the ground for the suspension of the latter has ceased.

Cause.—This is a disease of childhood, although it is rare in the very first year of life. In adults it is uncommon.

The strumous constitution, as indicated by the swollen nose and upper lip and sometimes by the enlarged lymphatics in the neck, as well as by the eczema just mentioned, is that most liable to this affection. Often, however, it will be found in strong children with apparently perfect general health; but even in them there is probably some allied irregularity of nutrition, of which the great tendency to recurrence of the eye affection is evidence.

Colonies of straw-colored micrococci may be found in the contents of the phlyctenulæ; but what etiological relationship, if any, to the production of the phlyctenulæ they possess is not yet known.

Treatment.—The solitary phlyctenula is best treated with the yellow oxid of mercury ointment in the strength of 10 grs. to ℥j of benzoated lard or vaselin, of which the size of a hemp-seed should be put into the eye once a day. Or a small quantity of pure calomel dusted into the eye once a day will also cure; but this remedy should not be employed if iodid of potassium is being taken internally, for then iodid of mercury is liable to be formed in the conjunctiva.

The miliary phylctenular conjunctivitis is best treated at first with cold or iced applications. Freshly prepared chlorin water (1 part Liq. Chlori., 9 parts water), to be dropped into the eye once a day, is recommended by some, and later on Liq. plumbi dil. or Sol. argent. nitr. (grs. v ad ℥j, and neutralized) applied to the everted conjunctiva; or, if the phlyctenular appearance predominate over the catarrhal, the yellow oxid of mercury ointment or insufflations of calomel may be preferred. I myself rarely employ any remedy other than the two latter, which I find applicable to all these cases.

When the cornea is slightly affected near the margin in cases of miliary phlyctenulæ, calomel, or the yellow oxid of mercury ointment, and warm fomentations, should be used.

Where a large pustule on the margin of the cornea has resulted in a deep ulcer, with tendency to perforate, and accompanied by much pain, I cannot too highly recommend paracentesis of the anterior chamber through the floor of the ulcer, the pupil having first been brought well under the influence of eserin to prevent prolapse of the iris. The good effect of this will be very soon apparent: the pain disappears, the patient sleeps, the ulcer becomes vascularized, and healing sets in. Cauterization of the ulcer in an early stage with the galvano-cautery is also good practice; but in

these cases I prefer the paracentesis. Many surgeons trust very much to eserin, warm fomentations, and a pressure bandage.

For the fascicular keratitis the yellow oxid of mercury ointment is in its place. When the crescentic infiltration is very intense it is well to touch it with the galvano-cautery. Division of the leash of vessels at the margin of the cornea has a beneficial effect.

For the ring ulcer a pressure bandage, under which an antiseptic dressing (boric or salicylic acid, or perchlorid of mercury) has been placed, is, perhaps, the best method of treatment. Warm fomentations promote vascular reaction, and may be used with benefit at each change of bandage.

For primary phlyctenulæ of the cornea, in the form of the minute gray superficial infiltration or ulcer, nothing beyond atropin, with warm fomentations, and a protective bandage to keep the eyelids quiet, should be used. When reparation of the ulcer has commenced, calomel or weak yellow oxid of mercury ointment may be employed.

For the large purulent phlyctenula, resulting in a large and deep ulcer, often situated at the center of the cornea, with hypopyon and iritis, warm fomentations (camomile, or poppy-head, at 90° Fahr., for twenty minutes three times a day), atropin, boric acid as ointment or powder, and a protection bandage form the treatment in the early stages. Here, also, I often puncture the ulcer with the very best results in respect of hastening the cure, and the galvano-cautery may be used with advantage. In the stage of reparation the yellow oxid of mercury ointment or insufflations of calomel are very useful.

In all forms of phlyctenular ophthalmia those favorite remedies, blisters, setons, and leeching, should be avoided. The first two worry the patient, give rise to eczema of the skin, and are not to be compared in their power of cure with the measures above recommended; while leeching gives, at best, but temporary relief, and deprives the patient of blood which he much requires.

For relief of the blepharospasm, in addition to the use of atropin, plunging the child's face into a basin of cold water is a most efficacious means. The face is kept under the water until the child struggles for breath, and this immersion is repeated two or three times in rapid succession, and used every day if necessary. It should always be used where the blepharospasm is severe, as the latter is not only distressing to the patient, but also an obstacle to the cure.

The general treatment, notwithstanding the so-called photophobia, should consist in open-air exercise before everything else, unless, indeed, there be an ulcer which threatens to perforate. It is not well to keep the patient's face or eyes covered with bandages and shades, nor to confine him to a dark room. A pair of dark-blue glasses are the best protection from strong glare of light; and shady places can be selected when the patient is out of doors. Cold or sea baths, followed by brisk dry rubbing. Easily assimilated food at regular meal hours, but no feeding between meals. Regulation of the bowels. Internally: cod-liver oil, maltine, iron, arsenic, syr. phosph. of lime, and such-like remedies are indicated.

The great tendency to recurrence is one of the most troublesome peculiarities of all kinds of phlyctenular ophthalmia: and in order to prevent this, so far as possible, it is important to continue local treatment until the eye is perfectly white on the child's awaking in the morning, and even for fourteen days longer. This prolongation of the treatment will also assist in clearing up opacities, as best they may be. For this after-course of treatment calomel insufflations should be used.

Nothing can be done for the opaque scars left on the cornea by ulcers when all inflammatory symptoms have subsided. If the ulcer have been very superficial the resulting scar in young children may disappear in the course of time. Deep ulcers cause more opaque and permanent scars, and ulcers which have perforated produce the greatest opacity. Some of the very disfiguring scars may be tattooed (see chap. vi.).

The degree of the defect of vision to which an opacity of the cornea may give rise depends, in the first instance, on the position of the opacity. If it be peripheral, the vision may be perfect; but if it be in the center of the cornea, sight may be seriously damaged. Even a slight nebula, barely visible to the observer, will cause serious disturbance of vision if situated in the center of the cornea; while in the same situation the very opaque scar of a deep ulcer will produce a proportionately greater defect. If a central, but not deep, ulcer should not become completely filled up in healing, and a facet remain, vision will also suffer much in consequence of irregular refraction, although there may be but little opacity.

REFERENCE.

[1] "Klinische Monatsblätter für Augenheilkunde," 1869, p. 465.

CHAPTER VI.

DISEASES OF THE CORNEA.

The importance of a knowledge of the diseases and injuries of the cornea depends on their great frequency, coupled with the fact that nearly every one of them is liable to leave behind it some opacity, with resulting defect of sight and disfigurement of the eye; while several of them are very apt to lead to complete loss of sight.

Inflammations of the Cornea.

From a clinical standpoint these inflammations will be most conveniently considered under the headings—(*a*) Ulcerative Inflammations and (*b*) Non-ulcerative Inflammations.

(*a*) Ulcerative Inflammations of the Cornea.—Before an ulcer can form in the cornea there must be a cellular infiltration of its tissue near its anterior surface; and this cellular infiltration is brought about, in most instances, if not in all, by the entrance into the cornea of certain micro-organisms, the gonococcus, or the staphylococcus pyogenes, or other as yet undescribed forms. One recognizes the existence of an infiltration by seeing an opaque spot in the cornea, with a dullness of the layers over it, and often of the corresponding part of the epithelium. Before long the epithelium covering the infiltration comes away, and soon the intervening layers of the true cornea break down, and then we have an ulcer established.

But although all ulcers of the cornea originate in an infiltration, yet when once established they assume great varieties of type, in consequence, probably, of a variety in the nature of the originating micrococcus. Some ulcers are purulent, others non-purulent; some tend to spread over the surface of the cornea, others tend to go deep into it; some attack by preference the central region of the cornea, while others are confined to its margin; some readily give way to treatment, and others are very obstinate or almost incurable. Again, some ulcerative corneal processes are attended by much circumcorneal injection, severe pain in and

about the eye, great reflex blepharospasm, and lacrimation; whilst others, which may really be more severe processes in so far as the integrity of the eye is concerned, can run their course with scarcely any injection of the eyeball, and with little or no distress to the patient.

Etiologically, corneal ulcers are primary or secondary. The primary ulcers are those in which the diseased process originates in the cornea, most commonly as the result of traumata, but also in phlyctenular keratitis, or as the result of corneal abscess, or where the nutrition of the cornea is interfered with, etc. Secondary ulcers are those which are the result of disease elsewhere, usually in the conjunctiva, as in acute blennorrhea and in conjunctival diphtheritis.

Corneal ulcers are more common in advanced than in early life. Indeed, in early life, unless in cases of blennorrhea neonatorum, and of phlyctenular disease, corneal ulcers are almost unknown. The greater liability to these affections in advanced life is due, no doubt, to a less active nutrition at that period in this already lowly organized part. Hence slight traumata, or the presence of a slight conjunctival catarrh, which would have no ill effect in a young person, may form the starting-point of a corneal ulcer in an old person or even in one of middle age. For the same reasons corneal ulcers are much more common in the lower orders than amongst the well-to-do; for the general nutrition of the poor is often defective, while they are more exposed to traumata than are the better classes.

The Diagnosis of the presence of a large corneal ulcer is very simple. Inspection of the cornea in ordinary daylight at once reveals the loss of substance, more or less extensive, deep, and infiltrated. If the ulcer be very small and shallow the difficulty is greater, especially if there be much blepharospasm. In such cases the surgeon must endeavor to inspect the cornea from different points of view, either by directing the patient to move his eye, or by moving his own head, until he succeed in obtaining such an incidence of the light as will display the minute loss of substance, with its margin, and more or less gray infiltrated floor; or he may employ the focal illumination with artificial light. An instillation of cocain may be necessary to facilitate the examination by diminishing the blepharospasm.

It is obviously important to decide at the outset, for the purposes of prognosis and of treatment, whether a gray spot in the

cornea be an infiltration (a collection of cells which may shortly become an ulcer), an ulcer, or a scar (a healed ulcer, or other loss of substance). The surface covering an infiltration, although flush with the general surface of the cornea, has usually a steamy appearance, due to some disorganization of the corneal epithelium, and has no polish. With an ulcer the appearances already described will be found. The surface of a scar is usually, although not always, flush with the general surface of the cornea, and it has a polished surface—*i. e.*, covered with normal epithelium, not rough, irregular, nor even steamy. In cases of corneal infiltration or ulceration there will be usually more or less pericorneal injection, pain, and photophobia, while with a mere corneal scar there will be no irritability of the eye.

A very beautiful method for ascertaining the presence and true extent of a corneal ulcer or traumatic loss of substance is the instillation of a 2 per cent. solution of fluorescin. Almost immediately after the instillation the tissue forming the floor of the loss of substance assume a greenish tint, which clearly differentiates it from the surrounding normal cornea.

The presence of *Hypopyon* (*ὑπό, under; πῦον, pus*) is the rule with some types of corneal ulcer. Hypopyon is a deposit of pus in the anterior chamber, and as the patient sits or stands it lies in the lowest part of the chamber, to which place it has gravitated. If the patient lies in bed (say, on the side of the affected eye) the hypopyon will of course change its position, and gravitate towards the outer side of the chamber. Sometimes the hypopyon is so small as to be detected with difficulty; and again it may fill the whole anterior chamber, completely obscuring the iris and rendering a diagnosis of the condition of the cornea difficult. It will be asked, From whence does the pus come which forms hypopyon in cases of corneal ulcers? It might be supposed that it is derived directly from the purulent floor of the ulcer, by passage of the pus-cells through the posterior layers of the cornea. But this is not so. No pus-cells do, or indeed can, pass through the membrane of Descemet. Moreover, copious hypopyon is often present when the corneal ulcer is quite small and non-purulent. The pus-cells which form hypopyon in cases of corneal ulcer come from the iris, in compliance with the law which causes leukocytes to wander out of blood-vessels in the neighborhood of an inflammatory focus, and to make their way towards that focus. When these leukocytes from the iris reach the anterior chamber they can go

no farther, owing to the barrier imposed to their progress by the membrane of Descemet. The pus forming a hypopyon is sterile. These interesting facts concerning the genesis and nature of hypopyon were discovered by Leber,[1a] whose observations have been corroborated by later observers.

The Dangers attendant upon Corneal Ulcers are, first of all, the




FIG. 52—(*Fuchs*).

opacities, the scars, which even the slightest of them are apt to leave behind.

Fig. 52 represents a section made through a deep ulcer in its progressive stage. At the margin of the ulcer the epithelium (*e*)




FIG. 53—(*Fuchs*).

and Bowman's membrane (b) ceases. The floor of the ulcer is seen covered with pus, which also infiltrates the corneal tissue in the neighborhood. As soon as cure commences the floor of the ulcer begins to get clear—*i. e.*, it becomes gradually less covered with pus—until it is finally quite free from it, and *pari passu* the sur-

rounding infiltration is absorbed. Then the epithelium, growing in from the margin (*m m*, Fig. 53) all round, gradually carpets over the floor of the ulcer, and underneath this newly formed epithelium the new tissue, which is to close the loss of substance, is laid down. This new tissue, however, is not corneal tissue, but is ordinary connective tissue, and is therefore opaque. Hence the deeper the ulcer, the more intense will be the resulting opacity. Bowman's membrane never becomes restored over the cicatrix.

The ulcers which are situated at the center of the cornea, in the pupillary area, are more serious for sight than those situated peripherally, as can be readily understood. The opacity left by a very superficial ulcer is slight, and is called a nebula; a somewhat more intense opacity is called a macula; and a very marked white scar is called a leukoma.

But a more serious danger connected with ulcers of the cornea than the opacities they leave behind is that of perforation of the cornea, to which some ulcers are very prone. For an account of the consequences of perforation see pp. 103, 147, and 171.

In the *Treatment* of primary corneal ulcers the student will soon perceive that a bandage, atropin, and warm fomentations play prominent parts; and these measures alone are sufficient to produce cure in the less severe cases.

The bandage should be put on with firm pressure,—but should not be made uncomfortably tight,—the eye having been previously padded out, especially at the inner canthus, so that equal pressure may be exercised on the globe all over. The support thus given to the cornea and front of the eye promotes the healing process in the ulcer, and the bandage is also useful by preventing the eyelids from rubbing over the ulcer, and by keeping small foreign bodies from it. In secondary ulcers, due to severe conjunctival processes, such as blennorhea, a bandage is contraindicated, because it retains the secretion, and therefore would do more harm than good.

Atropin in sufficient quantities to keep the pupil dilated should be employed. Iritis very often attends severe corneal ulcers, and here the indication for atropin is obvious. But rest of the affected part is, we know, an important element in preventing or in curing any inflammation; and in the affections we are now treating of, even where there is no iritis, atropin acts by procuring rest of the iris and of the ciliary muscle, the constant motion of

which would otherwise tend to augment the inflammatory process in the cornea.

Some surgeons use myotics (eserin, or pilocarpin) in preference to atropin in the treatment of corneal ulcers. They hold that their power of reducing the intra-ocular tension encourages healing of the ulcers; while they also think the more extended surface of iris presented facilitates absorption of the hypopyon. But it is doubtful whether myotics do reduce the normal tension, although they often have that effect upon abnormal tension; and my objection to them in these cases is that they increase, I believe, the tendency to iritis. Absorption of the hypopyon will only come about when the cornea begins to recover, whatever the treatment may be. I am not singular in this view of the use of eserin in corneal ulcers; yet a clear indication for myotics is given by the presence of an ulcer near the corneal margin which has a tendency to perforate, for here the myosis would assist in preventing prolapse of the iris, should perforation take place.

Warm fomentations promote the healing process by stimulating tissue-changes in the cornea. One usually orders them to be made with poppy-head water or camomile tea, although no doubt warm water would be equally efficacious. Hot solutions of 4 per cent. boric acid, or 1 in 5000 corrosive sublimate, may be used with advantage. The bandage having been removed, a compress of lint dipped in the stupe at about 120° Fahr. is laid upon the eye, and frequently replaced by fresh compresses out of the stupe, so that the one on the eye may be always hot. This is continued for half an hour at a time, and repeated every two or three hours. Or, the Japanese muff-warmer may be applied.

In an ulcer of a purulent or sloughing nature the insufflation on its floor of very finely divided xeroform powder is useful.

When more active measures than the foregoing are called for, the actual cautery, scraping, and paracentesis have to be relied on.

The actual cautery has of late years come much into use in the treatment of serpiginous and other purulent corneal ulcers. It acts by destroying the micro-organisms, which keep the process going. Either a thermo-cautère, in the form of a very fine point, or the galvano-cautery (Fig. 54), may be employed, and personally I prefer to work with the former. To the galvano-cautery a medium-sized bichromate of potash bottle-battery is attached, and the platinum wire brought to a red-heat. The eye

having been cocainized, the red-hot cautery is brought into contact with the whole surface of the ulcer, so as to thoroughly de-




FIG. 54.—The bolt *A* being pushed forwards, the current is completed, and passes through the platinum wire which forms the cautery. By pressure on the button *B* the current can be momentarily intercepted during use of the instrument.

stroy its superficial layer, and special attention is paid to any part of the margin of the ulcer where it seems inclined to spread to as yet healthy tissue. Fluorescin may be used to show the extent of the ulcerated surface. The cauterization can be repeated as often as the progress of the ulcer makes it desirable. It is well to perforate the cornea with the cautery, and to evacuate the aqueous humor and hypopyon; or this may be done with an ordinary paracentesis needle, after the cauterization is completed. My own experience of the cautery in these cases is extremely satisfactory. It seems to give the best percentage of cures with the least amount of opacity.

Thorough scraping of the floor of the ulcer with a small sharp spoon is a valuable method.

Paracentesis of the anterior chamber through the floor of the ulcer is another most valuable therapeutic measure for some corneal ulcers, and deserves a more routine application in these cases than is at present accorded to it; the more so as the little operation is simple and dangerless. But here are, I think, two impera-



FIG. 55.

tive indications for its use, namely: (1) If there be great pain. Very shortly after the operation, which for the moment increases the neuralgia, the patient experiences the greatest relief, and passes the first good night after many wakeful ones. (2) If perforation seem to be imminent. This may often be recognized by a bulging forwards of the thin floor of the ulcer; but sometimes it is not easily foreseen, and if there be any doubt on the point, paracentesis should be performed. It is important to forestall spontaneous perforation of the ulcer by this proceeding, because the opening made by the latter being linear it heals easily, and leaves but a slight scar without anterior synechia; while the natural opening would be a complete loss of substance, and, therefore, the more readily involve adhesion of the iris in the resulting comparatively extensive cicatrix. Other indications for the operation are increased tension and the presence of a large hypopyon.

Paracentesis of the anterior chamber is best performed by means of a paracentesis needle (Fig. 55), which is a somewhat

shovel-shaped instrument, with a shoulder or stop. If this be not at hand, a small keratome, or broad needle, or a Graefe's cataract knife with answer the purpose. The eye having been cocainized, a spring lid-speculum is inserted, the eye fixed with a fixation forceps, and the point of the paracentesis needle applied to the floor of the ulcer, in such a way that the plane of the little blade may be at an angle of about 45° with that of the floor of the ulcer. The point is pushed gently through the floor, and the plane of the blade is then immediately changed, so that, as the instrument is being advanced up to the shoulder, it may be almost in contact with the posterior surface of the cornea. The instrument should be withdrawn very slowly, in order that the aqueous humor may flow off gradually, and not with a rush. If these precautions be taken, there need be no danger of injuring the crystalline lens, or of having prolapse of the iris in the incision. If the latter should occur, it can usually be reposed with the spatula. It may happen that when the needle has been quite withdrawn a considerable portion of the aqueous humor may still remain in the anterior chamber, unable to escape owing to the valve-like closure of the wound. It should be evacuated by making the wound gape by gentle pressure with a spatula on its posterior lip. If it be desirable to tap the anterior chamber on the next day, it can be done by simply opening up the wound with a spatula, or with the probe-like instrument at the other end of the handle (Fig. 55), without the aid of any cutting instrument.

If the case does not come under the care of the surgeon until perforation of the ulcer, with prolapse of the iris, has taken place, the very important question as to the best method of dealing with the condition is presented. The same question arises in other forms of perforating ulcer. If the loss of substance occupies one-third or more of the cornea, with correspondingly large prolapse of iris, little can be done beyond the use of eserin—and here I would use eserin—to reduce the intraocular pressure, along with the application of a firm bandage; for in such cases the formation of a corneal staphyloma is almost inevitable. But if the ulcer and prolapse be small, an attempt may be made to free the iris, so that no anterior synechia may form, and in order that the cicatrix may be flat, and not raised over the surface of the cornea, and, consequently, exposed to injury. The importance of such an attempt lies in the fact that a corneal cicatrix with iris entangled in it—not merely adherent to its posterior

surface—affords a constant source of danger, especially if situated near the margin of the cornea; for in such eyes sudden and uncontrollable purulent inflammation of the iris and chorioid may come on after an apparently slight trauma, and end in total destruction of the eye. This event is due to septic infection reaching the interior of the eye through a superficial loss of substance, the direct result of the trauma. The surgeon's attention should, therefore, be directed to obtain at least as flat a cicatrix as possible, or, still better, a non-adherent cicatrix. The practice which I, as well as many other surgeons, have commonly followed, is to draw the prolapsed portion of iris lightly forwards with a forceps, and to snip it off level with the surface of the cornea; and then with a spatula to endeavor to free the iris from any adhesions it may have formed with the margin of the ulcer. Atropin or eserin, according to the position of the ulcer, is then instilled, and a bandage carefully applied. This proceeding is only of use when a fresh prolapse can be dealt with, before cicatrization sets in; and the result is often satisfactory in so far as the securing of a flat cicatrix is concerned, but an anterior synechia can rarely be avoided.

Da Gama Pinto has successfully employed the following method for obtaining a non-adherent cicatrix:—Having abscised the prolapsed portion of iris as above, and freed all adhesions to the margin of the ulcer with a spatula, he covers the opening in the cornea with a flap cut from the bulbar conjunctiva,—and this flap should be twice as large as the opening, in order to admit of its shrinkage,—and then pushes the flap into the opening with a blunt probe. A firm binocular bandage is applied—but no iodoform. The eye is not dressed until the third day, when the anterior chamber is often found restored, the iris all in its proper plane, and the conjunctival flap healed into the ulcer. Ultimately all trace of the flap disappears, and an ordinary non-adherent corneal scar is presented. I have employed this method, and in each case with a good result. Kuhnt also uses it, and he recommends, too,[1] a covering of the ulcer by means of a conjunctival flap of which a pedicle is retained, and which is drawn over the ulcer.

From time to time different types of corneal ulcers have been recognized and described. The following are the chief of them:

Simple Ulcer.—This may result from a slight trauma, or from the bursting of a phlyctenula. It presents the appearance

of a minute and shallow depression with a gray floor on the surface of the cornea. There is circumcorneal vascularity, especially at that part of the corneal margin nearest to which the ulcer is situated; the pupil is apt to be contracted, although iritis is not present, and there is often a good deal of pain, lacrimation, and photophobia.

Treatment and Prognosis.—The eye is to be bandaged, warm fomentations applied several times a day, and a drop of solution of atropin instilled night and morning. When of phlyctenular origin, stimulation with the yellow oxid ointment is indicated. Cure, with slight opacity remaining, comes about in a week or ten days. But occasionally this form of ulcer may pass over to the deep ulcer.

Deep Ulcer.—This is a purulent ulcer, and commences in a purulent infiltration of the cornea. It forms a tolerably deep pit in the cornea towards its center, the floor of the ulcer being covered with purulent deposit and detritus, and the corneal tissue immediately surrounding it being somewhat infiltrated with pus. The ulcer is generally round, but it may assume any shape. Hypopyon is often present, and a marked tendency to iritis exists. The pain is usually very severe, violent frontal neuralgia being a common symptom.

This ulcer has no great tendency to spread over the surface of the cornea, but has a very decided tendency to perforate through it. As it does not generally attain wide dimensions, the perforation it may produce is small, and gives rise to a small adherent leukoma rather than to a staphyloma. This ulcer seldom causes complete loss of the eye.

Causes.—This form of ulcer is a frequent one in purulent conjunctivitis, and it may be caused by the lodgment of foreign bodies, and other injuries of the cornea.

Treatment.—If the ulcer be due to a conjunctival process, the latter should be actively treated.

If the cause be other than conjunctival, a pressure bandage to give support to the ulcer is important, and periodical warm fomentations are most beneficial. Where the cause is conjunctival (purulent conjunctivitis), neither a bandage nor warm fomentations can be used. Atropin should be instilled several times daily. Antiseptic applications, especially xeroform, are useful.

Paracentesis of the anterior chamber through the floor of the ulcer is a proceeding always followed by improvement in the

condition of the eye, and is very important as a preventive of natural perforation. The actual cautery, too, is in its place here.

Serpiginous Ulcer (Ulcus Serpens, Saemisch's Ulcer). This also is a purulent ulcer, the characteristic of which is its tendency to creep over the surface of the cornea, especially in some one direction, rather than to strike deep into its tissue. Its position is chiefly central, and it presents a grayish floor, which is more intensely opaque at some places. One part of the margin takes the form of a curve, or of several closely placed curves, and at this place the margin becomes yellowish-white in color and somewhat raised, and the floor of the ulcer seems deeper in its neighborhood. Immediately around the ulcer the cornea is slightly opaque, but farther out it is normal.

The degree of pain and irritation varies much, being almost absent in some cases, while in others it is extremely intense. Iritis is apt to come on at an early period, and may pass into irido-cyclitis. Hypopyon is almost always present, and on the posterior surface of the cornea, from the region corresponding to the ulcer on the anterior surface, a line of pus is sometimes seen extending down to the hypopyon, and this was formerly taken as a proof that the hypopyon was formed by direct transmission of the pus-corpuscles through the cornea from the ulcer. The ulcer creeps over the surface of the cornea in the direction of the curved and intensely infiltrated margin. At a still later stage the whole cornea is apt to become infiltrated, and the entire margin of the ulcer to extend, and the anterior chamber becomes quite full of pus. Perforation now takes place, or may do so somewhat earlier. If the perforation be small, an adherent leukoma results; but if large, a staphyloma is gradually produced, or panophthalmitis may immediately follow on the perforation.

Causes.—Ulcus Serpens always has its origin in a trauma, which has produced, it may be, only a slight abrasion of the epithelium. In perhaps 50 per cent. of the cases chronic dacryocystitis is present, and in about 25 per cent. more there is ozena, and a considerable proportion of them occur in the warm summer months; and the agricultural population, especially during harvest-time, when injuries to the cornea are frequent, are very liable to it. The specific excitant of this ulcer is the pneumococcus, which can be found in its spreading margins, and which may also be found in the secretion from the diseased lacrimal sac.

Prognosis.—From the above description it will be seen that the process is a very severe one in many instances, and the prognosis unfavorable; yet some cases do recover useful, although damaged, sight under careful treatment, if it has been resorted to in time.

Treatment.—If the case be not severe, atropin, with protection of the eye, may cure in a few days. Here, too, some surgeons prescribe eserin, but I am opposed to its use. Warm fomentations should not be used, as they rather promote the activity of the diseased process; and the eye should not be bandaged, lest infective discharge be retained in the conjunctival sac. Antiseptic measures should always be employed, xeroform being the application most likely to prove of use. It may be employed either in the form of a strong ointment (gr. xxx ad ℥j) put into the eye, or it may be insufflated on the floor of the ulcer with a powder-blower. The floor of the ulcer may be washed with tincture of iodin, a solution of sublimate 1 in 5000, or other antiseptic solutions which do not act as caustics, the action of which on the healthy surrounding tissues might be difficult to control. Scraping the floor of the ulcer with a sharp spoon is a useful procedure. But it is in all respects wiser to deal with these cases, even the apparently mild ones, actively in the very commencement by means of one or other, preferably the second, of the two following methods.

Saemisch's Method consists in division of the ulcer with a Graefe's cataract knofe. Cocain having been applied, the point of the instrument is entered about 2 mm. from the margin of the ulcer in the healthy corneal tissue, and, having been passed through the anterior chamber behind the ulcer, the counter-puncture is made in the healthy cornea some 2 mm. from the opposite margin of the ulcer. The edge of the knife being then turned forwards, the section is slowly completed. The incision should divide the intensely infiltrated part of the margin in halves. The aqueous humor and hypopyon are evacuated, atropin is instilled, a bandage is applied, and the patient soon gets relief from pain. Every day, until healing of the ulcer is well established, the wound must be opened up from end to end with the point of a fine probe or spatula, the contents of the anterior chamber being thoroughly evacuated on each occasion, and atropin instilled. The result is that, in a vast majority of cases, the progress of the ulcer is arrested, and healing soon sets in. The

little operation should not be delayed long, but it may be employed with advantage even in late stages of the process.

But the Actual Cautery is the most valuable method of treatment for this ulcer. The infiltrated and undermined margin of the ulcer is the part which should be most thoroughly cauterized; but its floor, if much infiltrated, is also to be dealt with. The application of fluorescin just before the use of the cautery is of much value, as it enables the operator to clearly discern the whole of the diseased part requiring cauterization.

At the thinnest part of the floor of an extensive serpiginous ulcer it is desirable to make a perforation through the cornea with the point of the cautery; or, instead of this, when the cauterization is finished, the cornea may be paracentesed with a broad needle in a sound region beyond the ulcer. The object is to reduce the intra-ocular tension, and thus promote the nutrition of the cornea.

Subconjunctival injections of a four per cent. saline solution are found to assist the cure, although the explanation of this is not at present apparent.

Where there is dacryocystitis it is of the utmost importance at least to modify its virulence by treatment (*vide* chap. viii.); or, much better, to effect the radical cure by extirpation of the lacrimal sac. These ulcers require prompt relief before the time is passed when relief can be of any practical use; and the cautery undoubtedly affords the best prospect of aid of that kind. Even the cautery is too often ineffectual to arrest the progress of the ulceration.

Rodent Ulcer.—This is a rare and extremely dangerous form of ulcer, which was described by Mooren,[2] and it must not be confounded with the serpiginous ulcer. It is not a purulent ulcer. It appears as a small—sometimes even pinhead-sized—gray infiltration near the corneal margin, not differing in appearance from many a harmless catarrhal infiltration. This rapidly ulcerates. Other similar infiltrations appear in the neighborhood and at other parts of the margin, and ulcerate, and the ulcers coalesce into one, of which the margin nearest the center of the cornea is undermined. There is very great pain in some cases, and in others hardly any. The eyeball is injected. The ulcer does not go deeper than about one-fourth of the thickness of the cornea, and perforation seldom occurs. Occasionally a very small hypopyon is present. Before long the ulcer begins to

be vascularized and to heal, and finally leaves an intense cicatrix behind. Gradually the ulceration creeps round the cornea, and at the same time advances towards its center, healing going on the while in the oldest portions of the ulcer. This process goes on until, finally, the whole surface of the cornea has been eaten away, and cicatricial tissue substituted for it, its center being the last place affected, and then vision will have become reduced to finger-counting or to perception of light. Clearing up of the corneal opacity to a certain extent may subsequently take place, but cannot be reckoned upon. Yet in a few cases fairly good vision has been regained. The disease usually comes on in both eyes, although there may be an interval between the onset in each, of weeks, or months, or more. It attacks decrepit people of over middle life, but it also occurs in young persons and in those of apparently robust health. Its etiology is obscure. No specific micro-organism has as yet been discovered as the immediate cause. The progress of the disease is very slow, many weeks or even some months often elapsing before the surface of the whole cornea has been destroyed.

The onset at the edge of the cornea in the form of small gray infiltrations, the gray and shallow floor of the ulcer, its pale gray or almost white margin, the undermining of this margin (which may readily be ascertained by passing the point of a probe under it), and the steady advance of the ulceration towards the center and around the edge of the cornea, are the characteristics of this disease.

Treatment.—The general nutrition of the individual is to be improved, but reliance is mainly to be placed on local treatment, which should especially be directed to the undermined margin. The actual cautery may save the eye in some few cases. Scraping, tincture of iodin applied with a camel's-hair pencil, sublimate lotion, carbolic acid, with a bandage and the usual warm fomentations, may help in the treatment. The covering of the diseased part—after it has been well cauterized—or of the entire cornea, with a conjunctival flap, is a measure which is well worth the trial. Quite recently I observed in one case—the only case in which I have as yet tried it—that the application of absolute alcohol to the ulcer arrested its progress.

Prognosis.—From what has been said, it is evident that the prognosis in these cases is very unfavorable.

Marginal Ring Ulcer is a rare form, which commences as

a clean-cut, or but slightly infiltrated, yet rather deep, ulcer at the corneal margin. Its tendency is to extend along the margin of the cornea; and in some instances healing takes place in the older parts of the ulcer while it is still progressive at the newer parts. It may extend all round the cornea, and finally give rise to complete sloughing of the latter by cutting off its nutrition. This ulcer may result in children from a marginal phlyctenular infiltration (p. 134), but is more common in adults, or in aged people whose nutrition has fallen very low.

Treatment.—The actual cautery. Paracentesis through the ulcer, eserin having been first instilled. Insufflation of xeroform. Warm fomentations. A bandage. Quinin, iron, and strychnin internally, with nutritious diet.

Absorption Ulcer (Faceted Ulcer, Superficial Transparent Ulcer) is the term applied to a certain definite superficial ulceration which is accompanied by but little opacity and by no vascularization, and which is usually seated at or near the center of the cornea, where it presents the appearance of a shallow pit about 2 mm. broad, with rounded margin, its floor being covered with epithelium. If the eye be exposed to cold, wind, or other irritation, some circumcorneal injection makes its appearance, and there is lacrimation; but these symptoms soon pass off again. The healing process may take months to be completed, and slight opacity remains. Often the defect is never quite filled up, but a small facet is left, which is liable to interfere with vision.

The absorption ulcer does not tend to perforate, nor to spread over the surface of the cornea.

It occurs chiefly in childhood, and probably indicates malnutrition of the general system; some observers, indeed, think there is a close relationship between it and phlyctenular ophthalmia. It is also seen in granular ophthalmia, with and without pannus.

Treatment consists in atropin and protection, with a bandage in the early stages; and the yellow oxid ointment in the later stages. General treatment with suitable tonics is indicated.

Neuro-Paralytic Keratitis.— In paralysis of the Ophthalmic Division of the Fifth Nerve purulent infiltration with hypopyon and ulceration of the cornea is often observed. It was formerly believed that the fifth nerve had an influence over the nutrition of the cornea, and hence that this was a trophic process; but experiment has shown that this is not the case, and that the affection

is merely due to the loss of sensation, and consequent drying and disorganization of the epithelium, which renders it possible for septic infection of the cornea to take place. This disease, therefore, cannot be regarded as of neuropathic origin in the strict sense of the term.

Treatment consists chiefly in protection of the cornea by a bandage on the eye, or by keeping the lids fastened together with a dermic suture.

Infantile Ulceration of the Cornea, with Xerosis of the Conjunctiva, first described by von Graefe,[3] is a very rare affection. It attacks some marasmatic children early in the first year of life, making its appearance at or near the center of the cornea. Iritis always supervenes in severe cases. That portion of the bulbar conjunctiva which is exposed in the palpebral aperture at either side of the cornea undergoes slight epithelial xerosis, as in functional night blindness, due to retinal exhaustion (see chap. xvii.). Sometimes the xerosis of the conjunctiva is absent,[4] as Holmes Spicer has shown. Ulceration of the cornea soon comes on, through necrosis of the layers lying over an interstitial infiltration; and this ulceration spreads until it involves the whole of the cornea, except a very narrow margin. Finally, perforation, with prolapse of the iris, and panophthalmitis may supervene.

Both eyes become affected as a rule, although the disease usually attacks one eye some time before its fellow. The patients almost always die of diarrhea, pneumonia, etc.

Cause.—Streptococci have been found [5] in the corneal ulcer and in the conjunctiva, while a general invasion of the vascular system of the whole body is also present. To the latter circumstance are referred the symptoms which lead to a fatal termination.

Treatment is, unfortunately, of very little avail; but warm fomentations and the use of non-irritating antiseptic lotions, etc., are indicated, along with an antiseptic bandage. Such means as may possibly promote improvement of the general system are obviously called for.

Herpes Corneæ Febrilis.—Not only in herpes zoster ophthalmicus, but also in herpes febrilis (or catarrhalis) is a vesicular eruption liable to occur on the cornea. According to the late Professor Horner, herpes corneæ febrilis is a rather common affection, and, he believed, is often not recognized by ophthal-

mologists because it usually first comes under their notice when the secondary ulcers have formed. The following is Horner's description of the disease:

On the surface of the cornea of one eye is formed a group of clear vesicles, each from 0.5 to 1.0 mm. in diameter, their appearance being accompanied by much lacrimation, but without any swelling of the eyelid. They usually form in a line, which runs obliquely across the cornea, or sometimes in a vertical direction. Now and then they are arranged in trefoil shape or in a circle. The covering of the vesicles is short-lived, and, as already remarked, the resulting ulcer is that which the surgeon usually first sees. Even it, however, is thoroughly characteristic. On the surface of the clear cornea is an irregular loss of epithelium, along the margins of which may still sometimes be seen the shreds of the late covering of the vesicle. The margin of the region which is bared of its epithelium is dentated, and can only be mistaken for a traumatic loss of epithelium. The latter, however, would never present the peculiar "string-of-beads" appearance. The floor of the loss of substance is formed by the superficial layers of the cornea, and the anesthesia of the cornea is confined to this place, and does not, as in herpes zoster, extend to the rest of the cornea. The tension of the eye is generally reduced. Under favorable circumstances this loss of epithelium may be rapidly repaired; although even then more slowly than one of equal dimensions, but of traumatic origin. Usually the healing process is slow; and sometimes more or less intense opacities form in the area and at the margin of the ulcer, with hypopyon, iritis, etc., and the loss of substance becomes deep, with a dentated margin. This more unfavorable course is the result of secondary infection of the ulcer.

The subjective sensations are those of a foreign body in the eye, with lacrimation and photophobia, and are relieved immediately after the bursting of the vesicles.

The vesicular eruption is often regarded as irritation from a foreign body merely; or, occurring in the course of a serious disease (pneumonia, typhoid fever, intermittent fever, etc.), it passes wholly unnoticed, and its relationship to the latter remains unrecognized.

The only affection for which herpes corneæ is likely to be mistaken is phlyctenular keratitis; but the clear elevated vesicles will readily be distinguished from the flatter grayish mass of cells

which form the phlyctene. In herpes there is never—although often in phlyctenular keratitis—a vascularization of the cornea. The shape of the loss of epithelium after bursting of a herpes vesicle is characteristic. Phlyctenular keratitis is a disease of childhood, while herpes corneæ is rare under puberty.

The derangements of the system in which herpes corneæ febrilis occurs are naturally those in which herpes febrilis labii, nasi, etc., are found. These are more especially the inflammatory affections of the respiratory tract, from an acute catarrh of the Schneiderian mucous membrane to a severe pneumonia. On two occasions, with an interval of three years, Professor Horner saw herpes corneæ occur in the course of an attack of pneumonia in a boy. In just such cases herpes on the lips, ala nasi, external ear and eyelid of the same side are found; and in a case of double pneumonia in an adult occurred the only binocular herpes corneæ which Professor Horner had seen. He explicitly states that he had seen herpes corneæ in connection with whooping-cough, and often with intermittent and typhoid fevers.

But primary herpes corneæ—*i. e.*, unconnected with any other disease—is occasionally met with; and some patients are liable to recurrent attacks of it. It is accompanied by severe neuralgia in the frontal and temporal regions, and pain on pressure of the supraorbital notch may be present. There is much lacrimation. The upper lid is red and swollen. The bulbar conjunctiva, especially around the cornea, is much infected, and there may be a few vesicles on it. Over the surface of the cornea, but sometimes confined to some one district of it, there are a number of minute vesicles, some shreds of epidermis—the remains of ruptured vesicles—and round grayish-white superficial infiltrations not larger than a pin's head. The mucous membrane of the nostrils is also apt to be attacked, causing swelling of it, with much secretion, and the formation of scabs.

Treatment at an early stage, before the vesicles have burst or the loss of substance has become infiltrated, consists in protection of the eye, and, when infiltration has set in, in disinfection, with protection. If the vesicles give great pain they may be ruptured by dusting a little calomel into the eye, or by brushing it with a camel's-hair pencil wet with solution of boric acid, after which a well-fitting antiseptic bandage is applied. Cocain is valuable in these cases for relief of the pain. Atropin and warm fomenta-

tions should also be employed, and a weak yellow oxid ointment is of use in some cases. Where the nostrils are affected weak sublimate or other antiseptic washes should be applied to the Schneiderian mucous membrane.

Filamentary Keratitis (Fädchen-Keratitis).—This is very rare. It may occur with or without superficial injury to the cornea. Its name is due to the fine threads, like twisted spun-glass, several of which hang from the surface of the cornea, and give the condition its characteristic appearance. These threads never reach a length of more than 3 or 4 mm.

Different views are held as to the mode of origin of the threads. Fischer and Uhthoff [6] have observed that small vesicles, with clear or turbid contents, appear in groups upon part of the cornea, then burst, and from the center of each resulting depression a thread hangs out. The onset of the vesicles is accompanied by much pain and photophobia, and probably has its cause in some affection of the fifth nerve. The duration of an attack is usually short, but there may be sevral relapses at brief intervals, and finally the process ceases without permanent damage to the cornea. These same authors hold that the threads are composed of the peculiar fibrinous contents of the vesicles. But it has now been proved beyond doubt by the investigations of Hess [7] and Nuel [8] that the threads are composed of twisted proliferating epithelial cells, each thread ending in a bulbous enlargement caused by degeneration of the epithelium. A peculiar diseased condition of the corneal epithelium precedes the formation of the vesicles and threads.

Treatment.—The instillation of a 3 per cent. solution of chlorid of ammonium into the eye every two hours, by which the exfoliation of the epithelial growth is promoted and hastened. Protection of the eye with a bandage is important.

Bullous Keratitis.—Bullæ very rarely form on the cornea. They are seldom the primary condition, but usually depend on an interstitial diseased process in the cornea. This process may itself be a primary disease; but more commonly it, too, is secondary to deep changes in the eye, such as absolute glaucoma, irido-cyclitis, etc. I have a few times seen bullæ form on the cornea of otherwise sound eyes in persons whose health was in a debilitated state. The formation of a bulla is attended by much pain and photophobia, which disappear as soon as the bulla ruptures. One, or more than one, bulla may form at a time. After a day

or two they rupture, and their walls then hang in shreds from the surface of the cornea, and the seats of the bullæ present shallow depressions. These losses of substance heal without leaving any permanent opacity. After an interval of days or weeks another crop of bullæ appears, and runs the same course.

Treatment.—The bullæ should be opened, and their walls snipped away with a scissors, and a bandage applied. The recurrent attacks may cease after a length of time; but so far as treatment can influence them it can only be done by relieving the process in the cornea which gives rise to them. If it be a primary process warm fomentations, atropin, and a bandage, with remedies directed to correction of any fault in the general state of the health which may exist, are suitable; or if, as is more common, a deep ocular process (glaucoma, etc.) be the cause, the recognized treatment for this latter must be adopted.

Dendriform (*δένδρον, a tree*) **Keratitis.**—This is a rare affection, to which attention was first drawn by Hansen Grut, of Copenhagen. It is a very superficial and chronic ulceration, with but little infiltration of its margins or floor, and presents the appearance of a fine groove on the cornea. It spreads chiefly over the central region of the cornea by throwing out branches on either side. The pain and irritation are sometimes severe, and again but slight or quite wanting. Some slight permanent opacity may remain when cure has been effected.

The Cause has not been definitely ascertained, but the peculiar progress of the affection renders it almost certain that some special fungus is engaged.

It must, however, be stated that the opinion is strongly held by some that these ulcers result from a herpetic eruption on the cornea—in short, that they are the ulcers observed by Horner as the result of herpes corneæ febrilis (p. 156).

Treatment.—Scraping with a sharp spoon, with the subsequent application of 1 in 1000 solution of corrosive sublimate to the cornea, is recommended by some, also the application of pure carbolic acid to the ulcer with a fine camel's-hair pencil, care being taken to confine it to the ulcer, and the actual cautery is sometimes useful.

I can strongly recommend the application of absolute alcohol. A small bit of lint is folded to a point, and the latter is dipped in the alcohol. The ulcerated portion of the cornea is then rubbed with the point with such pressure as to take away the

epithelium, and, so far as possible, the rest of the cornea is avoided. Immediately afterwards the conjunctival sac is freely washed out with sterilized salt solution, to remove all surplus alcohol which would increase the subsequent pain. Usually one application is sufficient to produce cure, but some cases require it to be repeated after four or five days.

I have also found the application of a fine point of sulphate of copper to the ulceration produces good cures. It is less painful than the alcohol, because its action is easily confined to the ulcerated part.

Keratitis Aspergillina.— This disease was described by Leber,[9] and some seven or eight cases of it have been published. The appearance presented is that of an ulcer from 3 to 5 mm. in diameter, occupying a rather central position in the cornea. The surface or floor of the ulcer is of a pale grayish yellow, and is very irregular. A striking and characteristic appearance is the dryness of this surface, the copious discharge of tears flowing over it without seeming to wet it. The rest of the cornea is slightly opaque and dull, and there is a small hypopyon present. The conjunctiva is injected and swollen, and is covered with some mucous secretion. The eyelids are rather swollen. There is photophobia and often severe pain. Masses removed from the surface of the ulcer and examined with the microscope are found to be full of the aspergillus fumigatus. It may usually be ascertained that an injury has preceded the appearance of the ulcer.

Treatment.—The membranous mass which forms the floor of the ulcer should be peeled off, and the underlying surface cauterized and dressed with xeroform, after which a good and rapid cure takes place.

(*b*) Non-Ulcerative Inflammations of the Cornea.— **Abscess.**— This affection is on the borderland between the ulcerative and non-ulcerative inflammations of the cornea; for in one case it will result in an ulcer—usually the ulcus serpens—while again it will run its course without ulceration. The abscesses which are seated in the more superficial layers are those which go on to ulceration; those in the deeper layers are less likely to do so.

Abscess differs from infiltration in that the pus which forms it destroys the true corneal tissue—the fibrillæ and fixed corpuscles—and does not merely lie between them

Signs and Symptoms.—The appearance presented is that of a yellowish circumscribed opacity, more intense at its margin than at its center, seated at or near the middle of the cornea, and surrounded by a light gray zone. It is usually round in shape, but when situated near the edge of the cornea it is apt to be crescentic. The surface of the cornea just over the abscess is at first a little elevated over the general surface, but later on becomes flattened, owing to a falling-in of the normal layers anterior to the abscess; and the epithelium of the flattened part has a dull, breathed-on look. The rest of the cornea may also lose its brilliancy, although in a much less degree. Hypopyon and iritis are constant attendants upon corneal abscess. There is much injection of the conjunctival and ciliary blood-vessels. Severe pain in and about the eye and blepharospasm are common, yet occasionally a corneal abscess will be attended by but little pain or irritation.

Progress.—The abscess spreads through the cornea, usually in some one direction, and this direction is indicated by the yellowish opacity being more intense at the advancing side of the abscess. Before long, if the abscess be superficial, the layers of cornea covering it come away, and the condition is changed into that of the ulcus serpens already described. The deeper abscesses spread through the cornea more or less widely, and ultimately become absorbed, without having caused ulceration. But even these abscesses leave considerable opacity behind. The process which ends in ulceration is the more common of the two.

Etiology.—Abscess is the result of infection of the cornea with pyogenic organisms, which reach it either from without, through some traumatic loss of substance of the corneal epithelium, or from within, by the agency of the blood. The micro-organisms, which are introduced through a superficial loss of substance, may either have been on the foreign body which produced the injury, or they may have been present in the conjunctival sac. Infection through the blood is occasionally seen in some acute exanthematous diseases, such as scarlatina, measles, and smallpox; more especially in the latter in its convalescent stage.

Treatment.—Atropin, warm fomentations, and a bandage. But if these mild measures do not in a day or so arrest the progress of the abscess, resort must be had to the actual cautery.

Diffuse Interstitial, or Parenchymatous, Keratitis.—This af-

fection occurs most commonly between the ages of five and fifteen. It usually commences at some one part of the margin as a light grayish opacity, accompanied by slight injection of the ciliary vessels. The rest of the corneal margin soon becomes similarly affected; and then gradually the opacity extends concentrically into the cornea, or does so by sending in processes which afterwards become confluent. In this way the whole cornea becomes affected by degrees; and its epithelium acquires the breathed-on or ground-glass appearance, which is seen, also, in acute glaucoma. Occasionally the opacity commences at the center, and not at the margin of the cornea, as small gray spots, and extends towards the margin, which it often does not reach before clearing commences.

The opacity lies in the deep layers of the true cornea, and is slightly more intense in spots here and there. It is sometimes only a very light cloud, while again the cornea may be so opaque as to render the iris quite invisible. Along with the opacity, vessels form in the cornea in its posterior layers, but the degree of vascularization varies very much in different cases. In some the presence of vessels can only be ascertained by careful examination with a high convex glass (+ 16.0) behind the ophthalmoscope, or with the corneal microscope; while in others the new vessels are present in great numbers, and can be readily seen with the naked eye. In other cases, again, close leashes of vessels follow the tongues of opacity into the cornea, giving rise to the appearance known as the " salmon patch."

The infinite variety in the degree of opacity and in the amount and arrangement of the vascularization results in a great variety of the appearances in different cases. When the whole cornea has become opaque it begins to clear up at the margin, and the central portion becomes even more opaque than the margin had ever been—a fact which shows that the very cells which entered the cornea at its margin have advanced to its center. The clear margin gradually increases in width until only a rather intense central opacity is left. This central opacity slowly breaks up, and becomes absorbed, but not always completely; and then considerable and permanent impairment of vision may remain.

In severe cases iritis and chorioiditis are nearly always present, although the latter is not observable until the cornea has become clear enough to admit of an ophthalmoscopic examination. The disease, indeed, must be regarded, strictly speaking, as one of the

uveal tract, to which the posterior layers of the cornea, which are mainly diseased, belong.

The two forms above described, one commencing at the margin, the other at the center of the cornea, and more or less vascularized, but for the most part ultimately occupying the entire cornea, are those we are wont to find in children and young adults, and which, as will just now be stated, have congenital syphilis as their usual cause. But in older persons, up to thirty or thirty-five, milder forms of interstitial keratitis are met with. These rarely occupy more than a small region of the cornea, generally towards its center, either as a patch or as a ring of opacity, and with little or no vascularization.

The affection is often accompanied by a good deal of pain and blepharospasm, especially in the severe vascular forms, and there, too, the tension of the eye is apt to be temporarily reduced.

The acute stage of the disease lasts from six to eight weeks, or longer. But the entire process may not be completed for many months, and in one case which I saw the opacity did not begin to clear away for eleven months after the cornea was first attacked, the whole process extending over a period of two years.

Both eyes invariably become affected, although not always at the same time, the second eye being often not attacked until the inflammation in the first has made some progress, or, perhaps, not until it has undergone cure. It is important, in the very commencement of treatment, to acquaint the patient or his parents with the likelihood of this course of events.

In adults usually one eye alone is attacked, iritis is rare, the duration of the process is comparatively short, and complete clearing up is relatively frequent.

Causes.—The affection is more common in girls than in boys, and most frequently appears during second dentition, when the upper incisors are being cut, or at puberty. It depends upon some serious derangement of the general nutrition; and this, in over 50 per cent. of the cases, is inherited syphilis—a fact which was first pointed out by Mr. Jonathan Hutchinson. The children are often thin, anemic, and of stunted growth, with flat nose, cicatrices at the angles of the mouth, often more or less deaf; and the peculiarities of the incisor teeth, so well known from Mr. Hutchinson's description, are present in about one-half of the cases.

Occurring in adults, the affection is rarely due to inherited syphilis, although acquired lues may sometimes be taken as its

cause; while, again, it will often be impossible to assign any origin for it other than the universal one of exposure to cold, etc. Possibly some cases are due to tubercular disease.

Prognosis.—In children—in view of the possibility of an incomplete clearing of the cornea and the irregularity of its surface which the process may cause, as well as of the serious complications liable to supervene, and which may completely annihilate vision—the prognosis must be guarded (although by no means hopeless) in those cases where the opacity is very intense, or where there is much vascularity. Yet in the milder cases a very favorable prognosis may be given. I have never seen the affection recur, but it is said to do so very rarely.

In adults, as stated, the prognosis is much more favorable.

Treatment.—In the early stages no irritants should be locally applied. Atropin is important for the prevention of iritis or of posterior synechiæ; and the use of radiant heat, in the form of hot poultices or fomentations, or the Japanese warmer, promotes the nutrition of the cornea and hastens the cure by absorption of the cellular elements which form the opacity. A bandage should be worn. When the acute stage is ended the yellow oxid ointment may be employed with benefit for stimulating the absorbents to carry off what remains of the opacity. Massage may be used with advantage in both stages to disperse the infiltration. In severe cases I would advise a course of mercurial inunctions, continued for several weeks, care being taken not to allow stomatitis to exceed very moderate bounds. In mild cases a tonic plan of treatment, with iodid of iron and cod-liver oil, is the most suitable.

Counter-irritation, in the form of blisters to the temple or a seton in the scalp, is extensively employed by some surgeons. I have never adopted this treatment, as I doubt its value and am loath to add a worry to the troubles inseparable from so wearisome a disease.

Keratitis Profunda.—This also is an interstitial or parenchymatous keratitis. It presents the appearance of a grayish opacity in the deep layers of the cornea at or near its center, which gradually increases in intensity and extent, but never reaches the margin of the cornea. The epithelium of the cornea over the infiltration is dull. There is but little vascularization and no great tendency to inflammation of the uveal tract. After some weeks the infiltration gradually breaks up and becomes absorbed, leaving in severe

cases a good deal of opacity, and in less severe cases a fairly clear cornea. The affection is seen only in adults, and its etiology is unknown. It seems probable that it is the result of a localized derangement of the posterior epithelium of the cornea;[10] but if that be so, the ultimate cause of that derangement has not yet been suggested.

Treatment.—Atropin, warm fomentations, and bandage, with care of the general health.

Grating-like Keratitis.—This rare disease was described by Haab.[11] Its presence is very apt to be overlooked in the early stages, for by focal illumination the cornea may seem perfectly normal, and the rest of the eye, except perhaps for some slight distention of the anterior ciliary and larger conjunctival vessels, is healthy, and is not irritable. The patient complains of slight burning sensations (especially during use for near work), of some photophobia, and of somewhat diminished acuteness of vision. Transmitted light from the mirror displays in the illuminated pupillary area a number of fine radiating forked lines, recalling somewhat the appearance in some incipient cataracts, and between these lines there are innumerable fine dots. With the corneal microscope these dots and lines are found to be grayish, and to be situated closely under the epithelium. The surface of the latter is at this period in no way altered. A zone 2 or 3 mm. in width inside the corneal margin is quite free from both lines and dots, while the central region is occupied by dots alone, and contains no radiating lines. The latter are present in the intermediate zone only. In the further progress of the disease the lines advance into the center of the cornea, and the gray dots increase in number, the anterior ciliary vessels are more distended, vision sinks lower, and the patient may occasionally complain of some pain, with lacrimation, and swelling of the eyelids; but more commonly there is no irritation. The lines and dots now begin to show as slight elevations on the surface of the cornea, although under the epithelium, and the arrangement of the radiating forked lines begins to resemble the appearance of an iron grating. At a still later period the opacity in the center of the cornea becomes more intense and may even be leukomatous, while a marginal zone of the cornea remains fairly clear. Owing to the difficulty of distinguishing lines and dots, the diagnosis at this advanced stage may again be rendered doubtful, and the appearance is apt to be confused with that left in some severe cases of diffuse parenchymatous keratitis.

It is probable that the changes which produce the appearances described above commence in Bowman's layer, and consist in a hyaline degeneration of it, and that this process subsequently extends to somewhat deeper layers. The disease is apt to be found in more than one member and generation of a family. No constitutional or other cause can be assigned to it. It seems as a rule to commence in youth—in the fifteenth to twentieth year or thereabouts—and to progress slowly in both eyes during many years, leaving the patient almost blind in middle life.

Treatment is practically useless. Yellow oxid of mercury ointment has been employed, and warm fomentations, also galvanism, and hydrate of chloral eye-drops.

Guttate Keratitis.— This is a very similar disease to the foregoing, and, when our experience is greater, may prove to be one and the same with it. It was described by Groenow.[12] It differs from the grating-like keratitis in that no radiating lines are present in the cornea, but small gray patches and fine dots only. The small patches are of all shapes, and occupy the central region of the cornea, while the fine dots are present in every part, but more abundantly towards the center than elsewhere. These patches and dots lie in the most anterior layers of the cornea, and the epithelium over them becomes raised, giving an uneven and dull appearance to the corneal surface. Both eyes are affected. Pain or other irritation is almost absent. Most of the cases observed have been in men, and usually in young men. The cause is unknown. No relation to syphilis or other constitutional condition has been made out. Fuchs ascertained that the seat of the mischief is in those corneal cells which lie between the deeper and superficial layers, and which he found to contain mucin. Like grating-like keratitis, this is an exceedingly chronic disease, lasting many years, and finally causing much loss of sight. Similarly, too, it may so far be said to be incurable.

Keratitis Punctata.— Until recent years this term was only given to a condition which occurs in cyclitis, in iridocyclitis, and in sympathetic ophthalmitis, and which is not a primary disease of the cornea, and therefore does not come within the scope of this chapter. It will be considered under those headings (chap. x.); and it need only here be stated that it consists in the deposit, in the form of fine dots on the back of the cornea, of lymph derived from inflamed portions of the uveal tract, mainly from the inflamed ciliary processes.

Fuchs [13] has described a form of keratitis which he terms *Keratitis punctata superficialis,* and which has a good claim to that name. It begins with the symptoms of an acute conjunctivitis. Either at the same time, or some days or weeks afterwards, minute gray spots may be seen in the superficial layers of the cornea, the epithelium over the spots being somewhat raised up. The spots are often arranged in groups or rows, and may be scattered over nearly the entire cornea, or else confined to its central region. There may be but a few of them, or there may be a hundred or more, and one or both eyes may be affected. The initial irritative symptoms soon disappear; but the spots themselves remain for many weeks, or longer, and finally fade away entirely. It is more common in young people than in later life, and occurs usually in connection with a catarrh of the air passages; but it must not, by reason of this, be confounded with herpes of the cornea. The spots are often very faint, and hence can easily be overlooked, unless searched for with the oblique light. In this country the affection is rather rare, but I have seen several cases of it.

The Treatment should consist in atropin, bandage, and warm fomentations.

Sclerotizing Opacity of the cornea sometimes complicates scleritis, affecting the margin of the cornea in the neighborhood of the scleral affection, but not extending more than 2 to 3 mm. into the cornea, except in very severe cases. It is an intense white opacity situated in the true cornea, and is apt to remain as a permanent opacity, even when the scleritis undergoes cure. In such cases of sclero-keratitis iritis is often present.

Treatment.—Warm fomentations, massage, and the treatment of whatever diathesis (rheumatism, syphilis) may be taken as giving rise to the scleritis.

Ribandlike Keratitis (Transverse Calcareous Film of the Cornea; Calcareous Film of the Cornea).—This is a degenerative alteration which occurs chiefly in the corneæ of eyes destroyed by severe intra-ocular processes, such as iridocyclitis, sympathetic ophthalmitis, glaucoma, etc.

It also occasionally occurs as a primary disease in some persons of advanced life. In these latter instances glaucoma often comes on at a later period, or the corneal disease may be followed by iridocyclitis, or central chorioiditis. It seems probable that in these primary cases the cause of the degeneration is simply a loss of vital energy in the corneal tissue, due, it may be, to vascular

changes. These latter, in their turn, are held by Nettleship to be due in many instances to gout.

The disease occupies that transverse strip of the cornea which is uncovered in the commissure of the eyelids during waking. It usually commences on the inner margin of the cornea, but soon appears at the outer margin, and advances from each direction towards the center, where the two sections join. It presents the appearance of a grayish-brown opacity, with, in most cases, white calcareous deposits in and under the epithelium. In blind eyes which are constantly rolled upwards, the opacity is found not in the central transverse section of the cornea, but in the exposed lower third. The opaque masses consist of carbonate and phosphate of lime. Leber puts forward [14] the view that an abnormally abundant supply of phosphate of lime in the blood, and nutritive fluid of the cornea, is the cause of this condition, the rapid evapo-




Fig. 56—(*Pagenstecher*).




Fig. 57—(*Pagenstecher*).

ration on the exposed part of the cornea being the reason why the deposit takes place there. The deposit is at first in Bowman's membrane, but later on it may appear in the anterior layer of the true cornea, and in the epithelium.

Treatment.—Some improvement may be effected by scraping away the chalky deposit.

Ectasies of the Cornea.

Staphyloma of the Cornea is the result of a perforating ulcer of the cornea. The ulcer, having healed, may present a weak cicatrix, which becomes bulged forwards by even the normal intraocular tension (Figs. 56 and 57). If the iris be not involved in this cicatrix the anterior chamber will be made deeper (Fig. 57).

Staphyloma corneæ, in which the iris is involved, is probably a

more common condition than the above. When a corneal ulcer is large, a correspondingly large portion of iris is liable to become prolapsed into it, and to form a bulging mass. This may burst and collapse, and a flat cicatrix may be formed; or, if it do not rupture, it may form what is termed a partial staphyloma of the cornea and iris, the latter becoming consolidated by the formation of a layer of connective tissue over it.

If the whole, or a very large part, of the cornea be destroyed by an ulcer, the iris is completely exposed. It soon begins to be covered with a layer of lymph, which gradually becomes converted into an opaque cicatricial membrane. Should this not be strong, the normal intra-ocular tension is sufficient after a time to make it bulge; or, increased intra-ocular tension may arise in consequence of further changes within the eye, and then bulging of the pseudo-cornea all the more surely comes on, and the condition is termed total staphyloma of the cornea. Sometimes a total staphyloma has a lobulated appearance, owing to the pseudo-cornea having some of its fibers stronger than others, and hence the name given to the condition (from σταφυλή, *a bunch of grapes*), and which has in time become applicable to almost any bulging of the cornea or sclerotic. Such staphylomata are apt to gradually increase to a very large size.

Treatment.—In cases of partial staphyloma, where a clear portion of the cornea remains, an iridectomy is frequently indicated for the reduction of the tension—so that further bulging may be arrested—as well as for the sake of the artificial pupil, which may improve sight, in cases where the normal pupil is obliterated by corneal opacity. When, sight having been lost, the staphyloma is very bulging, or when total staphyloma is present, enucleation of the eyeball, or one of the following operative measures, must be adopted.

Abscission.—A Beer's cataract knife being passed through the base of the staphyloma, with its edge directed upwards, the upper two-thirds of the staphyloma are separated off, while the remaining third is detached by means of a scissors. If the lens be present it must now be removed. The wide opening becomes filled up with granulations and cicatrizes over.

In de Wecker's [15] method the opening is closed with conjunctival sutures. He begins the operation by separating the conjunctiva all round the margin of the cornea, and by then loosening it from the eyeball nearly as far back as its equator. Four

sutures (*a, b, c, d*) of different colors are then passed through the conjunctiva about 2 to 3 mm. from the margin of the wound, as represented in Fig. 58. In order to keep the field of operation clear the ends of two of these sutures are laid over on the nose while the others are laid over on the temple. The staphyloma is now abscised, and the sutures drawn together and tied.

The foregoing and other methods of abscission are only applicable where the tension is either low or normal. If it be high, the liability to intra-ocular hemorrhage during the operation makes enucleation, evisceration, or Mules' operation more suitable pro-




Fig. 58.

ceedings. Indeed I, and probably most surgeons, would now employ one of the two latter operations in all these cases.

Evisceration was proposed about the same time by Professor Graefe, of Halle,[16] to prevent death from meningitis after the removal of suppurating globes, and by Mules,[17] of Manchester, chiefly to take the place of enucleation in cases of sympathetic ophthalmitis. Many surgeons are opposed to its employment in those cases, but for staphyloma of the cornea it cannot meet with any such opposition.

The cornea is removed by making an incision with a Graefe's knife, so as to include one-half of the corneo-scleral margin, and by completing the circumcision with scissors. All the contents of the globe are then evacuated by means of Mules' scoop, care

being taken to remove the chorioid unbroken by carefully peeling it from the sclerotic margin backwards, until it is only held at the lamina cribrosa. The scoop is then used to lift out the separated unbroken chorioid and then other contents of the globe.

Finally, the margins of the sclero-conjunctival wound are drawn together with a few points of suture. The whole proceeding should be done with strict antiseptic precautions, chief among which is the free use of irrigation with a 1 in 5000 solution of corrosive sublimate before, during, and after the operation, the interior of the globe being most carefully washed out with the solution in a full stream. The result is a fairly good and freely movable stump for the application of an artificial eye.

Mules' Operation.—This proceeding—a modification of the foregoing—was also proposed by Mules [18] for cases of threatened sympathetic ophthalmitis, and, like simple evisceration, has not yet met with universal acceptance in those cases, because many fear that it does not afford sufficient protection against sympathetic ophthalmitis. I do not altogether participate in this feeling. In cases of staphyloma, however, and in some other conditions where the questions of sympathetic ophthalmitis in the other eye, or of a new growth in the eye to be operated on, do not enter into consideration, no proceeding is, in my opinion, more satisfactory, at least in young persons, than this beautiful one of Mules'. The prothesis it gives is almost perfect. Its object is to provide a still better stump for the artificial eye by the insertion into the scleral cavity of a hollow glass ball, called an artificial vitreous humor. It is performed as follows:

The cornea is removed—the conjunctiva having first been freed from the scleral edge towards the equator of the eyeball—and the contents of the eyeball evacuated, as in simple evisceration. The opening is now enlarged vertically, to admit of the introduction of one of the glass spheres. This introduction is best effected by means of a special instrument designed for the purpose by Mules. The spheres are made in several sizes to suit different cases, and it is well not to use the largest which will fit into any given eye. The sphere having been inserted, the margins of the sclerotic opening are united vertically by some points of interrupted suture, for which purpose I prefer silk to catgut, as the latter is apt to undergo absorption before complete union has taken place. The conjunctival opening is then closed by another set of sutures placed at right angles to the sclerotic line of closure. Similar antiseptic

precautions are required as in simple evisceration, and care must be taken that all bleeding in the cavity has ceased before the glass sphere is inserted. Before the lids are closed the anterior surface of the globe is well covered with boric acid or xeroform. A firm antiseptic bandage is applied. I do not dress the eye for forty-eight hours, and subsequently once every twenty-four hours, using the sublimate solution freely. There is generally some reaction, consisting of chemosis, swelling of the eyelids, and pain, and sometimes these symptoms are very marked, especially if too large a sphere has been employed. In the course of a week or so this all passes off, and a very perfect stump is obtained.

The danger that the glass sphere may get broken by a blow upon the eye has been put forward as an objection to this method. No doubt it is an accident which may occur, and would then necessitate the enucleation of the eye; but no case of the kind has as yet been recorded, although the operation has been in use for




Fig. 59.

eighteen years. Silver spheres, instead of those of glass, have been sometimes employed to obviate the danger referred to.

I can recommend this procedure, and I use it frequently. With a well-fitting glass eye, the cosmetic result it gives is infinitely better than that produced either by complete enucleation or by evisceration of the eyeball. It is, I think, more uniformly successful in young people than at more advanced ages, and, therefore, I do not recommend it for persons over twenty-five. To insure success an important point is to take care that the glass globe be not too large—it should be an easy fit for the cavity of the sclerotic. In case the sutures give way, and the sclerotic opening gapes, an attempt may be made to reclose it with new sutures, but the attempt is not often successful. As a rule the glass globe must in that event be removed, and the case then becomes one of simple evisceration.

Conical Cornea, or **Keratoconus.**—In this the cornea is al-

tered in shape to that of a cone. The change is due to a gradual and slowly advancing atrophic process in the cornea, at or near its center, in consequence of which the normal intra-ocular tension acts on it so as to distort it into the form represented in Fig. 59. If the apex be touched with a probe its extreme thinness may be ascertained. Tweedy [19] holds that there may be some congenital weakness in the center of the cornea as the result of its mode of development, while Panas [20] and Elschning [21] are disposed to seek the cause of keratoconus in a chronic degeneration of the membrane of Descemet. The cornea remains clear, except sometimes just at the apex of the cone, where a slight nebula may be present. The position of the apex of the cone is often not quite central, and is then most commonly either in the lower outer or lower inner quadrant of the pupil. The condition is easy of diagnosis in its advanced stages by mere inspection of the cornea, especially in profile, but in its commencement it may not be so readily detected.

In the early stages, when light is thrown on the cornea from the ophthalmoscope mirror, as in retinoscopy, the corneal reflex will be noticed to be smaller at the center, owing to the greater curvature there. Moreover, a dark shadow, circular or crescentic in shape according to the incidence of the light, appears between the corneal margin and the corneal center; and, finally, when the fundus is examined its details will be seen distorted.

The process begins in early adult life, progresses slowly, never leads to rupture or ulceration of the cornea, and, finally, after many years, ceases to progress, but does not undergo cure. Both eyes are apt to become attacked, one after the other. The disturbance of vision is very great, owing to the extreme irregular astigmatism produced.

Treatment.—In the early stages, or in slight cases, an improvement in vision may be obtained by means of concave, spherical, or sphero-cylindrical glasses; for, as is evident, the change in shape of the cornea must cause the eye to become myopic. The refraction of the central portion of the cornea may be ascertained by retinoscopy, with the aid of a stenopeic disc in the trial frame, as recommended by Mackay.[22] At a later period these glasses are of little use. Hyperbolic lenses have been employed, but, although they may raise the acuteness of vision, there are obvious difficulties in the way of the practical every-day use of them. A stenopeic slit renders assistance in some cases.

Glass shells, which are known as contact glasses, have been introduced by Fick for the temporary relief of irregular refraction; they are worn in contact with the eye, and may enable some patients to work for hours at employments which they could not otherwise carry on.

A few cases are reported in which the keratoconus was much reduced and vision greatly bettered by instillations of eserin and the application of a pressure bandage, continued for several months.

But it is upon operative measures we must chiefly rely in this affection for any practically useful improvement in sight.

Von Graefe's Method consists in flattening the cornea by the production of an ulcer on the apex of the cone, and the resulting cicatricial contraction. From the surface of the cornea, a little to one side of the apex of the cone, a morsel of corneal substance is removed with a cataract knife, care being taken not to open the anterior chamber. On the second day after this proceeding the wound is touched with mitigated lapis (solid), and this is repeated every third day for a fortnight or three weeks. Paracentesis of the anterior chamber is then performed through the floor of the ulcer, and the aqueous humor is evacuated every second day for a week, after which the healing process is allowed to take its course. A bandage must be worn during the whole course of the treatment. Finally, when the contraction and consequent flattening are completed, a narrow optical iridectomy may be necessary, in consequence of the central, or almost central, and rather intense corneal opacity.

In Bader's Method a small elliptical flap of the cornea at its apex is removed, and the margins are brought together by one or two fine sutures. The sutures are omitted by many surgeons as useless, and as liable to cause irritation. Opinion is divided as to whether the ellipse should lie vertically or horizontally in the cornea. An anterior synechia is unavoidable in a large number of the cases, and a subsequent optical iridectomy is always required. I have myself no experience of this operation, but it is said to be attended with unusual risk of suppuration of the cornea, going on to destruction of sight.

Sir William Bowman's Method consisted in cutting a disc on the apex of the cornea with a small trephine, and then severing this disc with forceps and cataract knife. Cicatrization of the wound produces the desired flattening of the cone.

Mr. Stanford Morton informs me that he employs, with satisfactory results, a modification of an operation described by Higgins.[23] He excises an elliptical piece from the apex of the cone by transfixing it from above downwards, about midway between its base and apex, with a long narrow rigid Graefe's knife, of which the edge is directed forwards and inclined slightly to the right. The knife is pushed onwards until it cuts its way out a little to the right of the apex of the cone, the aqueous escaping at the same time. The flap thus formed is then lifted up well by a fine forceps, and the knife—with its edge now turned to the left—is passed beneath the points of the forceps, and by cutting forwards the flap is excised. The sides of the wound should be steep and incline towards each other at an angle of 30° to 45°. The apex of the cone being usually somewhat downwards and inwards, or outwards, from the center of the cornea, it is important, in order to prevent an anterior synechia, to avoid carrying the incision too far downwards. The eye should be well soaked with atropin both before and after the operation, and a firm compress kept on for some time after the wound has healed.

Multiple puncturings of the apex of the cone with a fine cataract needle have been employed. The summit of the cone is transfixed from three to six times at each sitting, and this may be repeated at intervals of two weeks or more. The first effect of the punctures is to allow some of the aqueous humor to escape, and then the eye is firmly supported with a bandage. The pupil is kept under the influence of eserin. Eventually a network of cicatricial tissue forms, which flattens the cone without giving rise to much corneal opacity.

A satisfactory proceeding, and one now very generally adopted, is the application of the electro- or thermo-cautery at a red heat to the apex of the cone. By this means a contracting cicatrix is produced, which brings about a general flattening of the cornea, while the operation is practically free from risk. The cauterization must be strictly confined to a small area at the apex of the cone, and the cornea should not be perforated with the cautery. The operation may be repeated over the same area at intervals of ten to fourteen days, to bring about a more intense cicatrix.

Sir A. Critchett lays much stress on the graduated application of the cautery. He first applies the cautery at a black heat to the whole area intended to be cicatrized; within this area a little more

is destroyed at a slightly increased heat, while the very apex is touched with a cautery at a dull red heat. One sitting is sufficient.

After the cicatrization following on cauterization is completed, the scar is to be tattooed, and an optical iridectomy will usually be required, especially if the cone has been quite central. The cases in which the apex of the cone has an eccentric position are those most benefited by cauterization, because the resulting scar interferes less with vision than where it is central.

Tumors of the Cornea.

Primary tumors of the cornea are extremely rare. Epithelioma and sarcoma have their origin not in the cornea, but in the limbus conjunctivæ (p. 129). Dermoid tumors are usually seated partly on the conjunctiva and partly on the cornea (p. 128). Yet a very few cases of papilloma, epithelioma, and fibroma are recorded as taking their origin in the cornea. Corneal cysts also occur.

Injuries of the Cornea.

Foreign Bodies in the Cornea, such as morsels of iron, stone, coal, etc., are amongst the most common accidents of the entire body. The pain caused by these foreign bodies is very considerable, as can be understood, when the rich nerve supply of the cornea is remembered.

The dangers which may follow on the presence of a foreign body in the cornea depend partly upon the infection or non-infection of the foreign body, and partly upon the depth at which it is buried in the cornea. The deeper a foreign body lies the more difficult will be its removal, and the greater must be the laceration of the cornea caused by its removal. A foreign body which carries infection will be more likely to set up serious inflammatory reaction than one which is aseptic or nearly so. For this reason it is important to ascertain, if possible, the origin of the foreign body, although an apparently aseptic origin must not set all fear on this point at rest. Atoms of hot metal are from their temperature aseptic.

Many foreign bodies are so small as to defy detection until the cornea is searched with the oblique light—an aid which should al-

ways be made use of whenever the symptoms or history in the remotest way suggest the presence of a foreign body.

A foreign body which lies quite superficially in the epithelium is easily removed by gentle wiping with a clean camel's-hair pencil, or soft cloth. Those which lie deeper require instrumental interference.

The eye having been thoroughly cocainized, the patient is seated, and leans his head against the chest of the surgeon, who stands behind him. With the index-finger of the left hand the surgeon then lifts the upper lid of the injured eye, pressing the margin of the lid upwards and backwards, while with the second finger he depresses the lower lid in a similar manner; and between these two fingers he can, to a great extent, control the motions of the eyeball. The foreign body is now to be pricked out of the cornea with a special needle, with as little injury of the general surface as possible, the patient all the while directing his gaze steadily at some given point. If the foreign body be deep in the layers of the cornea, it must be dug out, as it were; and a minute gouge is made for this purpose. In the case of a morsel of iron or steel which has lain for some time in the cornea, a small ring of rust will be seen surrounding the late seat of the foreign body after its removal. This rust-ring is in the true cornea, and must be carefully scratched away, or the recovery, by necrosis of the affected part, will be much slower, and the resulting opacity much greater.

Care must be taken not to infect the cornea in the removal of a foreign body, and, consequently, thorough antiseptic precautions, especially as regards the instrument used, must be taken. After the foreign body is removed, the place where it was seated should be washed with a 1 in 5000 solution of corrosive sublimate. A bandage is worn until the epithelium is regenerated—*i. e.*, for a day or two.

Every surgeon and general practitioner should possess the two small instruments required for the removal of superficial corneal foreign bodies, and should understand the use of them.

The magnet is of no use whatever for the removal even of superficially seated foreign bodies of steel or iron in the cornea.

Sometimes a foreign body in the cornea will be so long as to protrude somewhat into the anterior chamber, and there is danger that, in the attempts at removal, it may be pushed farther on, and fall into the anterior chamber. Here it is necessary to pass a

keratome through the cornea, and behind the foreign body, so as to provide a firm base against which to work, or the keratome may be made to push the foreign body forwards.

The wing-cases of small beetles and scales of seeds may get into the eye, and adhere to the cornea by their concave surface for several days.

Simple Losses of Substance or **Abrasions** of the surface of the cornea, involving the most anterior layers of the true cornea, or perhaps merely the epithelium, are very common from rubs or scratches with branches of trees, finger-nails, etc., etc. These injuries heal readily by protecting the eye with a bandage; but when neglected, or if septic matter have been introduced when the injury occurred, or if it be present in the conjunctiva or lacrimal sac, these losses of substance are capable of forming the starting-point of corneal abscess (p. 161), ulcus serpens (p. 151), etc.

A remarkable condition known as **Recurrent Abrasion,** or **Disjunction of the Cornea** (and also as Traumatic Keratalgia, and Recurrent Traumatic Keratalgia) is sometimes observed to follow upon abrasions of the cornea. Healing of the primary lesion having taken place in an apparently normal manner, the patient, after an interval of days, weeks, or even months, is seized with severe pain, similar to that experienced on the occasion of the injury, on awakening in the morning. On examination of the eye a loss of the epithelium, which may be greater or less in extent than was the primary loss, is found at the seat of the original lesion, or it may be elsewhere on the cornea. Or, more rarely, instead of a loss of epithelium, the latter may be raised up like a vesicle, or bulla. Examination of such cases has shown that the epithelial covering of the whole cornea may easily be lifted off with a forceps; in short, that the cohesion between epithelium and Bowman's membrane all over the cornea has become imperfect. Care of the eye by means of a bandage enables the renewed loss of substance to be rapidly repaired; but after a period of quiescence, another attack takes place on awaking in the morning or in the course of the night, and such attacks may continue to recur even for several years. It is characteristic of the affection that the attacks always take place on awaking—a circumstance which is explained by the slight adhesion between palpebral conjunctiva and corneal epithelium formed during sleep, so that on the lifting of the eyelid the loosened epithelium is torn away or lifted in a bulla-like shape. Examination of the corneal sur-

face, after one of these attacks, by the usual methods (focal illumination with magnifying glass, and fluorescin), may fail to reveal the presence of a loss of substance, and then, according to Szili,[24] it may be discovered by means of transmitted light from a plane mirror, which will display the defect as a black mark.

The cause of disjunction of the cornea has not been ascertained. The condition has occasionally been noted in cases where no trauma had occurred, and this suggests the possibility of the eyes in the traumatic cases being such as already had a tendency to the derangement, whatever it may be, and which became aggravated by the injury.

Schoeler[25] suggests that the tendency to recurrence is due to very minute foreign bodies, which lie on the surface of the original loss of substance when the epithelium grows over it, and prevents a correct adaptation of the epithelium to Bowman's membrane, between which the tears are consequently enabled to percolate, and to separate off the epithelium even to a greater extent than did the original trauma.

Treatment.—Cocain, owing to the disorganizing effect it has on the corneal epithelium, should be used as sparingly as possible. A carefully applied bandage is important, and Szili states he has found dionin useful. He also has used in many cases an operative measure—namely, the removal with the forceps of the entire corneal epithelium or as much of it as would easily come away. He believes that this has had the effect of inducing a sounder union to take place.

Schoeler, acting on the theory he holds concerning the cause of the condition, brushes weak chlorin water over the loss of substance with a camel's-hair pencil, taking care at the same time to brush off all the loosened epithelium. His objct is to wipe away all small foreign bodies which may lie under the loosened epithelium preventing proper union. He follows this up with the use of atropin ointment and compresses of very weak chlorin water, and states that he has never seen a second relapse occur after this treatment.

Blows on the Eye are liable to cause corneal bullæ, the walls of which consist of Bowman's membrane and the epithelium. In some cases these bullæ contain blood derived, no doubt, from the ruptured canal of Schlemm. Such bullæ may also form after burns with lime, etc.

Another condition caused by blows on the eye is hemorrhagic discoloration of the true cornea, which presents a greenish or a reddish-brown color in the cornea. Hemorrhage in the anterior chamber is always present at first. At first, too, the discoloration occupies the whole cornea, and after a time begins to clear up from the margin towards the center. The prognosis for vision is good, if the eye is otherwise sound, but the absorption of the coloring matter in the cornea is excessively slow, and as much as two or three years or more may elapse before the process is complete. Treacher Collins [26] has ascertained that the peculiar discoloration in these cases is due to the presence of hematoidin, which he thinks enters the cornea from the hemorrhage in the anterior chamber through Descemet's membrane by a process of diffusion. He did not find any red blood-corpuscles in the cornea.

Burns of the Cornea.—Burns of the cornea from lime are not uncommon. The lime acts as a caustic, and destroys the cornea more or less deeply, with resulting more or less intense permanent cicatricial opacity. The lime, moreover, enters into chemical combination with the corneal tissues in the form of the albuminate of lime, which causes further opacity.

Treatment.—As soon as possible after lime has entered the eye it should be removed as thoroughly as possible by means of forceps and free washing out with water; or, better still, with saturated solution of sugar, which forms, with whatever loose lime may be present, an insoluble substance that may be readily removed.

The removal of such albuminate of lime as remains fastened deeply in the cornea is difficult to effect. When the irritation of the eye has subsided, the method proposed by Guillery [27] may be employed. It consists in the application to the cornea, by means of an eye-bath, of a warm 2 per cent. solution of chlorid of ammonium, which dissolves out the lime. The bath is to be used several times a day for half an hour each time, and the strength of the solution may be gradually increased, according as the patient can bear it, up to 20 per cent.

Perforating Injuries of the Cornea.—In these cases the injury done is rarely to the cornea alone, and at the first inspection the attention of the surgeon is occupied less with the state of the cornea than with the question as to whether, and to what extent, deeper parts of the eye (lens, vitreous humor, etc.) are involved. Another very important point which has often to be decided is whether or not the foreign body which has perforated

the cornea is contained in the eye. But these matters belong to future chapters.

A perforating wound of the cornea which does not involve any other part is serious in proportion to its extent, and to the probability of its being infected. Every perforating corneal wound is followed by loss of the aqueous humor, which flows away through it, and by consequent collapse of the anterior chamber; but in itself this is not a serious event. Short wounds close almost at once (and through them indeed very little of the aqueous humor may flow off), the aqueous humor is rapidly restored, and no harm is done to the eye beyond a slight opacity, which, if in the pupillary area, may cause some defect of vision; or, should the wound be situated more peripherally, and should the iris have lain against the cornea for a while, an anterior synechia may form.

Long wounds, which may even occupy the cornea in its entire diameter without directly involving any other organ of the eye, are almost certain to be complicated by prolapse of the iris between the lips of the wound, and when healing takes place the prolapsed portion becomes permanently incarcerated in the cicatrix. At the least this incarceration, if allowed to remain, causes irregularity in the curvature of the cornea, and consequent irregular astigmatism. But it may be the starting-point of a staphyloma of the cornea, it may become the cause of glaucomatous intraocular tension, or, if at any time a slight trauma with loss of substance of its surface should occur, it may take on septic inflammation, which may spread rapidly to the deeper uveal structures, leading to panophthalmitis and loss of the eye.

Treatment.—In small uncomplicated perforating wounds, without prolapse of iris, the aqueous humor being still wanting, atropin should be freely used if the wound be towards the center of the cornea, or eserin if it lie toward the periphery, with the object of preventing adhesions of the iris to the posterior aspect of the wound, and a bandage should be applied to the eye.

In recent injuries of this kind, in which there is a prolapse of the iris, the latter, if not very large, may sometimes be reposed with a spatula or fine probe, aided by the action of atropin or eserin, according to the position of the wound. But in many instances this attempt will prove futile, while in those in which there is suspicion of septic infection, it is unwise to make it at all. In either circumstance the prolapsed portion of iris should be snipped off at its base. It is not enough to abscise a portion of the sum-

mit of the prolapse. The prolapsed iris should be seized with an iris-forceps, drawn forward so as to loosen any adhesions between it and the lips of the wound (or the adhesions may previously be separated by a probe passed round the edges of the wound), and cut off close to the cornea. This affords the best chance of the iris receding into the anterior chamber without any of it adhering in the cornea.

In cases which are not recent the adhesions between cornea and prolapsed iris will have become so firm that it will not be possible to separate them by any means, and the prolapse must be left to become cicatrized over, attention being directed to keep the tension of the eye low by means of eserin, punctures made in the prolapse, and ultimately an iridectomy, so as to produce a firm and flat cicatrix.

Opacities of the Cornea.

Nebula, Macula, Leukoma.—These terms are applied to opacities of varying degrees in the cornea, which are the result of some diseased process, or which are consequent upon an injury. The first term is used for very slight opacities, often discoverable only with oblique illumination. Macula indicates a more intense opacity, recognizable by daylight. Leukoma is a completely non-translucent and intensely white opacity, the result almost always of an ulcer, which has destroyed most of the true corneal tissue at the affected place; indeed, it is often the result of an ulcer which has eaten its way through the cornea. In these latter cases the iris may have become adherent in the corneal cicatrix, and then the term adherent leukoma is employed.

Very often eyes with an old-standing nebulous condition of the cornea are myopic. It is probable that this myopia is produced by the habitual close approximation of objects to the eye, owing to the diminished acuteness of vision from the opacity of the cornea.

Treatment.—Little or nothing can be done to reduce these opacities. In slight and fresh cases massage with the yellow oxid of mercury ointment may render them less intense.

In case of a nebulous cornea a stenopeic apparatus often improves the sight. This consists of a metal plate with a small central hole or slit, which is placed before the patient's eye in a spectacle-frame; and by this arrangement a large portion of the

rays which pass through irregular parts of the cornea, and which merely confuse the sight, is cut off. Should the opacity be dense, and situated in the center of the cornea, portion of the margin having remained clear, an iridectomy will in some instances improve the sight.

The Operation of Tattooing was first proposed by de Wecker, and is a valuable proceeding for improvement of the appearance of the eye in cases of leukoma. It is also an extremely useful method for improvement of the sight where the nebula occupies only part of the pupillary area of the cornea. In these cases much disturbance of sight is caused by the dispersion of the light which makes its way through the nebula; and when by tattooing the scar all light is prevented from getting through, brighter and distincter vision is enjoyed with the part of the cornea, opposite the pupil, which is absolutely clear.

In the case of a leukoma either the whole surface of the leukoma may be tattooed or only part of it—*e. g.,* its center, in order to represent a pupil.

The material used is fine Indian ink rubbed into a very thin paste. The eye having been cocainized, the leukoma is spread over with this paste, and then covered with innumerable punctures by means of de Wecker's multiple tattooing-needle, each stab of which carries into the cicatricial tissue some of the black pigment. The coloration continues sufficiently intense for some months, but then often begins to get pale, owing, probably, to the pigment falling out of the punctures. A method of tattooing, by which the pigmentation lasts longer, is performed with de Wecker's single-grooved needle. The pigment is placed in the groove of the instrument, which is then passed into the leukoma, a long canal being made in a plane parallel to its surface. On withdrawal of the needle the pigment remains behind. A large number of such canals must be made in close proximity to each other until the desired intensity of color is obtained. Some operators remove the corneal epithelium over the part to be tattooed, in order to facilitate the entrance of the coloring matter into the true cornea.

In cases where the whole cornea is leukomatous, and, consequently, where no restoration of sight can be obtained by means of an artificial pupil, *Transplantation of a Portion of Clear Cornea* from a rabbit's eye, or from a freshly enucleated human eye, has been repeatedly performed by ophthalmologists in various parts

of the world. Very many of these operations have been perfectly successful in a surgical sense—*i. e.*, in so far as the healing-in of the transplanted flap was concerned; but, with a few exceptions, they all ended in disappointment, in consequence of the flap not retaining its transparency. In the course of a week or two the transplanted portion invariably becomes as opaque as the leukoma had been before. The mode of proceeding consisted in removing a portion of the leukoma with a trephine, and in then cutting a disc with the same instrument out of the clear cornea to be utilized, and inserting it into the opening in the leukoma.

Various theories were formed to account for the occurrence of the opacity in the transplanted flap, but into these it is unnecessary to enter here. Von Hippel [28] came to the conclusion that the onset of the opacity was due to the entrance of the aqueous humor into the substance of the cornea, owing to the solution of continuity in its posterior epithelium; Leber's experiments [29] having shown that, unless this epithelial layer be intact, the transparency of the cornea cannot be maintained. Von Hippel, acting on this theory, applied a trephine to the leukoma as deep only as the posterior elastic lamina, and then dissected off the superficial layers contained within the ring, leaving the posterior elastic lamina and posterior epithelium. With the same trephine he then excised a disc of its entire thickness from a rabbit's cornea, and applied it to the wound. Iodoform was dusted over this, and a bandage applied. Healing took place readily, and twenty months afterwards the flap continued transparent, and vision $= {}^{20}$. Von Hippel has had some other successful cases.

Arcus Senilis.—This is a change which is developed in the cornea without previous inflammation. It presents the appearance of a grayish line a little inside the margin of the cornea and all round it, most marked above and below, and never advancing farther towards its center. It is most common in elderly people, but is sometimes seen in youth, and even in childhood. No functional changes are caused by it, nor does it interfere with the healing of a wound which may be made in that part of the cornea. Arcus senilis is caused by a hyaline degeneration of the corneal cells and fibrillæ, and is not a sclerosis, as is stated by some authors.

References.

[1a] "Die Entstehung der Entzundung." Leipzig, 1891.

[1] "Ueber die Verwertbarkeit der Bindehaut in der praktischen und operativen Augenheilkunde."

[2] "Ophthalmiatrischen Beobachtungen," p. 107.

[3] "Von Graefe's Archiv," xii. 2. p. 250.

[4] "Trans. Ophthal. Society, U. K.," xiii. p. 45.

[5] "Von Graefe's Archiv," xxxiv. 4. p. 250.

[6] "Bericht der Ophthalmologischen Gesellschaft." Heidelberg, 1889.

[7] "Von Graefe's Archiv," xxxviii. 1. p. 160; and xxxix. 2. 199.

[8] "Archives d'Ophtalmologie," xiii. 4. p. 193.

[9] "Von Graefe's Archiv," xxv. 2. p. 285. See also *Ibidem,* xlii. 1. pp. 114, 131.

[10] "Bericht der Ophthalmologischen Gesellschaft." Heidelberg, 1898, p. 71.

[11] "Zeitschrift für Augenheilkunde," II., pp. 235, 354.

[12] "Von Graefe's Archiv," xlvi. 1. p. 85. See also *Ibidem,* liii. p. 423, and "Zeitschrift für Augenheilkunde," viii. p. 340.

[13] "Textbook of Ophthalmology."

[14] "Bericht der Ophthalmologischen Gesellschaft." Heidelberg, 1897, p. 53.

[15] "Chirurgie Oculaire," p. 188.

[16] "Centralbl. für Augenheilkunde," 1884, p. 378.

[17] *Ibidem,* 1884, p. 378; 1885, p. 32.

[18] "Trans. Ophthal. Society, U. K.," v. p. 200.

[19] *Ibidem,* xii. p. 67.

[20] "Archives d'Ophtalmologie," 1885, p. 348.

[21] "Klinisch. Monatsblätter für Augenheilkunde," 1894, p. 25. See also "Bericht der Ophthalmologischen Gesellschaft." Heidelberg, 1898, pp. 75, 78.

[22] "Ophthalmic Review," 1893, p. 317.

[23] "British Medical Journal," 1880, p. 623.

[24] "Von Graefe's Archiv," xli. 3. p. 486.

[25] "Centralblatt für Augenheilkunde," 1901, p. 161.

[26] "Trans. Ophthal. Society, U. K.," xv. p. 69.

[27] "Archiv für Augenheilkunde," xliv. p. 311.

[28] "Bericht der Ophthalmologischen Gesellschaft." Heidelberg, 1886, p. 54.

[29] "Von Graefe's Archiv," xix. 2. p. 87.

CHAPTER VII.

DISEASES OF THE EYELIDS.

Erythema, erysipelas, phlegmonous inflammation, and abscess are all liable to attack the eyelids, but require no special observations in this work. It should merely be stated that erysipelas of the eyelids may extend to the connective tissue of the orbit, and ultimately give rise to atrophy of the optic nerve.

Eczema.—This is very often seen on the eyelids, most frequently in connection either with general eczema of the face or with phlyctenular ophthalmia. The lacrimation in phlyctenular ophthalmia increases the eczema, which then, by causing contraction of the skin of the lower lid, produces eversion of the inferior punctum lacrimale, and this, in its turn, causes increased lacrimation, and thus a vicious circle is set up.

Atropin infiltration of the eyelid, from long use of solution of atropin in some persons, is often accompanied by a moist form of eczema of the lids and face.

Treatment should consist in the daily removal of the scabs in such a way as to cause no bleeding of the surface underneath; and for this purpose a warm solution of bicarbonate of potash is useful. The place should afterwards be well dried, and painted with a strong solution of nitrate of silver (gr. xx ad ℥j), and a boric acid ointment (gr. xxx ad ℥j), or the following, applied over this: Ol. Cadin, *m* xv; Flor. Zinc, gr. xx; Lanolin, ℥ij.—M. If the inferior lacrimal punctum be everted, the canaliculus should be slit up.

Herpes Zoster Ophthalmicus is a herpetic eruption of the skin in the region supplied by the ophthalmic division of the fifth nerve of one side.

Occasionally in the same case the second division of the fifth nerve may be affected, and, yet more rarely, the third division as well. One or two cases, too, have been published in which the zoster affected each side of the face.

But by far the most common case is the simple herpes zoster ophthalmicus, in which only the region supplied by the ophthalmic

division of the fifth nerve is affected, and of this region it is usually that portion alone which pertains to the supra-orbital and infratrochlear branches that is involved. The number of vesicles varies much; there may be but one, or there may be several, or they may be so numerous as to become confluent.

The appearance of the eruption is often preceded by a feeling of general discomfort, gastric disturbance, and high temperature. Yet more commonly is the eruption preceded by supra-orbital neuralgia, which is often of intense severity. This pain usually continues, but may cease, after the eruption comes out, and sometimes it persists even for months after the eruption disappears. Photophobia, due to the irritation of the fifth nerve, is not uncommon at the commencement of the affection. Along with the appearance of the herpes the skin of the forehead becomes red and swollen, and the appearances are often mistaken for erysipelas, but the strict limitation of the eruption by the middle line of the forehead is of itself sufficient to indicate the diagnosis. The upper lid is somewhat edematous and red, and droops a little over the eye, and this is much more marked when the skin of the eyelid itself is the seat of vesicles.

The contents of the vesicles soon become purulent, the latter gradually dry up, and form crusts, under which are more or less deep ulcers, and as these penetrate the corium they, on healing, leave permanent scars, which at first are red, and later become of a glistening white. The entire eruptive process lasts about three weeks, and when it is completed the sensibility of the affected skin remains dull for a considerable time. Herpes zoster ophthalmicus is more common in advanced life than in youth, but it may appear at any age, even as young as the sixth month after birth.

The disease is not associated with danger to the eye, unless keratitis comes on, or, what is much more rare, unless iritis, cyclitis, or chorioiditis appear. Hutchinson laid down the rule that the cornea does not become affected unless the eruption appears on the region (the side of the nose) supplied by the nasociliary nerve, this being the nerve which gives the long root to the ciliary ganglion, as well as the long ciliary nerves. The rule holds good in most instances, but there are many exceptions to it. The conjunctiva is almost always slightly chemotic and injected, or there may be true conjunctivitis; but vesicles are not often seen on it.

There is considerable variety in the forms of keratitis liable to

occur in herpes zoster ophthalmicus—viz., herpetic vesicles, phlyctenulæ, bullæ (any of which may go on to ulceration), superficial opacity without loss of substance, and parenchymatous opacity, either diffuse or punctate. The superficial opacities without loss of substance may disappear completely. Parenchymatous opacity either clears away altogether, or remains as a slight nebula; while ulceration leaves, at the least, some opacity; or, if it become septic, may seriously endanger the eye. Anesthesia, more or less well marked, attends the corneal affections, and remains for a long time after they recover.

Iritis is very uncommon in herpes zoster ophthalmicus, and is usually of a mild type, and iridocyclitis and chorioiditis are still more uncommon. They seldom occur unless keratitis is also present, and, like the keratitis, for the most part only when the cilionasal nerve is implicated.

Herpes zoster ophthalmicus is due to an inflammatory process in the Gasserian ganglion, as Head and Campbell have shown, and in the opinion of these authors the skin eruption is caused by intense irritation of the ganglion cells. The lesion in the Gasserian ganglion is similar to that found in the posterior root ganglion in zoster of the trunk and limbs. Head and Campbell [1] believe the affection to be an acute specific disease—a view suggested by the facts that it occurs in the course of recognized infective diseases, that it occurs endemically and epidemically, and that it rarely occurs a second time. It is probable that the affection may also have a toxic origin, as when arsenic has been taken for a long time, and in carbonic oxid poisoning.

Treatment.—It is doubtful whether treatment has any influence in curing or in controlling the severity of an attack of herpes zoster ophthalmicus. Quinin in full doses should be given, and a 1 per cent. cocain ointment made with equal parts of vaselin and lanolin should be smeared lightly over the affected part. Complications in the cornea or uveal tract are to be dealt with on the principles laid down in the chapters on diseases of those organs. The patient, unless the attack be a very mild one, should remain in bed.

Primary Syphilitic Sores occur on the eyelids, usually near the margin of the upper or lower lid or at the inner or outer canthus. The first appearance is generally a small red swelling which the patient calls a "pimple," and which ulcerates and becomes characteristically indurated about its base. The margin of the ulcer

is clean-cut, and its floor somewhat excavated and covered with a scanty grayish secretion. Occasionally there is no ulcer present, but the entire lid is swollen, greatly indurated, purple, and shiny; and then the diagnosis may be rendered difficult. The pre-auricular and submaxillary glands are almost always swollen; and this is a valuable, although not altogether positive, diagnostic sign, as it is seen also in tubercular diseases of the conjunctiva. The occurrence of the sore is followed by the usual constitutional symptoms of syphilis. Very rarely is there any permanent damage done to the eyelid.

The most common modes of infection are by a kiss from a syphilitic mouth, or by a dirty finger.

Treatment.—Locally, sublimed calomel by Kane's method, dusting with finely powdered iodid of mercury, or the black wash may be used; while the usual general mercurial treatment is employed.

Secondary Syphilis gives rise to ulcers on the margins of the lids, to loss of the eyelashes (madarosis), and to the secondary skin affections which attend it in other parts of the body.

In **Tertiary Syphilis** ulcerating gummata of the lids sometimes are seen, accompanied by remains of previous iritis or keratitis.

Vaccine Vesicles on the eyelids are produced by accidental inoculation at the intermarginal part of the lid; or on the outer surface of the lid, if the skin be abraded by the finger-nail or otherwise. Sometimes the vesicle develops into a large ulcer with yellowish floor and hard and elevated margin. There is much pain, much swelling of the eyelid, and chemosis.

Although distressing for a week or so while it lasts, the affection is not a dangerous one, further than that a cicatrix in the skin is left behind, and the eyelashes at the affected part are lost.

Treatment.—A warm chlorate of potash lotion (gr. v ad ℥j.) is the best application.

Rodent Ulcer (Jacob's Ulcer).—This disease commences as a small pimple or wart on the skin near the inner canthus, or over the lacrimal bone, as a rule; but it may also originate in any other part of the face. The scab or covering of the wart is easily removed, and underneath is found a shallow ulcer with a well-defined, indurated margin; the skin surrounding the diseased place being healthy, and continuing so to the end of the chapter. The progress of the disease is extremely slow, extending over a great number of years, and in the early stages the ulcer may even seem to heal for a time, but always breaks out again. In mild

cases the ulceration may remain superficial; but more usually it strikes deep, in the course of time eating away every tissue, even the bones of the face and the eyeball. The latter is often spared until after the orbital bones have gone.

The disease is an epithelial cancer of a non-malignant or purely local kind. There is no tendency to infiltration of the lymphatics. It is rarely seen in persons under forty years of age.

Treatment.—Extirpation of the diseased part affords the best chance of relief for the patient. Recurrence of the growth is the rule, but this should not deter from operative measures, nor even from the renewal of them, as they afford much comfort to the patient and prolong his life. Even in advanced stages operation is frequently called for. The application of chlorid of zinc, or of the actual cautery, should be employed, after the disease has been as thoroughly removed with the knife as is possible.

Bergeon's Treatment.—This consists in the internal administration of 5 grains of chlorate of potash three times a day, with the local application of a saturated solution of chlorate of potash to the ulcer, and by aid of it remarkably good cures, even though not always permanent ones, can be effected. It is well, in many cases, to scrape the ulcer before applying the solution. The process must be repeated daily, or at least every second or third day. It is certainly painful, but not unbearably so. Sometimes a green slough is produced, and when this is the case there is generally some surrounding inflammation, which should be allowed to subside before going on with the treatment. As the healing process does not begin until the diseased tissue has been removed, the progress may seem slow for the first week or fortnight; but no case resists the treatment altogether if it be persevered with. While the chlorate of potash destroys the disease, it does not act injuriously on the delicate epithelium, which begins to grow in from the margin as healing sets in, and it should therefore be continued until the whole surface has healed. Another fortunate peculiarity is that it has no effect on the normal conjunctiva, and may be used without fear if this membrane be involved in the disease.

Marginal Blepharitis (*βλέφαρον, eyelid*), or **Ophthalmia Tarsi,** is nothing else than eczema of the margin of the eyelid. It is found either as **Blepharitis Ulcerosa** (Eczema Pustulosa), or as **Blepharitis Squamosa** (Eczema Squamosa). In the former small pustules form at the roots of the eyelashes, and these, having lost their covering, become ulcers, which scab over. The whole

margin of the lid may then be covered with one large scab, in which the eyelashes are matted, and under which the lid will be found swollen, red, and moist, with many minute ulcers and pustules. Many eyelashes come away with the scab, and others are found loose and ready to fall out.

The disease is chronic, and is most commonly seen in strumous children. It is frequently accompanied by phlyctenular ophthalmia, or by simple conjunctivitis, which may have been its cause, or which promotes it by keeping the margin of the lid constantly wet.

If neglected, ulcerous blepharitis is liable to produce trichiasis by giving a false direction to the bulbs of the cilia.

Many ophthalmologists hold that blepharitis is often caused by ametropia, especially by hypermetropia or hypermetropic astigmatism, in consequence of the incessant efforts of accommodation. I cannot go so far; but it may be that, if blepharitis be once set up, such anomalies of refraction may help to keep it going.

The Treatment of Ulcerous Blepharitis consists, in the first place, in the careful removal of the scabs without causing any bleeding of the delicate surface underneath. Such bleeding indicates that the newly formed epithelium has been torn away, and it is important, therefore, to soften the scabs by soaking the eyelid with olive oil, or with a solution of bicarbonate of potash, before removing them. Any pustules found under the scab should be punctured, and all loose eyelashes taken away, and the ulcers touched with a fine point of solid mitigated lapis. The surface should then be well dried by pressure, not by rubbing, with a soft cloth, and the following ointment (Hebra) applied: R. Ol. Rusci (or Ol. Juniperi) 3ss, Hydrarg. Ammon. Chlor. gr. iv, Vaselin alb., Lanolin aa 3ij. This ointment is to be continued until healing is thoroughly established. In many mild cases a boric acid ointment (gr. v ad 3j of vaselin or of lanolin) will be found efficacious instead of the above, and a white precipitate ointment of from 1 to 2 per cent. acts well. A creolin ointment suits many cases—viz., Creolin, 1 to 5 min.; Aq., 3ij; Lanolin, 3vj.

Or, again, after the scabs and loose eyelashes have been removed as above, the margins of the eyelids may be freely bathed with a wash of ten to twenty minims of creolin to eight ounces of water, and after this the creolin ointment may be applied. I have found this method very successful. But in all cases, whatever the lotion or ointment ordered may be, the ulcers should be touched with

mitigated lapis, as above recommended, and all loose eyelashes removed.

All complications with conjunctival affections or lacrimal obstruction must be attended to, and the patient's general system carefully improved. Any error in refraction should be suitably corrected.

Squamous Blepharitis comes on after the ulcerous form has passed away; or it is found as a primary affection, especially in chlorotic women. The margin of the lid is somewhat swollen and red, and covered with loose epidermic scales. It is an extremely chronic affection.

The Treatment of Squamous Blepharitis is also an ointment of Hebra's:

℞ Emplast. Diachylon Co.,* ʒ ij; Ol. Olivar, q. s.;

or the Boric Acid ointment may be used.

Chlorosis, if present, is to have suitable remedies.

Phtheiriasis (*φθείρ, a louse*) **Ciliorum.**—The pediculus pubis occurs on the eyelashes. It gives rise to excessive itching and burning sensations, and the consequent rubbing produces excoriations of the margin of the lid. The lice occupy chiefly the roots of the eyelashes, while the shafts of the cilia are covered with their brown egg-capsules, and this gives to the cilia the peculiar appearance of being covered with dark brown powder, which enables the diagnosis to be easily made. The fully developed parasites, as well as the eggs, may be more readily seen by aid of a strong convex glass.

Treatment.—With a cilium forceps the pediculi may be to a great extent, or completely, removed, as well as some of the eggs from the cilia. This proceeding repeated daily, along with the application of mercurial ointment, or of a weak red precipitate ointment, to the margin of the eyelids morning and evening, will soon effect a cure.

Hordeolum (*hordeum,* a grain of barley) or **Stye,** is a circumscribed purulent inflammation situated at the follicle of an eyelash. It commences as a hard swelling, with more or less tumefaction and edema of the general surface of the lid, and often with some chemosis, especially if it be situated at the outer canthus. In

* Emplast. Diachylon Co. is made as follows: Emplast. Litharg. B. P., 12 parts; Cornflour, 1 1-2 part; Ammoniac, Galbanum, Turpentine, of each 1 part.

its early stages there is much pain associated with it. It gradually suppurates, and may then be punctured or allowed to open of itself.

Styes frequently come in rapid succession one after the other, and then, probably, a constitutional disturbance exists as the cause. In the earliest stage cold applications may be successful in putting back a stye, but, later on, warm stupes will hasten the suppuration and relieve the pain. Habitual constipation is a common source of hordeolum, and should be met by the occasional use of cascara sagrada, aperient mineral water, or other mild laxative. Sulphid of calcium, 1-10 gr. every hour, or 1-2 gr. twice a day, for an adult, has been recommended (D. Webster) as a specific in these cases.

Chalazion (*χάλαζα, hail*), **Meibomian Cyst,** or **Tarsal Tumor,** is probably a granuloma in connection with a Meibomian gland, and not a mere retention cyst. Micro-organisms have been found by some observers in these tumors, but what relation exists between them and the tumors is a matter upon which opinions differ. Chalazion has its origin in a chronic inflammatory process in the connective tissue surrounding the gland, which usually passes off without having attracted the attention of the patient. The tumors vary in size from that of a hemp-seed to that of a hazelnut, causing a marked and very hard swelling in the lid. They occasionally open spontaneously on the conjunctival surface, giving exit to contents which are usually viscid or grumous, but sometimes purulent.

Treatment.—No application can bring about absorption of these tumors. The lid should be everted, the tumor opened by a single incision from the conjunctival surface, and its contents thoroughly evacuated by aid of a scoop or small sharp spoon. Difficulty is sometimes experienced in finding the point in the conjunctiva corresponding to the tumor, but it is usually indicated by a dusky or grayish discoloration. Immediately after the evacuation, bleeding into the sac often takes place, and causes the tumor to remain for a day or two as large as before—a fact of which the patient should be warned. The operation may occasionally require to be repeated two or three times. The interior of the sac should not be touched with nitrate of silver; and the incision and evacuation should never be made through the skin, because more or less disfigurement from the scar would result.

More than one chalazion is often present at a time, and some people become liable to them periodically during a number of years.

Milium (*milium,* a millet seed) presents the appearance of a

perfectly white tumor, not much larger than the head of a pin, in the skin of the eyelid. It is a retention tumor of a sebaceous gland, and can readily be removed by puncture and evacuation.

Molluscum, or **Molluscum Contagiosum.**—This is a white tumor in the skin of the eyelid, which may attain the size of a pea. At its summit is a depression, which leads to an opening into the tumor, through which the contents can be pressed out. It is probably a diseased condition of a sebaceous gland, and contains altered epithelial cells and peculiar bodies, termed molluscum corpuscles, which are of a fatty nature. Many such tumors may form in the lids at the same time.

It is held by some observers that this affection is contagious, although in what way is not clear, inasmuch as experimental rubbing of the contents of a molluscum into the skin has not given rise to the tumors.

Treatment.—Each separate tumor must be evacuated by simple pressure, or after it has been opened up with a knife or scissors.

Telangiectic Tumors, or **Nevi,** of the eyelids occur congenitally.

Treatment.—Small tumors of this kind may be destroyed by touching with nitrate of silver or hydrochloric acid, or by performing vaccination on them. Large tumors may be ligatured or treated with the galvano-cautery, and electrolysis is a very effectual method in many cases.

Xanthelasma (*ξανθός, yellow; ἔλασμος, a layer*) is the term applied to yellowish plaques raised slightly over the surface of the skin, with very defined margins. The patches are generally bilateral and symmetrical, and are most frequently situated in the neighborhood of the inner canthus. The shape of these plaques is extremely irregular, and they may attain the size of a shilling or larger. The appearance is caused by hypertrophy of the sebaceous glands, with retention of their contents, and fatty degenration of the subcutaneous connective tissue.

Treatment can only consist in removal by careful dissection, and this is hardly to be recommended except in extreme cases.

Palpebral Chromidrosis (*χρῶμα, color; ἵδρωσις, sweating*).—The phenomenon of an exudation of pigment upon the eyelids, of which about fifty cases have been recorded, has given rise to much discussion. The opinion held by many is that these cases are always the result either of deception in hysterical individuals, or of accidental circumstances, such as the exposure of a patient

with seborrhœa palpebrarum to an atmosphere loaded with coal-dust or pigmentary matter, in some manufacturing district. Of the fact that the appearance has occurred under both of these conditions there can be no doubt. There would seem also to be evidence that some genuine cases of color-sweating on the eyelids



Fig. 60.

have been observed; but they must be extremely rare. The discoloration is blue or black, and occurs in the form of fine powder upon the skin of one or both eyelids of both eyes. It can be wiped off, and is said to begin to reappear after a short interval. The subjects of it have been chiefly young girls, but it has also been seen in women of advanced years, and even in middle-aged men.

The Treatment in a genuine case may consist in the application of a lotion of liq. plumbi and glycerin; and, internally, iron, quinin, and arsenic, along with the regulation of the general system, particularly in respect of any uterine derangement.

Epithelioma, Sarcoma, Adenoma, and **Lupus** are all seen in the eyelids, but require no special description here.

The condition known as **Solid Edema,** or **Elephantiasis Lymphangiodes of the Eyelids,** is well represented in the accompanying picture (Fig. 60 *) of a case under the care of Sir A. Critchett. It is a chronic tumefaction of the eyelids, most marked in the lower lids. The skin covering the swelling is smooth and pale, and suggests to the eye the skin of an edematous lid; but on palpation the swelling is found to be much more resistant than simple edema. There is, almost invariably, a history of recurring attacks of facial erysipelas. These give rise to a permanent alteration of the lymph channels, and, each attack leaving its trace, an ever-increasing hypertrophy of the tissues of the eyelids takes place.

Treatment.—Operative measures have been adopted in many instances with satisfactory results, both cosmetically and as regards the functions of the eyelids; but, unfortunately, in those cases which have remained under observation sufficiently long, the former condition gradually returned, as in Sir A. Critchett's case. Multiple punctures, collodion, pressure, etc., and many internal remedies have been tried in vain.

Clonic Cramp of the Orbicularis Muscle, or of a portion of it, is often seen, and is popularly known by the name of "life" in the eyelid. It is frequently due to overuse of the eyes for near work, especially by artificial light, or if there be defective amplitude of accommodation.

Treatment should consist in the regulation of the use of the eyes for near work, and the correction by glasses of any defect in the accommodation.

Blepharospasm, or **Tonic Cramp of the Orbicularis Muscle,** is commonly the result of irritation of the ophthalmic division of the fifth nerve by reflex action, as in phlyctenular ophthalmia and some other corneal and conjunctival affections; or from foreign bodies on the conjunctiva or cornea, etc.; or it may continue for some time after the relief of any such irritation. It occurs, also,

* Taken by permission of the Council from "Trans. Ophth. Soc., U. K.," vol. xix.

independently of such causes, and is then difficult to account for, unless as a hysterical symptom. Yet even in these obscure cases the spasm is probably often a reflex from the fifth nerve, and it will be found that pressure upon the supra-orbital nerve at the supra-orbital notch may arrest the spasm; or, if not there, then pressure on the infra-orbital, temporal, malar, or inferior alveolar branch may have the desired effect.

Treatment.—If the cause of the reflex cannot be ascertained, or if it has passed away, and if the cramp be still very distressing, stretching or resection of the branches of the fifth nerve, from which the reflex proceeds, may be tried.

Ptosis (*πτῶσις, a fall*), or **Blepharoptosis,** is an inability to raise the upper lid, which then hangs down over the eyeball. It is either congenital or acquired; and in the latter case is most usually the result of paralysis of the branch of the third nerve supplying the levator.

Persons affected with ptosis involuntarily endeavor to raise the eyelid by an overaction of the frontalis muscle. The drooping lid and elevated eyebrow give a peculiar and characteristic appearance.

The Causes of Paralytic Ptosis are similar to those of paralysis of other branches of the third pair, more especially exposure to cold draughts of air while the body is heated, and syphilis or rheumatism affecting the branch to the levator palpebræ in its course. It may also be due to cerebral disease (see chap. xvii.). The branch to the levator may be paralyzed alone, or in conjunction with other third-nerve branches, and the loss of power may be partial or complete.

The Treatment of a recent case of ordinary paralytic ptosis depends upon its cause. If this be syphilis, then a course of mercurial inunctions or of iodid of potassium; if rheumatism, salicylate of soda or iodid of potassium—with, in either case, protection of the eye and side of the head by means of a warm bandage. Cases in which these remedies have failed, and which have become chronic, often demand operative treatment. Attempts have been made, with success in some cases, to obviate the inconvenience of ptosis by giving support to the lid by wire splints worn like an eyeglass or attached to the upper edge of spectacle-frames.

Ptosis due to a cerebral lesion rarely comes within the scope of treatment.*

* The value of ptosis as a localizing symptom in cerebral disease will be treated of in chap. xvii

Operative treatment is indicated in cases of paralytic ptosis—where other measures have produced no result—in ptosis adiposa,




Fig. 61.




Fig. 62.




Fig. 63.




Fig. 64.

Fig. 61 shows intended track of wire.

Fig. 62.—Wire loop D, with needle ends passed upwards in thickness of tarsus, and turned out at B B, to be taken subcutaneously to E E by ptosis needles (C C) passed from above eyebrow.

Fig. 63 shows wire at E in eye of ptosis needle above eyebrow; D, point of needle threaded with wire projecting at middle of lid before being drawn up to E.

Fig. 64.—Completed except sinking wire knot permanently.

and in congenital cases. A very common proceeding consists in the excision of a sufficiently large oval piece of integument, its

long axis lying in the length of the lid, with the subcutaneous connective tissue and fat, and, in paralytic cases, a small portion of the orbicular muscle. The fold of integument to be abscised is seized by two pairs of forceps—one of them held by an assistant—at the inner and outer ends of the lid, and by this means the necessary size of the fold is estimated. The abscission of the fold is performed with a pair of scissors, the margin of the wound lying close to the points of the forceps. The subcutaneous tissue, etc., is then removed, and the edges of the wound drawn together by a few points of suture.

Mules' Operation.[2]—A Knapp's clamp is applied to the lid to be operated on. The edge of the tarsus is then grooved one-third of an inch in length and deeply enough to permit of perfect and easy healing over the wire bend (Fig. 62, D); the grooving should be behind the bulbs of the lashes, to prevent their being directed abnormally inwards. After grooving, the Desmarres ring may be removed, and the lid drawn tense with the lid-forceps. Next, at each end of the groove (Fig. 62, A A), one sewing needle with wire is passed upward half the width of the lid in the center of the tarsus, so as to give a strong bite for the wire bend, then thrust forwards and outwards through the tarsus (at B B) and skin, the wire being drawn through the small incision made there. The ptosis needles (C C) are now pushed deeply from above the brow, a quarter of an inch apart, into the tiny apertures made for them by a thrust of the sclerotomy knife, then underneath the eyebrow downwards to the incision (B B) in the skin of the lid, through which the wire has already passed, and their points are also turned out there (Fig. 63, D). The wire is cut from the sewing-needles, threaded through the eyes in the points of the ptosis needles, withdrawn through the small apertures in the skin of the lid, and brought out above the brow at the points of entry of the ptosis needles (E); the wire bend is then drawn firmly into the grooved edge of the lid to allow of immediate healing, a little iodoform dusted on the groove and small incisions, the wire ends shortened, the apertures dressed with collodion and iodoform, and a pad and bandage applied. After all swelling of the lid has passed, the permanent effect may be secured by tying the wire after careful adjustment of the lid, and sinking it into a small skin incision made for the purpose (Fig. 64).

Pagenstecher's Method is as follows: Its object is to enable the patient to derive more benefit from the effort of his frontalis

muscle, which he is constantly making with so little result, by transferring its action more directly to the eyelid. A needle carrying a thick ligature is entered under the skin of the forehead about half an inch above the center of the eyebrow, and passed subcutaneously as far as the margin of the eyelid at its middle point. The suture is closed, not very tightly at first, but each day somewhat more tightly, until it has cut its way through the skin. As the result of this a cicatrix is formed in the course of the ligature




FIG. 65.

which gives the frontalis much more power over the eyelid. I have tried this method, but I have not been satisfied with it.

Birnbacher's Operation is an improvement on former attempts to connect the tarsus with the frontalis by cicatrices. An incision, with its convexity upwards, is made in the skin corresponding to the upper edge of the tarsus. Three sutures with a needle at each end are passed through the upper border of the tarsus, so as to form three loops, one central and two lateral; the two needles of the central loops are passed vertically upwards under the skin, and are brought out quite close to one another in the eyebrow. The lateral loops are treated in the same way, but are made to diverge on each side from the central one, instead of being parallel. The

ends of the threads are tied over a small roll of lint, and tightened until the edges of the lids just touch when the patient closes the eye. They may be left in from twenty to twenty-five days.[3]

Panas' Method.[4]—The object of this operation is to bring about a union between the lid and the frontalis muscle by forming a flap in the former, which is fastened to the skin of the forehead and to the surface of the muscle.

Before the operation commences, and while it is in progress, an assistant applies his hand firmly to the patient's forehead, in such a way as to prevent shifting of the skin of the eyelid over the un-




Fig. 66.

derlying tissues, which would interfere with the exactitude of the proceeding.

A horn lid-spatula is inserted under the lid. Fig. 65 explains how the eyelid flap is formed. The horizontal incision along the top of the flap has a slight convexity upwards, is not quite an inch long, lies over the orbital margin, and goes through all the tissues down to the periosteum. Another incision, parallel to this one, rather more than an inch long, is made along the upper border of the eyebrow and as deep as the periosteum. The flap of skin and muscle is now dissected from the tarsus down to its ciliary border, but the suspensory ligament of the lid must not be interfered with. The bridge of tissue between the two horizontal incisions is now to be undermined without injury to the periosteum or suspensory ligament. The flap is then drawn up under the bridge by means

of the sutures (*a a'*) and secured to the upper edge of the upper incision. Inasmuch as the traction exercised by the flap when so fixed tends to produce ectropion of the lid, two lateral sutures (*b b'*) are applied deeply through the suspensory ligament and conjunctiva to the exclusion of the skin, and are attached, like the other sutures, to the upper lip of the upper incision, thus counteracting the tendency to ectropion. Fig. 66 shows the effect of the operation.

Fuchs has published [5] some cases of bilateral ptosis in elderly people, which were due, in his opinion, to primary atrophy of the




Fig. 67. Fig. 68.

l, levator palpebræ; *o*, orbicularis.

levator palpebræ muscles. The eyelids were elongated and thinned, so that the eyeball showed plainly through them. The loss of power had in each case been very slowly increasing for many years.

Congenital ptosis is generally present in both eyes. It is due in some cases to an imperfect development of the levator palpebræ, and in others to an abnormal insertion of this muscle, its tendon being attached to the tarsus too far back. Either Birnbacher's or Panas' operation may be employed, and Everbusch has proposed the following proceeding more particularly for congenital ptosis:

Everbusch's Operation for Congenital Ptosis[6] (Figs. 67 and 68).—The object of the operation is to increase the power of the levator by advancing its insertion, or rather by doubling it down over the tarsus, to which it forms fresh adhesions. Snellen's lid-clamp is applied, the plate being passed well up into the fornix; and, before the ring is screwed down, the skin of the lid is drawn down, so that its prolongation just under the eyebrow may be forced into the instrument. The skin and the underlying orbicularis are now divided in the entire width of the lid, parallel to its free margin, and at a distance half-way between this margin and the eyebrow. The skin and the subjacent muscle are then sepa-




Fig. 69.

rated up, both upwards and downwards, for 4 mm. in each direction, so that the insertion of the levator may be well exposed. A suture with a small curved needle at either end is then introduced by means of one of these needles horizontally into the tendon at its insertion, and near the center of the latter, in such a way that about 2 1-2 mm. of the tendon may be included in the suture. Each needle is now passed vertically downwards between the tarsus and orbicularis, and brought out at the free margin of the lid at a distance from each other of about 2 1-2 mm. Two more such double sutures, one in the temporal, the other in the nasal, third of the tendon, are similarly applied. The margins of the horizontal skin and muscle wound are now drawn together, and then the three

sutures are closed tightly. It is desirable to slip glass beads over the ends of the sutures before tying them, to prevent cutting into the margin of the lid. Both eyes are bandaged, and the sutures left in for a week or more.

Hugo Wolff's Operations for Congenital Ptosis by advancement of the levator palpebræ superioris:

Method I. (Fig. 69).—An incision of about 2 cm. in length is made through the skin of the upper lid in a position corresponding to the upper border of the tarsus, and the lips of the wound are each dissected up for a distance of 3 mm. By this means the orbicularis is laid bare. In the center of the wound, and at the upper margin of the tarsus, a fold of the orbicularis of about 1 cm. in width, with the subjacent levator tendon, is seized in the forceps. This fold is isolated in a vertical direction by a few strokes with the scissors, and is undermined. Two strabismus hooks are then passed under it, and placed so that one of them lies close to the insertion of the levator tendon in the tarsus.* The amount by which it is desired to raise the lid is measured off on the tendon from its insertion, and Schweigger's strabometer is convenient for this purpose. (If, for example, the palpebral opening in the normal eye is 10 mm., while that in the eye to be operated on is 3 mm., it will be required to raise the drooping eyelid 7 mm.) At the point found by this measurement two sutures, each with two needles, are applied in the muscle, and the latter is divided immediately below the ligatures. The four needles are then passed through the stump of the tendon at its insertion, and through the portion of orbicularis which covers it; the sutures are tied, and cut off short, and the skin wound is then closed by a few sutures.

Method II.—From the conjunctival surface. The eyelid is everted in the usual manner; the upper margin of the tarsus is then seized in the double-legged fixation forceps, and the lid is rolled over again, and by this means the field of operation is brought into view. To prevent bleeding a lid-clamp is now applied. On the right margin (as looked at by the operator) of the field of operation a fold of conjunctiva is raised and divided vertically, and the conjunctiva is undermined upwards, downwards, and to the left, and divided by a horizontal incision about 2 cm. in length. The conjunctival flaps are turned upwards and

* H. Wolff states that he finds the insertion of this tendon is 5 mm. below the upper margin of the tarsus.

downwards, and then the levator, with Müller's muscle lying on it, is exposed to view. With the forceps a portion in the center of the muscle, about 1 cm. wide, is seized and isolated from its bed by a few vertical strokes with the closed scissors. Two strabismus hooks are inserted under the isolated portion of muscle, one of them lying close to the convex margin of the tarsus, and then the clamp and fixation forceps are removed. In place of the strabismus hooks Wolff's spatula, with millimeter scale engraved on it, can be used. At the desired distance from the upper margin of the tarsus (which must precisely represent the difference between the palpebral opening in the sound eye and that in the faulty eye) two catgut sutures, with two needles each, are tied in the muscle, so that each knot will include more than half of the isolated portion of the muscle. The muscle is then divided close below the point of ligature; the needles are passed through the stump close to the upper margin of the tarsus, which is also included; the sutures are closed, and cut off short. The conjunctival wound is then also closed.

A remarkable condition is *Congenital Ptosis, with Associated Movements of the Affected Eyelid, during the action of certain muscles.* There are only about thirty cases of this on record. It is most commonly the left lid which is affected, and the paralysis may be congenital or acquired. Three conditions have been observed—viz., (1) elevation of the drooping lid when the eye is adducted, (2) when the eye is abducted, or (3) when the mouth is open. A synchronous contraction of the pupil has been noticed in some cases, while in some the elevation of the lid occurs also with a lateral motion of the jaw, and with deglutition. Gowers' explanation is that in these cases the levator is not wholly supplied by the third nerve, but partly also by nerve fibers, which take their origin in the nucleus of the fifth pair, and which also supply the external pterygoid and digastric muscles. But this theory does not hold good in all cases, for Bull [8] describes a case in which the lid was raised when the head was bent back, thus stretching the digastric. He regards these as associated or reflex movements. In some instances the lid can be raised voluntarily on closing the other eye. Needless to say, no remedy can be applied for relief of this condition.

The term ptosis is also given, although not very correctly, to cases in which increased weight of the lid causes it to droop, as in conjunctival affections, or where a tumor has formed in the eye-

lid, or where there is a hyperdevelopment of the subcutaneous fat.

Lagophthalmos (*λαγώς, a hare,* as it was supposed that this animal sleeps with its eyes open; *ὀφθαλμός*), or inability to close the eyelids, is most commonly due to paralysis of the portio dura, and is then associated with the other symptoms of the latter affection. On an effort to close the lids being made, the eyeball is rotated upwards under the upper lid, owing to the associated action of the superior rectus; and in sleep this upward rotation also occurs—a fact which explains, to a great extent, the immunity of the cornea from ulceration in many of these cases. Lagophthalmos may also be due to orbital tumors pushing the eyeball forwards, to exophthalmic goitre, to staphyloma, or to intra-ocular growths distending the walls of the eyeball—in all of which conditions the eyelids are often mechanically prevented from closing over the eyeball, or can be closed only by a strong effort of the will. The danger to the eye depends upon the tendency to ulceration of the cornea from its dryness, caused by exposure to the air, and from foreign substances not being removed from it by nictitation.

In cases of non-paralytic lagophthalmos protection of the cornea by keeping the eyelids closed with a bandage, or by inserting a few epidermic sutures in the margins of the eyelids, to draw them together, should be our first care. Tarsorrhaphy may be employed in those cases where circumstances indicate that it would be useful —*e. g.,* in some cases of exophthalmic goitre, or of staphylomatous eyeball.

In paralytic cases, the primary cause of the paralysis (syphilis, rheumatism, etc.) must be treated so long as there is a prospect of restoring power to the muscle. Locally, galvanism and hypodermic injections of strychnia may be employed. During cure the cornea should be protected as above. In incurable cases the opening of the eyelids must be reduced considerably in size by an extensive tarsorrhaphy, or by the method proposed by Pflüger.

The Operation of Tarsorrhaphy consists in uniting the margins of the upper and lower lids in the neighborhood of the external commissure, so as to reduce the size of the opening of the eyelids. The commissure should be caught between the finger and thumb, and the edges of the lids approximated, so as to enable the operator to form an estimate of the required extent of the operation. A horn spatula is then passed behind the commissure, and the

necessary length of the margin of each lid, including the bulbs of the cilia, abscised with a sharp knife. The raw margins are then brought together with sutures.

Pflüger's Method [9] consists in passing one, two, or even three double sutures subcutaneously around the eyelids, about 5 mm. from their margins. The ends are drawn together, so that the eye is concealed by the pouch thus formed, and tied. From time to time the sutures are tightened, until finally they cut through, and by this means a subcutaneous ring-cicatrix is produced. Should the first ring-cicatrix not sufficiently close the eyelids, the operation can be repeated even more than once again. The method is tedious and painful.

Symblepharon (*σύν, together; βλέφαρον, the eyelid*) is an adherence, partial or complete, of the eyelid to the eyeball. It is usually the result of burns of the conjunctiva by fire, acids, or




Fig. 70.




Fig. 71.*

lime. The shortening of the conjunctival sac, which is seen as the result of pemphigus or of granular ophthalmia, and which I have above described under the heading of Xerophthalmos, is not properly called symblepharon. If the symblepharon interfere seriously with the motions of the eyeball, or if it cause defect of vision by obscuring the cornea, it becomes desirable to relieve it by operation. Should it consist of a simple band stretching from lid to eyeball, it may be severed by ligature, and if the band be broad, two ligatures may be employed, one for either half. A symblepharon which occupies a considerable surface cannot be got rid of in this way; and for such cases a transplantation procedure like that of Teale [10] or of Knapp [11] may be employed, the great difficulty in dealing with these cases being the tendency there

* Mr. Teale now makes his flaps, as in Fig. 71, wider than he originally did. I have to thank him for altering this drawing with his own hand for this work.

is to reunion of the surfaces, unless one or both of them be carpeted with epithelium.

In *Teale's Operation,* if we suppose the case to be similar to that represented in Fig. 70, an incision is carried along the line of the margin of the cornea at *A,* through the whole thickness of the symblepharon, and the lid is dissected off from the eyeball as far as the fornix. Two conjunctival flaps are now formed, as at *B* and *C* in Fig. 71, and one of them (*B*) is turned to form a covering for the wounded surface of the inside of the eyelid, while the other (*C*) is used to cover the bulbar surface (Fig. 72), the flaps being held in their places by fine sutures. That part of the symblepharon which is left adherent to the cornea soon atrophies and disappears. No great tension of the flaps should exist as they lie in their new positions.

Teale, again, has suggested the formation of a bridge-like conjunctival flap above the cornea, and the removing of it across the




Fig. 72.




Fig. 73.

latter to cover the loss of substance situated below. After the sutures to keep the flap in its place have been introduced, the latter is separated at its bases.

A simple plan, which would be applicable to such a case as that depicted in Fig. 70, where the adhesion is not very extensive, and perhaps even to some more extensive ones, consists in dissecting the conjunctival process off the cornea, and then turning it down on the raw inner surface of the under lid, and fastening it there with a suture or two. I have done this with complete satisfaction.

Harlan's Operation.[12]—This is especially applicable to extensive symblepharon of the lower lid, and differs from the foregoing operations in that it provides a covering of skin, and not of mucous membrane, for the raw surface of the under lid. Operations on the same principle have been proposed by Snellen and by Kuhnt.

An incision (A B, Fig. 73) through the whole thickness of the eyelid, and corresponding in length to the latter, is made along the lower margin of the orbit. Below this a skin flap (C D) is then formed. The flap is dissected up, and the incisions are carried a little more deeply as A B is approached, to enable the flap to turn the more readily. The flap is then turned up as on a hinge, slipped through the buttonhole, and sutured securely to the inner surface of the under lid. After a time the skin surface turned towards the eyeball becomes considerably modified, so as to be somewhat like mucous membrane. The bare space left by the removal of the strip of skin is covered without strain by making a small horizontal incision (D E) at its outer extremity, and forming a sliding flap.

Transplantation Operations.—The transplantation of a portion of mucous membrane from the lips or cheek, as employed by Stellwag, is a useful method for many cases of extensive symblepharon. The chief precautions necessary for success in this proceeding are: That the flap to be transplanted be not applied in its new position until all bleeding at the latter place has ceased. That the flap be nothing more than mucous membrane, all submucous tissue being carefully removed. That it be sufficiently large to cover the defect without any stretching; and it should be remembered that the flap shrinks to two-thirds of its size after being detached from its own bed. That the flap be kept moist and warm during the period—as short as possible—which may elapse between its detachment and its adjustment. And, finally, that it be kept firmly in its new position by a sufficient number of points of interrupted suture. It is seldom possible in extensive cases of symblepharon to do all that is necessary at one sitting.

A drawback to mucous membrane flaps, where two opposing surfaces have to be covered, consists in the liability there is, when the superficial epithelium of the mucous membrane is thrown off, for the surfaces to unite. Hence some operators prefer to transplant Thiersch grafts, which are not open to this objection; or, mucous membrane may be used for one surface and Thiersch grafts for the opposite surface. In order to keep the grafts in position it is desirable, when possible, to insert an artificial eye, or some such support, into the socket.

Blepharophimosis (*βλέφαρον, eyelid; φίμωσις, narrowing*) is a contraction of the outer commissure of the lids, with consequent diminution in size of the opening between the latter; and is com-

monly due to shortening of the skin, from long-continued irritation of it, caused by the discharge in a case of very chronic conjunctivitis.

It is remedied by a *Canthoplastic Operation.* The outer commissure is divided in its entire thickness, in a line which is a prolongation of the line of junction of the lids when closed, by a single stroke of a strong straight scissors, one blade of which has been passed behind the commissure. The integumental incision should be made a little longer than that in the conjunctiva. An assistant then draws the upper lid up and the lower lid down, so as to make the wound gape. The conjunctival margin and the dermic mar-




FIG. 74—(*de Wecker*).

gin are now united in the center by a point of suture (C, Fig. 74), while two more sutures (A and B) are applied, one above and the other below the first. This operation is also employed in cases of granular ophthalmia and of purulent conjunctivitis when it is desired to relieve the pressure of the lid on the globe.

Distichiasis (*δίς, twice; στίχος, a row*) and **Trichiasis** (*τρίχος, a hair*).—The first of these terms indicates the growth of a row of eyelashes along the intermarginal portion of the lid in addition to the normal row; while trichiasis indicates a false direction given to the true cilia. Both conditions are often found coexisting, and often, too, they are present along with entropium. They may both be produced by chronic blepharitis or by chronic granular oph-

thalmia. It has been commonly held that cicatricial contraction, giving a false direction to the hair follicles, is the immediate cause of these conditions; but Raehlmann has recently [13] shown that the false cilia are developed as buds or offshoots from the follicles of the cilia, and primarily from the cuticle of the free margin of the lid. The latter mode of development is a novel discovery by Raehlmann, which he seems to have definitely proved by his pathological investigations. His view is that hyperemia of the margins of the lids, and inflammation of a proliferating type are what give rise to this primary development of hairs. The symptoms they produce, and the dangers to the eye attendant on them, are due to the rubbing of the irregular eyelashes on the cornea, which pro-




Fig. 75.

duces pain, blepharospasm, and opacity of the cornea, or even ulceration of it.

Operations for Distichiasis and Trichiasis:

Epilation.—The false cilia may be pulled out with a forceps; but this cannot be regarded as a cure, for the hairs grow again.

Electrolysis has been proposed by Charles Mitchell, of Missouri,[14] and by Arthur Benson, of Dublin.[15] A needle is attached to the negative pole, and its point passed into the bulb of the eyelash to be removed, the positive pole being placed on the temple. On closure of the circle, if the battery be working properly, bubbles of gas should rise up round the needle, and a slough forms at the root of the hair, which becomes loose, and is removed. It does not grow again, for the bulb is destroyed. Each hair must be separately operated on. The proceeding is very valuable where only a few cilia are to be dealt with.

Illaquetio.—Snellen has revived this ancient operation for cases where only a few isolated hairs are out of order. Both ends of a

bit of very fine silk thread are passed through the eye of a fine needle, so as to form a loop. The needle is now entered as close to the point of exit of the hair as possible, and the counter-puncture is made in the position which the hair should normally occupy in the row of its fellows. The needle is drawn completely through, as also the ends of the thread, but the loop not as yet. Into the loop the eyelash is now inserted by aid of a fine forceps, and by traction on the ends of the thread, loop and eyelash are drawn through the tunnel. Unfortunately the eyelashes fre-




Fig. 76.

quently regain their abnormal position by reason of their own elasticity.

Excision.—When some half-dozen hairs close together are growing wrong, the simplest and best plan is to completely remove them by excision of the corresponding portion of the ciliary margin. A fine knife is passed into the intermarginal region at the place corresponding to the hairs to be dealt with, and a partial division of the lid into two layers, as in the Arlt-Jaesche operation (*vide infra*), is effected. A V-shaped incision in the skin of the lid is then made, including the erring hairs, the whole flap is excised, and the margin of the loss of substance drawn together with sutures.

In cases of distichiasis or trichiasis involving the whole length of the eyelid, removal of the marginal portion of skin containing the bulbs of all the eyelashes, true and false (Flarer's operation), is not to be recommended—unless, occasionally, in the under lid —because it unnecessarily deprives the eye of an ornament, and of a protection against glare of sun and foreign bodies.

Transplantation, or *Shifting,* of the marginal portion of the integument containing the hair bulbs, true and false, is a preferable proceeding in these complete cases. One of the oldest and most valuable operations of this kind is that of Jaesche, modified by Arlt. It is performed as follows: Knapp's, or Snellen's, clamp (Fig. 75) having been applied to prevent bleeding, the lid in its whole length is divided in the intermarginal part into two layers




Fig. 77.




Fig. 78.

(Fig. 76), the anterior containing the orbicular muscle and integument with all the hair bulbs, the posterior containing the tarsus and conjunctiva. The incision in the intermarginal portion is about 5 mm. deep. A second incision is now made through the integument of the lid, parallel to its margin, and from 5 to 7 mm. removed from it. This incision also extends the whole length of the lid. A third incision is carried in a curve from one end to the other of the second incision. The height of the curve is proportional to the effect required, varying from 4 mm. to 7 mm. The piece of integument included between the second and third incisions is dissected off with forceps and scissors, without any of the underlying muscle being touched, and the margins of the loss of substance are brought together by sutures. By this procedure

the lower portion of integument, containing the hairs and their bulbs, is drawn up and away from contact with the cornea.

Many double transplantation operations have been proposed; of these:

Dianoux's Operation[16] is as follows: Snellen's (or de Wecker's) clamp is applied (omitted in figures for simplicity), and an incision (Fig. 77) is made parallel to the free margin of the lid, about 4 mm. from it, extending the whole length of the lid, and penetrating to the tarsus, but not through the latter. The ciliary portion of the lid marked off by this means is now detached from the tarsus by an incision in the intermarginal portion of the lid, as in the Arlt-Jaesche operation. An incision through the skin alone is then made about 3 mm. above the first incision and parallel to it,



Fig. 79.

but extending some 2 mm. beyond it at either extremity. The skin flap is separated off from the underlying muscle, except at either end, where it is left attached. The underlying portion of the muscle is then separated from the tarsus, and allowed to retract upwards. A forceps is passed under the ciliary flap (Fig. 77), and the skin flap is seized and drawn down into the position of the former (Fig. 78), where it is made fast by three sutures to the margin of the tarsus. The ciliary flap is moved up, and carefully stretched upon the tarsus bared of the orbicularis, the latter being drawn back with a strabismus hook, and the flap is secured in its place by sutures to the tarsus. An antiseptic dressing is applied, and the sutures may be removed on the third day. Although the wounded surface of the ciliary flap does not become

vitally united with the epidermic surface of the skin flap, yet no practical ill result follows.

A real objection lies in the circumstance that occasionally the cutaneous hairs on the transplanted flap irritate the cornea, and these hairs, being much finer than cilia, are more difficult to deal with.

Vossius' Operation.[17]—If, for example (Fig. 79), the whole extent of the right upper lid be affected with trichiasis, a horn lid-spatula (the clamp will not answer) is passed under the lid, and held by an assistant. An intermarginal incision is made, as in the Arlt-Jaesche operation, about 3 mm. to 4 mm. deep. This incision is then prolonged through the skin merely, over the external com-




Fig. 80.

missure for 5 mm. to 6 mm. It is then turned upwards at an angle of about 35° with the free margin of the lid, and a flap about 5 mm. wide is marked out with the knife in the usual crease or fold of the upper lid. A narrow, sharp, and pointed scalpel is then thrust under the flap at its base, and carried towards its inner end, so as to separate it off without the aid of forceps, scissors, or any other instrument. The margins of the wound thus made are brought together with four or five sutures, and the flap turned down and secured in the gaping intermarginal incision by means of four or five sutures between each of its edges and the corresponding palpebral margin. One suture fastens the free end of the flap in the median corner of the wound. The position of the cicatrix, just in the fold of the upper eyelid, prevents

its causing any disfigurement. Were the case one of partial trichiasis, the intermarginal incision should extend a little beyond the point where the abnormal condition ceases. If it be the inner half only of the margin of the lid which is affected, the intermarginal incision is prolonged towards the nose, and the flap so formed that its base lies over the inner canthus. The flap heals in readily, and, although it shrinks somewhat, secures a wide intermarginal portion. The same drawback in connection with the cutaneous hairs on the transplanted flap holds good here as in Dianoux's operation.

Van Millingen's Operation[18] consists in splitting the eyelid, as in the Arlt-Jaesche operation, from end to end, sufficiently to produce a gap (*B*, Fig. 80) 3 mm. in width at the central part of the lid, and gradually becoming narrower towards the canthi. The gap is kept open by sutures passed through folds of skin on the upper lid (*a a a*), by means of which also the lid is prevented from closing for twenty-four hours at the least. As soon as the bleeding has ceased, a strip of mucous membrane of the same length as that of the lid, and 2 to 2 1-2 mm. in breadth, is cut out with two or three snips of a curved scissors from the inner surface of the patient's under lip, and is placed at once into the gap in the intermarginal space. It should then be pressed into position with a pledget of cotton-wool steeped in sublimate solution (1 in 5000). According to Van Millingen, sutures are superfluous; but I like them for the sake of security, and I do not find that they do harm. The eyelid is then covered over with a piece of lint, on which is spread a thick layer of xeroform vaselin, and on this is placed a wad of cotton-wool and a bandage. Both eyes should be bandaged. The sublimate lotion is used for disinfecting the eye and lip during, before, and after the operation. The dressing should be renewed once in twenty-four hours, and the sutures in the upper lid ought not to be removed before the second day.

Van Millingen does not think it advisable to transplant small strips of mucous membrane if the trichiasis be partial. He regards this condition as only the commencement of complete trichiasis, and therefore recommends, even in these cases, the filling up of the entire length of the intermarginal space with a flap of mucous membrane. In cases of shortening of the conjunctival surface, in which it has been reduced to 1-2 cm., a strip of mucous membrane measuring 4 mm. in width at the center may be transplanted.

The strip to be transplanted is generally taken from the angle of the lip and from the line of demarcation between the dry and moist surfaces of the lip. A couple of fine sutures, which serve to unite the margins of the wound in the lip, arrest the bleeding at once, and accelerate union of the part, which is generally completed in twenty-four hours.

The transplanted tissue in this instance being free from hairs, the method is not open to the objection referred to in Dianoux's and Vossius' operations, while it is equally effectual in permanently providing a good intermarginal space and in thus permanently relieving the condition.

Entropium (*ἐν, in; τρέπω, to turn*), or **Inversion of the Eyelid,** is due to some organic change in the conjunctiva or tarsus, or to spasm of the palpebral portion of the orbicular muscle.

A large proportion of the former class of cases is the result of chronic granular ophthalmia, and is most common in the upper lid.

Spastic entropium occurs in the under lid only. It is frequent in old people (senile entropium) from relaxation of the skin of the eyelid, and is also produced by the wearing of a bandage after operations, etc., and by edema of the conjunctiva in inflammation of that membrane.

Treatment.—Organic entropium, in which tarsus of the upper lid is not distorted, can often be corrected by one of the methods described for trichiasis and distichiasis. But many of these cases are accompanied by, or rather are due to, abnormal curvature with hypertrophy of the tarsus.

In all such cases the operation must include an attack on the tarsus itself, or the result will be abortive. Indeed, I have little doubt that much of the disappointment experienced in the treatment of entropium has been due to imperfect appreciation of this fact.

Streatfield's Operation is as follows: The clamp having been applied, an incision is made through the integument of the eyelid parallel to its margin, 2 mm. distant from the latter, and extending its whole length. The muscle is dissected up so as to lay bare the tarsus, and then a wedge-shaped piece, 2 mm. wide and the length of the lid, the edge of the wedge pointing towards the inner surface of the lid, is excised from the tarsus. A corresponding portion of muscle and skin is also removed, and the wound is allowed to heal by granulation. The shrinking of the resulting

cicatrix causes the marginal portion of the tarsus to return to its correct position.

Snellen's Operation.—Snellen's clamp (very similar to Knapp's, which can equally well be used) is applied. About 3 mm. from the margin of the lid, and parallel to it, an incision is made through the skin alone, extending the whole length of the lid. The orbicular muscle is exposed by dissection of the skin upwards, in order to promote retraction of the latter, and along the edge of the lower margin of the wound a strip about 2 mm. broad of the orbicular muscle is removed, and the tarsus to the same extent exposed to view. A wedge-shaped piece corresponding to the exposed part of the tarsus is now excised from it with a very sharp scalpel or Beer's cataract knife, the edge of the wedge pointing towards



Fig. 81.




Fig. 82.

the conjunctiva, which latter, however, is left intact. The hypertrophy of the tarsus, which is always present, facilitates this procedure. A silk suture carrying a needle on each end having been prepared, one needle is passed from within outwards through the band of muscle and integument left at the margin of the lid. The second needle is also passed from within outwards through the upper lip of the tarsal loss of substance, and then from within outwards through this same marginal band, at a distance of about 4 mm. from the point of exit of the first needle. The ends of the structure are now tied together, a small bead having first been strung on each to prevent it from cutting through the skin. Three such sutures are employed. The accompanying figures (81 and 82) make the foregoing description more intelligible.

Green's Operation.[19]—An incision is made on the inner sur-

face of the lid in a line parallel to, and about 2 mm. distant from, the row of openings of the Meibomian ducts. It is carried through the conjunctiva and whole thickness of the tarsus, and should extend, in cases of complete entropium, from near the inner to the outer canthus. A strip of skin about 2 mm. broad, and tapering to a point at each end, is now excised from the lid,—the lower margin of the strip being 1 1-2 mm. above the line of the eyelashes. The muscle is left intact. Fine silk sutures are applied in the following manner by aid of a No. 12 glover's needle bent to an arc of about a third of a circle: The needle is first introduced a little to the conjunctival side of the row of eyelashes, and is brought out just within the wound made by the excision of the strip of skin (*A*, Fig. 83); it is then drawn through, inserted again in the wound near its upper margin, and




Fig. 83.

passed deeply backwards and upwards so as to graze the front of the tarsus, and emerge through the skin a centimeter or more above its point of entrance (*B*, Fig. 83). On tying the two ends of the thread together, the skin-wound is closed, and the loosened lid-margin is at the same time everted and brought into a correct position. Three sutures generally suffice for the accurate adjustment of the lid-margin. In the spaces between and beyond the sutures it is often practicable and advantageous to turn the eyelashes upwards against the front of the eyelid, and fix them there by means of collodion. The stitches should be removed at latest on the day after the operation, the line of suture being then strengthened by collodion, or, in case the cilia are very short, a few short fibers of cotton are used with the collodion.

Berlin's Operation.—Knapp's clamp is applied. The first incision lies 3 mm. above the margin of the lid, extends its whole length, and divides it in its entire thickness, including the conjunctiva. The skin and muscle at the upper edge of the wound

are pushed or dissected up so as to expose the tarsus. The upper edge of the tarsal incision is now seized at its center with a finely toothed forceps, and an oval piece with the adherent conjunctiva, about 2 to 3 mm. wide in its widest part, and in length corresponding with that of the eyelid, is excised from it with a fine scapel. The wound is closed with three sutures through the skin. If it be thought desirable to increase the effect, a skin-flap may be excised from the lid. The objection to this operation, that a portion of the mucous membrane is removed, is not of importance. Except for an occasional granulation forming on the bulbar aspect of the wound, I have found the operation free from inconvenience, and its result satisfactory, and in most instances permanent.

Spastic Entropium, as the result of bandaging, usually disap-




FIG. 84.

pears when the use of the bandage is given up, or, if the bandage must be continued and should the inverted lid cause irritation, an epidermic suture at the palpebral margin and fastened to the cheek below will give relief.

Senile Entropium of the lower lid is, of the spastic kinds, the one which most commonly demands operative interference. The methods in general use for it are:

The Excision of a Horizontal Piece of Skin, with a portion of the underlying orbital part of the orbicular muscle, so as to give rise to sufficient cicatricial contraction to draw the margin of the lid outwards.

The application of *Subcutaneous Sutures (Gaillard's Sutures).* —The point of a curved needle carrying a silk suture is entered in the center of the lid near its margin, passed deeply into the

orbicular muscle, brought out at a point some 10 mm. below, and the suture tied tightly. Two more similar sutures, one on either side of the first and about 5 mm. distant from it, are placed. The sutures are allowed to remain for a week or more, and the resulting cicatrization brings the lid into its position.

Von Graefe's Operation.—3 mm. from the margin of the lid an incision is made, as in Fig. 84, through the skin, and a triangular skin flap (*A*) excised. The edges *B* and *C* of the triangle are dissected up a little and brought together by three points of suture, while the horizontal incision is not sutured. The size, especially the width, of the triangular flap to be excised is proportional to the looseness of the skin. When a very marked effect is desired, the flap to be removed is given the shape as represented at the right




Fig. 85.

of the figure. I have found this proceeding extremely satisfactory, and its result, as a rule, permanent.

All the foregoing and other such measures produce a good result at the time, but are sometimes followed by recurrence of the entropium. Holtz believes the cause of this to be that the cicatrix, be it dermic or dermo-muscular, upon which the result depends, has no *point d'appui;* and consequently, while it may draw the eyelid out, it is just as liable to draw the skin of the cheek up, and thus neutralize its desired effect. He proposes the following ingenious operation:

Holtz's Operation.[20]—A horn spatula is inserted under the lid, and then, at 4 to 6 mm. below the margin of the latter, a horizontal incision is made through the skin from the inner to the outer

end of the lid. This incision is at the boundary between the palpebral and orbital portions of the orbicular muscle, and just over the lower margin of the tarsus. An assistant then draws the upper edge (*a*, Fig. 85) of the wound upwards with a forceps, while the surgeon draws the lower edge (*b*) downwards, in this way exposing and stretching the orbicular muscle. A few strokes of the knife in the direction of the incision are now sufficient to separate the palpebral portion (*l*) of the muscle from the orbital portion (*p*), and to lay bare the lower edge of the tarsus (*t*), which has a yellowish tendinous appearance. That part of the palpebral portion of the muscle which covered the lower edge of the tarsus, and which was drawn up with the palpebral edge of the first incision, is now removed with forceps and scissors, to the extent of about 2 mm. in width, through the whole length of the lid. All such muscular fibers, also, which may still adhere to the lower third of the tarsus must be carefully cleaned off, and now the palpebral skin may be brought into union with the tarsus. Four sutures are generally applied, about 5 mm. apart. The needle is passed through the palpebral skin, close to the margin of the wound (at *a*). The bare tarsal edge is then seized in the forceps, the needle placed perpendicularly on it (at *d*), and carried through it by a short downward curve until its point appears (at *c*) below the tarsus in the tarso-orbital fascia (*f*). The needle is now passed out through the lower edge of the incision (at *b*), care being taken that none of the fibers of the orbital portion of the muscle are included in the suture. Upon the suture being tightly closed, the edges of the skin wound are drawn into the tarsus, and become adherent to it. The sutures may be removed about the third day. If the first incision be placed too far from the margin of the lid, there will be no result, as the traction upon the palpebral skin will be too slight. If the incision be placed too close to the margin, the traction may be so great as to interfere with the union of the skin and tarsus. In this operation the tarsus affords the fulcrum, which Holtz thinks is wanting in other methods. The tarsus of the lower lid is sometimes very little developed, and then I find the result of the operation may be disappointing.

Ectropium or **Eversion of the Eyelid.**—Of this there are two chief kinds: (1) Muscular, or Spastic, which affects the lower lid only; (2) Cicatricial.

Muscular ectropium may be caused by edema of the conjunctiva, which everts the edge of the lower eyelid, and this eversion is

increased and encouraged by spasm of the palpebral portion of the orbicular muscle, so that the term palpebral paraphimosis might be given to the condition. In the recent stage it may generally be remedied by a properly applied bandage, combined with the suitable conjunctival measures. In chronic cases operative measures may be required.

Muscular ectropium is often seen in old people, and is then given the name of Senile Ectropium. Here it is due to atrophy of the palpebral portion of the orbicularis of the lower lid and relaxation of the skin of the face. When these have resulted in slight eversion of the inferior punctum, a flowing of tears is produced, causing excoriations of the skin and edge of the lid, which then, in their turn, increase the tendency to ectropium. If the condition be not extreme, with secondary changes in the conjunctiva, slitting up of the canaliculus, with the use of a boric ointment for the lids and mild astringents for the conjunctiva, will give much relief. In pronounced cases a more active treatment of the conjunctiva, and the performance of tarsorrhaphy, or the latter preceded by the application of Snellen's sutures, or Kuhnt's operation, are demanded. Muscular ectropium is also caused by paralysis of the orbicular muscle.

Snellen's Sutures.—A silk ligature is threaded at either end with a needle of moderate size and curve. The point of one of these needles is passed into the most prominent point of the exposed and everted conjunctiva, and brought out through the skin 2 cm. below the edge of the lower lid. The other needle is entered in the same way 5 mm. from the first, and made to take a nearly parallel course, the points of exit on the cheek being 1 cm. apart. Equal traction is applied to each end of the suture, while the lid is assisted into its place by the finger. The suture is tied on the cheek, a small roll of sticking-plaster having been inserted under it, to protect the skin from being cut. Two, or even three, such sutures may be required, and they are allowed to remain for several days.

Kuhnt's Operation[21] for Senile Ectropion is an admirable one. It consists in splitting the lower lid in its central third, so that the conjunctiva and tarsus are left in the posterior layer, while the anterior layer contains the orbicularis and the skin. A triangular piece, the base of which is formed by the margin of the lid, is then excised from the posterior layer, and the margins of the loss of substance in the latter are brought together by three or

four points of suture. Lest they should give way too soon, it is necessary to place these sutures very securely. A puckering of the anterior layer, opposite the line of sutures in the posterior layer, is produced, but subsequently disappears, and a suture which unites the most prominent point of the pucker and the margin of the tarsus assists in this. Or, if the lid be split, say, to an extent twice as long as the base of the triangular piece to be excised, the puckering can be distributed at either end of the incision. By reason of the shortening of the lid as the result of this excision the




FIG. 86.

eversion is corrected. I have repeatedly used this operation, and always with gratifying results.

Argyll Robertson's Operation [22] has been designed for those cases of ectropium which result from long-continued chronic inflammation of the conjunctiva of the lower lid. He thinks the difficulty in severe cases of this kind depends upon the abnormal curvature which is gradually acquired by the tarsus. The following is his description of the operation, from which he has obtained satisfactory results:

The materials required are:

1. A piece of thin sheet-lead about 1 inch long and 1-4 inch broad, rounded at its extremities, and with its cut margins

smoothed. This piece of lead must be bent with the fingers to a curvature corresponding to that of the eyeball.

2. A waxed silk ligature about 15 inches long, to either extremity of which a long, moderately curved needle is attached.

3. A piece of fine india-rubber tubing of the thickness of a fine drainage-tube.

The operation is performed by perforating the whole thickness of the lid with one of the needles at a point (*b*, Fig. 86) one line from its ciliary margin, and a quarter of an inch to the outer side of the center of the lid. The needle, having been drawn through (at *a*), is passed directly downwards over the conjunctival surface of the lid till it meets the lower conjunctival fornix, through which the needle is thrust—the point being directed slightly forwards—and pushed steadily downwards under the skin of the cheek, until a point (*d*) is reached about 1 inch or 1 1-4 inch below the edge of the lid, when the needle is caused to emerge, and the ligature is drawn through. The other needle is, in like manner, thrust through the edge of the lid at a corresponding point (*b'*) a quarter of an inch to the inner side of the middle of the lid, then passed over the conjunctival surface of the lid, through the fornix, and downwards under the skin, till the point emerges at a spot (*d'*) a quarter of an inch outwards from the point of exit of the first needle (*d*). The ligature is kept slack, or is slackened so as to permit of the piece of lead being introduced under the loops of the ligature that pass over the conjunctival surface of the lid, and of the piece of india-rubber tubing (*c*) being slipped under the loop at the edge of the lid (between *b* and *b'*). The free ends of the ligature are now drawn tight, and tied moderately tightly over a lower part of the india-rubber tube; the excess of india-rubber tube is cut off—about three-quarters of an inch beyond the ligature—and the operation is complete.

The result of the procedure is that the edge of the lid is made to revolve inwards over the upper edge of the piece of lead, while the tarsus is caused to mold itself to the curve of the lead, and the eyelid at once occupies its normal position. A certain amount of redness and edema of the lid follows the operation, and suppuration occurs in the track of the ligature; but as the india-rubber tube yields somewhat to the tension on the ligature, the resulting irritation is moderate, so that the apparatus need not be removed for five, six, or seven days, by which time the tarsus has become pretty well fixed in its new curvature. A slight relapse

may occur when the apparatus is removed, but this is readily amenable to treatment by astringent applications.

The suppuration occurring in the tracks of the ligature leads to cicatricial formation, which appears to impart a degree of rigidity to the lid that helps to keep the latter in its new position.

Kenneth Scott's Operation.[23]—The external canthus and tissues beyond are thoroughly divided by a pair of strong scissors; the lower eyelid, which is usually the affected one, is then seized and its margin stretched sufficiently outwards, parallel to the border of the other lid, so as to restore the palpebral aperture to its proper appearance; the portion of eyelid margin thus made to extend beyond the site of the external canthus is removed, along with its contained eyelashes, by slicing it with a sharp knife.



Fig. 87.




Fig. 88.

The upper and lower eyelids are then brought together, so that the original outer extremity of the one approximates exactly to the new extremity of the other eyelid. They are secured in this position by passing a silver wire suture vertically downwards through the substance of the upper lid, continuing it out through that of the lower one, and then twisting the ends firmly together. Two of these retaining stitches may be introduced close together if necessary. The edges of divided skin, along with the deeper muscular tissues, including that part which recently formed the outer end of the affected eyelid, are simply stitched together with a continuous fine silk suture.

No dressing other than a repeated dusting with some fine

antiseptic powder need be used. The silk stitches may be removed in six days' time, the silver ones being left in for five or six days longer. Dr. Scott states there is never any puckering apparent beyond the newly formed canthus, and the small linear cicatrix is lost amongst the other lines often found there.

Cicatricial Ectropium is caused by scars from wounds or burns, or from caries of the orbit, and can only be relieved by operation.

Wharton Jones's Operation is as follows: The cicatrix is circumscribed by a V-shaped incision (Fig. 87), and the skin made thoroughly movable in its neighborhood. The edges of the wound are now brought together so as to form a Y (Fig. 88).

Arlt's Operation for cases due to caries of the margin of the orbit.—If the cicatrix be situated at *e* (Fig. 89), the incisions at *a b* and *b c* are made through the skin and muscle, so that an acute, or at most a right, angle is formed at *b*. The margin




Fig. 89.

of the lid from *c* to *d* is excised. The cicatrix is completely undermined, and the triangle dissected up from *b* to the margin of the tarsus, so that the lid can be readily put into its position, and the edge *c b* of the flap united to *d c*. The size of the exposed surface on the cheek can, according to Arlt, be diminished by drawing its edges together after the manner of a hare-lip, but possibly the transplantation of a piece of skin from the arm to fill the gap might be a better plan.

The foregoing and similar operations are difficult or impossible in many cases where there has been great destruction of the skin of the eyelids and surrounding parts by burns, ulcers, etc., and at best the deformity is liable to recur. Transplantation of skin

from different parts of the body is in these cases a more promising proceeding. A description of the method is given in the next paragraph but one.

Ankyloblepharon (*ἀγκύλη, a string; βλέφαρον, an eyelid*) is a uniting of the upper and lower eyelids along their margins. It may be partial or complete, and often goes with symblepharon. Like the latter, it is usually caused by burns and ulcers.

The condition can only be relieved by operation, of which the result is often unsatisfactory, owing to the difficulty of preventing reunion taking place. To avert this it is always necessary to cover the wounded surface with conjunctiva or skin.

The Restoration of an Eyelid.—It is an extremely rare event for the whole substance of one or both eyelids to be destroyed by lupus, or other ulceration, or by accidents, which do not at the same time injure the eyeball seriously. In this rare event the eyeball, especially if the upper lid be destroyed, is exposed; the patient is subject to extreme discomfort; and, owing to ulceration of the cornea, the eye is ultimately lost.

The formation of an eyelid from the skin of the forehead or cheek in these cases is a most disappointing proceeding, and one the description of which does not, I consider, come within the scope of this book. Indeed, my own feeling in such a case would be to recommend enucleation of the eyeball, provided the fellow eye were good, rather than to propose a plastic operation which at the best would give but an imperfect result.

Fortunately the class of cases with which we commonly meet are essentially different in their nature; for in them the whole thickness of the eyelid is not destroyed. They are usually the result of burns (epileptics and children falling in the fire) and scalds which only destroy the integument of one or both eyelids. A granulating surface replaces the skin, and when healing commences the shrinking process draws the free margin of the upper eyelid up towards the eyebrow, and that of the lower lid down towards the cheek, the conjunctival surface of the eyelids becoming everted, and the cornea exposed, as the eyelids cannot now be closed. We have a satisfactory method for dealing with these cases.

In the first place the eyelid—let us suppose it to be the upper eyelid—is dissected down into its place to the utmost limit, so that the most extensive raw surface possible may be obtained. The margin of the lid is now fastened to the cheek with three

points of suture. A portion of skin, suited as regards shape, and one-third larger (to allow for shrinkage) than the raw surface of the eyelid, is then taken from the inside of the arm, and after being carefully freed of all its subcutaneous fat and connective tissue, is laid upon the raw surface and fastened to it by a large number of fine sutures around the margin. Or, if the margin of the skin surrounding the raw surface be dissected up, the edge of the graft can be slipped under it, and secured in its place by this means. A non-irritating antiseptic dressing is applied, and the graft usually heals on in the course of a few days. This method of grafting was introduced by Wolfe and Lefort, and I have employed it many times with most satisfactory results.

It is most important to preserve and utilize any part of the eyelid which remains, especially its ciliary border with the eyelashes.

The flap sometimes becomes separated from the wounded surface by oozing of blood or serum from the wound, and then sloughs, and it is important not to apply the graft until all hemorrhagic oozing has ceased. To obviate any such separation, Wickerkiewicz has employed secondary transplantation with satisfactory results. The flap is applied to the wounded surface from two to five days after the latter has been prepared, while during the interval the wounded surface has been protected with moist antiseptic dressings. He states that union by first intention occurs readily by this method.

The transplantation of a flap with pedicle from the forehead, temple, or cheek is also used to repair an eyelid; but, owing to the thickness of the integument, the result is cosmetically less satisfactory than that given by a graft from the arm, while the tendency to shrink is quite as great.

Sachs prefers Thiersch grafts to dermic grafts. He states they are more easily applied to the raw surface, and do not differ in color from the surrounding skin when healing is completed. He is careful to obtain one continuous graft of the whole size of the wounded surface.

Injuries of the Eyelids.—All kinds of injuries of the eyelids (contusions, incisions, burns, etc.) are common.

In consequence of the looseness of the integument, edema and ecchymosis, one or both, are often seen in a marked degree as the result even of slight injuries.

Owing to the direction of the fibers of the orbicularis, an in-

cised wound of the eyelid, if in the vertical direction, will gape, while a similar wound in the horizontal direction will not do so. Hence the scar left after the former wound is apt to be very visible, but that after the latter may be almost imperceptible. If the eyelid be divided vertically in its entire thickness, unless union by first intention can be obtained, a deep furrow is left in the eyelid, and, perhaps, at its margin an unsightly coloboma.

Emphysema of the eyelids is sometimes seen after a blow on the eye, and is a sign of fracture of the orbit, with a communication between the subcutaneous connective tissue of the eyelids and the nose, the ethmoid sinus, the frontal sinus, or the antrum of Highmore. An emphysematous lid is swollen, soft, and crepitating to the touch.

Ecchymosis of the lower lid, usually with ecchymosis of the lower conjunctiva, after falls or blows on the head, is a sign of fracture of the base of the skull, the blood making its way along the floor of the orbit.

Simple ecchymosis of the eyelids from blows, commonly known as "Black Eye," never gives rise to further complication. It requires some fourteen days or more, according to the quantity of blood extravasated, before the eye recovers its normal appearance.

Treatment.—Injuries of the eyelids, of whatever kind, are of course treated upon general surgical principles. Incised wounds should be carefully and neatly drawn together with sutures as soon after the injury as possible, and with antiseptic precautions. Emphysema may be assisted in its absorption by the application of a rather tight bandage, and directions should be given to the patient to blow his nose as gently as possible, so as to avoid recurrence of the emphysema.

Epicanthus is a congenital deformity, usually binocular, which, in the most pronounced cases, consists in partial paralysis of the levator palpebræ (ptosis) and of the rectus superior, with a narrow palpebral fissure, and a fold of integument at the inner canthus concealing the caruncle from view, and giving the appearance of great breadth to the bridge of the nose. The term is also used for cases in which the integumental fold at the inner canthus is the only abnormal condition, and this deformity can be somewhat diminished by the removal of an oval piece of skin from the bridge of the nose, its long axis being vertical and its width varying according to the effect required. When the margins of

the wound are brought together the abnormal folds are diminished in width.

Congenital Coloboma of the upper lid, sometimes associated with a dermoid cyst of the limbus of the cornea corresponding to the cleft in the lid, and even congenital absence of the eyelids, have been occasionally observed.

References.

[1] "Brain," 1900, p. 353.
[2] "Practitioner," November, 1898.
[3] "Centralblatt für praktische Augenheilkunde," 1892, p. 129.
[4] "Archives d'Ophtalmologie," Janvier-Février, 1886.
[5] "Von Graefe's Archiv, xxxvi. 1. p. 234.
[6] "Klinische Monatsblätter für Augenheilkunde," 1883, p. 100.
[7] "Archiv für Augenheilkunde," xxxiii. p. 125.
[8] "Archives of Ophthalmology," xxi. p. 354.
[9] "Klinische Monatsblätter für Augenheilkunde," 1902, p. 527.
[10] "Roy. Lond. Ophthalmic Hospital Reports," xiv. 1. p. 270.
[11] "Von Graefe's Archiv," xiv. 1. p. 270.
[12] "Ophthalmic Review," ix. p. 351.
[13] "Von Graefe's Archiv," xxxvii. 2. p. 66.
[14] "Trichiasis and Distichiasis, their Nature and Pathology, with a Radical Method of Treatment," 1882.
[15] "British Medical Journal," December, 1882.
[16] "Annales d'Oculistique," 1882, p. 132.
[17] "Bericht der ophthalmologischen Gesellschaft." Heidelberg, 1887, p. 42.
[18] "Ophthalmic Review," 1887, p. 309.
[19] "Trans. American Ophthalmological Society," viii. p. 167.
[20] "Klinische Monatsblätter für Augenheilkunde," 1880, p. 149.
[21] "Beiträge zur operativen Augenheilkunde." Jena, 1883.
[22] "Edinburgh Clinical and Pathological Journal," December, 1883; and "Ophthalmic Review," iii. p. 47.
[23] "British Medical Journal," September, 1896.

CHAPTER VIII.

DISEASES OF THE LACRIMAL* APPARATUS.

Malposition of the Punctum Lacrimale.†—Inversion of the punctum accompanies entropion of the lower eyelid, while eversion of it is present with ectropium of the lid. A slight eversion, quite sufficient to cause epiphora, may exist without any marked ectropion of the lid, and it is these cases which more properly belong to this chapter. They are the result generally of some chronic, although it may be slight skin affection of the lower lid, which draws the inner end of the latter a little away from the eyeball.

The prominent symptom of this and of all the following lacrimal affections is lacrimation, or epiphora (*ἐπιφορὰ δακρύων, a sudden burst of tears*), a flowing of tears over the cheek.

Stenosis and **Complete Occlusion of the Punctum Lacrimale.**—Either of these conditions may result from conjunctivitis or from marginal blepharitis, although they may not appear for a length of time after those affections have passed away, and the original affection may have been so slight as to have escaped the observation of the patient. In stenosis the size of the punctum may become so extremely minute that even the normal flow of tears is too copious to pass through it. Complete occlusion is probably only a more advanced stage of stenosis.

The treatment, in cases of eversion of the punctum, of stenosis, and of complete occlusion is similar, namely, the opening up of the punctum, and its conversion into a slit: this is done with a Weber's knife (Fig. 90), the probe-point of which is passed into the punctum in cases of eversion, forced into the small opening in cases of stenosis, or forced through the usually thin covering of the punctum in cases of occlusion. In doing this the lower lid should be stretched tightly by a finger of the surgeon's left hand

* *Lacrima,* a tear.

† In this chapter, and elsewhere in the book, the terms punctum lacrimale and canaliculus refer to the inferior passage, unless it be otherwise expressly stated.

placed near the external canthus. The edge of the knife being now directed towards the eyeball, the instrument is pushed on a little into the canaliculus, until 2 mm. of the latter has been opened up, and it is then withdrawn. If the edge of the knife be directed outwards in this proceeding, the incision comes to lie on the outer edge of the intermarginal portion of the lid, and not in contact with the eyeball; consequently the tears are not carried away, and the disfigurement produced is considerable. A slitting up of the whole, or the greater part, of the canaliculus in these cases is unnecessary, and interferes with the physiological action of the tear passage. For two or three days after the little operation, it is necessary to pass a probe along the portion of the canaliculus which has been slit up, to prevent union taking place.



FIG. 90.

Obstruction of the Canaliculus.—The canaliculus may be diminished in its caliber, or entirely closed, by contraction the result of inflammation which had extended to it from the conjunctival sac. It is not possible to diagnose the presence of either of these conditions, which may be associated with stenosis or occlusion of the punctum lacrimale, except by the introduction of a very fine probe into the canaliculus. The passage may also be obstructed by an eyelash, a chalky deposit, or a mass of leptothrix.

The diagnosis of leptothrix in the inferior canaliculus is made by the following signs and symptoms: Lacrimation; the presence of a creamy-yellow discharge at the inner canthus, without dacryocystitis; congestion of the caruncle and neighboring parts of the conjunctiva. On everting the inner end of the lower lid, the region corresponding to the canaliculus is seen to be rounded and swollen on its conjunctival aspect. The lacrimal punctum is enlarged, stands out from the eyeball when the patient looks up, and is filled with the creamy exudation. On pressure, a hard cylindrical mass can be felt in the canaliculus. At a later stage, severe purulent inflammation of the canaliculus comes on, with marked swelling of the eyelid in the neighborhood; while the greenish-yellow dacryolith contained in the canaliculus usually consists of leptothrix or streptothrix, and actinomyces have also been found.

Treatment.—Where there is merely diminution in the caliber

of the passage, the introduction of probes, increasing in size, is frequently sufficient to effect a cure. If dilatation fail, recourse must be had to slitting up the canaliculus; but, if it can possibly be avoided—that is, if a less extended opening will answer—the passage should not be slit up in its entire length. At least 3 mm. of its median end ought to be left intact, as otherwise regurgitation of tears from the lacrimal sac is liable to trouble the patient ever afterwards. If the canaliculus be completely closed by adhesions, so that a fine probe cannot be pushed through it, it becomes necessary to rip it up with the point of any small knife, following the known course of the passage from the outside. If the canaliculus be closed as far as the opening into the sac, or if only at that point, the obstruction must be pierced with the point of a fine knife. A great difficulty in all these cases is to keep the passage patent when once formed. A plan which affords tolerable certainty of this is the frequent passage of probes into the sac until the tendency to closure seems to have ceased; but even under favorable conditions recurrences of the closure are apt to occur.

Leptothrix in the lower canaliculus is readily cured by slitting up the passage and evacuating its contents.

Stricture of the Nasal Duct is usually the result of simple swelling of its mucous membrane in a catarrhal attack, which has originated in the nasal mucous membrane; or of membranous or cicatricial contraction of the mucous membrane resulting from long-continued catarrh. It also occurs in consequence of disease of the bones of the nose—*e. g.,* in syphilis, acquired or congenital, and from blows which fracture the bridge of the nose.

Treatment.—Stricture due to acute inflammatory swelling of the mucous membrane should be treated by the injection of weak alum or other astringent solutions into the lacrimal sac, or down the nasal duct, by means of an Anel's syringe, and attention should be paid to the nasal mucous membrane. Probing here should not be attempted, lest it injure the delicate swollen mucous membrane of the duct.

Membranous or cicatricial strictures are best treated by means of probes in the manner proposed by Sir William Bowman. The inferior canaliculus is slit up to a slight extent so as to admit the point of one of Bowman's smallest probes, which is given a curve to suit that of the nasal duct. With the fingers of the left hand the surgeon stretches the lower lid, and, entering the probe into

the canaliculus, pushes it gently along its floor until the point reaches the lacrimal bone forming the posterior wall of the sac. The point being kept pressed against this bone, the direction of the probe is now altered by carrying its free end upwards towards the bridge of the nose until the point in the lacrimal sac is directed towards, or aimed at, the sulcus between the ala of the nose and the cheek. The probe, then, is in a position corresponding to the prolonged axis of the nasal duct, down which it is pushed with a slow and gentle motion. Any obstacles met with on the way are overcome, if possible, by an increase of the pressure; but if at any part of the proceeding much difficulty be encountered, rather than that any violence be used, all further manipulation should be postponed to another day, and it will often be found that at the second or third visit the probe is passed with comparative ease. Thicker probes are gradually introduced at successive sittings until the largest size has been reached.

The most common seats for membranous or cicatricial strictures of the nasal duct are at its entrance into the sac, where it is narrowest; and at its lower end, where it is most exposed to catarrhal processes spreading from the nostril.

Where there is reason to think that the stricture is due to chronic catarrhal swelling of the lining mucous membrane of the duct, astringent injections into the canal, in addition to the probing, are of use.

Otto Becker used very fine probes, which he passed by the upper canaliculus. Weber's probes are conical, and of very large caliber at their thickest part. Their inventor passes them by the superior canaliculus, but many other surgeons pass them by the lower. I do not employ these probes, because, when passed into the nasal duct, their thickest part, which is 3 to 4 mm. in diameter, corresponds with the upper end of the duct, which is its narrowest part, being only 3 mm. in diameter: consequently the probe becomes more or less impacted at this place at each operation, and is apt ultimately to give rise there to hypertrophy of the periosteum, and finally to stricture; so that, while the immediate effect of their use is good, the ultimate result is often the reverse. When used by the inferior canaliculus, their size makes it necessary to slit that passage in its entire length, and the entrance of the passage into the sac must be enormously dilated by so large an instrument, both of which circumstances are most undesirable. Indeed, I entirely agree with those who reject large probes, of

whatever shape or however introduced, in the treatment of lacrimal-duct strictures.

To prevent closure of the duct when once made free, Arthur Benson (Dublin) advocates the use of leaden styles, removable by the patient. He first divides the canaliculus (by preference the upper one), and dilates the stricture with probes in the ordinary way, and then introduces into the duct a piece of leaden wire 1.5 mm. to 2 mm. in diameter, cut to length, and smoothed off at the ends. The upper end is curved so as to lie out on the cheek. The style is at first removed daily, and the duct syringed, until any existing inflammation and discharge have almost ceased. The intervals are then increased; and as soon as practicable the patient is taught to remove the style and to replace it himself. When he is able to do this easily, he is directed to leave the style out for some hours each day, and finally to wear it only at night.

Stilling has proposed an operation, which he calls strictu-rotomy, for the cure of membranous obstructions in the duct. Having slit up the canaliculus, and ascertained with a probe the position of the stricture, Stilling passes his knife, with the cutting edge directed forwards, down the duct and through the stricture; he then withdraws it a little, turns the edge in another direction, and pushes it again through the stricture, and performs this maneuver a third time before removing the knife. On subsequent days large probes are passed. This method has never gained much acceptance.

Very obstinate membranous strictures can sometimes be freed by electrolysis.

The most favorable cases of stricture for cure are those due to inflammatory swelling of the mucous membrane, and next in order come those caused by membranous or cicatricial contraction, while those due to bony obstructions must be regarded as incurable.

Now and then cases of persistent lacrimation will be met with, in which the nasal duct and the rest of the lacrimal apparatus are in perfect order. These cases are often due to a catarrhal affection of the nasal mucous membrane, slightly involving the very lowest extremity of the nasal duct. Applications directed towards relief of the nasal affection are here indicated.

Blennorrhea of the Lacrimal Sac, or **Chronic Dacryocystitis,** is commonly caused, in the first instance, by stricture of the nasal duct. In consequence of this stricture the tears and the normal

mucous secretion of the lining membrane of the sac are retained, and offer favorable conditions for the development of the micro-organisms, which are constantly present on the surface of the eye, and are carried into the lacrimal sac by the tears. The decomposing contents of the sac set up inflammation of its mucous membrane, with discharge of a muco-purulent nature.

But cases of lacrimal blennorrhea are often met with in which no stricture of the nasal duct is found. In many of these cases there has been a stricture due merely to catarrhal swelling of the lining membrane of the duct, which has subsided in the course of time without treatment, and the duct has then again become free, while the lacrimal blennorrhea to which the stricture gave rise continues. It is very probable, however, that lacrimal blennorrhea may occasionally come on where there has never been a stricture of the nasal duct, and merely as an extension of catarrh from the nostrils, especially in cases of ozena, or as an extension of catarrh from the conjunctiva.

The patients usually complain of nothing more than epiphora. Those who are more observant of themselves may have noticed a swelling, which we call a lacrimal tumor or mucocele, in the region of the lacrimal sac; and also that the conjunctival sac, especially when the swelling is pressed upon, becomes now and then more or less filled with a somewhat thick and opaque discharge, which obscures the sight until wiped away. Occasionally there is no lacrimal tumor, for the contents of the sac may not be copious enough to distend it markedly.

In order to ascertain in each case of epiphora whether or not lacrimal blennorrhea be present, the surgeon presses with his finger over the lacrimal sac, when, if there be blennorrhea, the discharge will be evacuated through the puncta into the conjunctival sac. In those cases in which there is no longer a stricture of the nasal duct, the discharge may pass downwards into the nose, and the patient will feel it in his nostril, out of which he can blow it.

Conjunctivitis must sometimes be regarded, not as the cause, but rather as the effect of a lacrimal blennorrhea, by reason of the decomposing discharge from the lacrimal sac making its way into the conjunctival sac. Blepharitis, too, is seen as a further result of irritation from the discharge in old-standing cases.

Treatment.—It is important, in the first place, to ascertain whether there be a stricture of the nasal duct, and for this purpose

water should be injected by means of an Anel's syringe through the canaliculus into the duct. If the fluid make its way freely into the nose or pharynx, it may be taken for granted that the nasal duct is not obstructed; but if instead of passing through—or only under high pressure—it distends the lacrimal tumor to a greater size, a stricture may be assumed. If stricture of the nasal duct be present, it must be relieved, or all other measures will prove futile. Should there be no stricture, and also before and after any existing stricture has been freed, the treatment consists in the very frequent pressing out of the contents of the sac by the patient, so that no distention of it may occur: and in doing this he should endeavor to cause the discharge to pass down the nose rather than into the eye; while the surgeon, having, if necessary, dilated the canaliculus, injects astringent solutions into the sac daily, to relieve the catarrh. I find that protargol, in a 15 or 20 per cent. solution, is the best application for introduction into the lacrimal sac. The latter should first be washed out with a physiological salt solution.

The caustic treatment (recommended further on for acute dacryocystitis) is often of the greatest benefit in these chronic cases. Any existing conjunctivitis or nasal catarrh should be treated. But there are many cases in which no treatment short of obliteration or extirpation of the lacrimal sac will bring about a radical cure of this very troublesome complaint.

Obliteration of the sac is desirable in some very chronic cases. where repeated attacks of acute inflammation and fistula occur, or where there is constant discharge, and disease of bone, or when all other methods have failed to relieve the patient. This can be done by the application of a galvano-cautery to the lining membrane of the sac. In my experience, obliteration of the lacrimal sac is a most unsatisfactory proceeding.

Extirpation of the Lacrimal Sac is a preferable proceeding, and for it Kuhnt's method [1] is probably the best.

An incision down to the bone and about 2.5 cm. long is made over the anterior lacrimal crest. It begins 4 mm. above the internal palpebral ligament, and ends 5 mm. below the commencement of the bony lacrimal duct. As a rule the palpebral ligament is not divided by this incision, and has now to be separated with the scissors close to its insertion. If there be bleeding from the small arteries, they are seized and twisted. The temporal lip of the wound is now drawn aside, and the fibrous capsule of the sac

carefully incised along the crest, care being taken that the sac itself is not opened. The anterior wall of the sac, grayish blue and shining, then becomes exposed in the cavity. To remove the sac the surgeon separates its inner wall from the periosteum with the closed blunt ends of the scissors; then the fundus of the sac, along with the strong fibrous capsule which is here adherent to it, is drawn forwards, and with a few strokes of the scissors is separated from its bed, and the scissors are passed behind the sac (Fig. 91), and the posterior wall is similarly separated. The




Fig. 91.

temporal lip of the wound is now drawn forwards and outwards as much as possible by an assistant, while the surgeon draws the fundus of the sac inwards and forwards, and frees the outer and what remains attached of the anterior surfaces. Finally, the sac is cut off close to the bony canal, and, if there be no close stricture present, the mucous membrane of the duct is curetted with a sharp spoon. A small drain is inserted and the wound is closed by two or three deeply placed sutures, special care being taken to secure the palpebral ligament to its insertion.

Should the sac be opened during the operation, its total extirpa-

tion will be rendered much more difficult; and if a portion of the sac be left behind, the object of the operation is likely to be frustrated by a return of suppuration. If the sac has not been got out in its entirety, the suspicious places must be destroyed by curetting. There is very often a good deal of troublesome bleeding during the operation, which must be controlled by compression, torsion, and ligation, so that the operation may be carried out with the precision upon which its success depends. Where there is a fistula of the sac, or where there has been phlegmonous dacryocystitis, the operation is rendered much more difficult by reason of adhesions. Czermak considers it unnecessary and undesirable to divide the palpebral ligament, because in his opinion it is likely to lead to some disfigurement, no matter how carefully the sutures may be applied. He finds that the fundus of the sac can be got out without division of the ligament.

The lacrimation, after extirpation of the sac in cases of chronic dacryocystitis, is usually less than before. Should it continue to any distressing degree, removal of the lacrimal gland is indicated.

Acute Dacryocystitis (*δακρύω, to weep; κύστις, a bladder*). —Acute inflammation of the lacrimal sac most usually comes on when chronic lacrimal blennorrhea is already present. Caries of the nasal bones may cause it, and it occurs idiopathically, probably as the result of exposure to cold.

The region of the lacrimal sac and the surrounding integument become swollen and red, and these conditions often spread to the lids, giving an appearance which is sometimes mistaken for erysipelas; but the history of the case, showing the previous existence of lacrimal obstruction, etc. will assist the diagnosis. Great pain accompanies the inflammatory process. Gradually the region corresponding to the lacrimal sac becomes the most prominent one of the swelling, and the abscess, pointing there, opens. When the pus has been discharged the inflammation subsides, and the opening through the skin may either close, the parts resuming their normal functions, or the opening may remain as a permanent fistula.

The difference between chronic blennorrhea of the lacrimal sac and acute dacryocystitis, besides the fact that one is a chronic and the other an acute inflammatory process, is that the former process is confined to the mucous membrane of the sac, while in the latter the submucous tissue is involved, with phlegmonous inflammation as the result.

Treatment.—In the early stages poultices and purgatives should be employed. As soon as palpation of the sac indicates the presence of pus it must be evacuated. This can be effected either through the canaliculus, by opening it up to its entrance into the sac, or by an incision through the integument over the sac. The latter is the better method, as it admits of free access to the interior of the sac. The next day the walls of the sac are to be freely touched with solid mitigated nitrate of silver; or a plug of cotton-wool soaked in a strong solution of nitrate of silver may be inserted into its cavity, and left there for some hours; or various astringent solutions may be injected into the sac. The aim of the treatment, whatever it may be, is to secure a rapid return of the mucous membrane to its normal condition. If stricture of the nasal duct be present, it must be treated *pari passu.* By these means the discharge from the sac is arrested, and the external opening gradually closes.

If a fistula should form, it may be brought to close, in many cases, by simply freeing an existing stricture of the nasal duct; or it may be necessary to pare its edges, and bring them together by sutures; or, especially if there be a long fistulous passage, the galvano-cautery, in the form of a platinum wire, can be applied with advantage.

Dacryoadenitis (*δακρύω, to weep; ἀδήν, a gland*), or **Inflammation of the Lacrimal Gland,** occurs in an acute and in a chronic form, but is extremely rare in either. I have seen one case of acute purulent dacryoadenitis, but no instance of the chronic affection. Swelling and hyperemia over the gland and of the whole lid, with chemosis of the conjunctiva and much local pain, increased on pressure, are the most marked symptoms of acute dacryoadenitis. When suppuration has taken place the abscess may open into the conjunctiva, as it did in my patient, or through the skin. In the latter case it is liable to leave a fistula behind it; and, indeed, the chronic form may also, it is said, lead to fistula.

Numerous cases of chronic enlargement of both lacrimal glands have been recorded. Good results have been obtained by administration of potassium iodid or mercury in some cases.

Treatment in the early stages consists in poultices and purgatives. When pus has formed, the abscess may be opened through the skin or from the conjunctiva.

Hypertrophy of the Lacrimal Gland is also of rare occur-

rence. It may attain such dimensions as to push the eyeball out of its position. It can only be dealt with by—

Extirpation of the Lacrimal Gland.—This is performed by making an incision through the integument under the outer third of the orbital margin; the fascia under this is dissected up, the gland drawn out with a hook, and dissected out with a scalpel. Or, if it be considered sufficient to remove the palpebral portion only, this can be done from the conjunctival surface. It can be seen in the upper fornix, by separating the lids widely at the outer canthus, while the patient looks well down and to the nasal side. This partial removal may be performed for persistent lacrimation when other means fail.

Tumors of the Lacrimal Gland.— See chap. xix.

Reference.

[1] "Ueber die Therapie der Conjunctivitis Granulosa," p. 74. Jena, 1897.

CHAPTER IX.

DISEASES OF THE SCLEROTIC.

Inflammation of the sclerotic is not a common disease, although the diagnosis "scleritis" is often made by inexperienced persons, every redness of the white of the eye being taken for inflammation of the sclerotic. Beginners are warned against this error. Iritis, cyclitis, and conjunctivitis as well as scleritis, cause redness of the white of the eye.

The diagnosis from conjunctivitis is easily made by observing whether the conjunctival vessels can be moved over the affected part or not; while in iritis and cyclitis the ciliary injection is confined to the part immediately surrounding the cornea. Moreover, in iritis the appearance of the iris itself is conclusive; and in scleritis, as will just now be seen, the appearances are characteristic.

Scleritis attacks only that part of the sclerotic which is anterior to the equator of the eyeball, and is either superficial or deep. The superficial form is known as episcleritis. Yet it is not always possible to distinguish between these two forms in a given case, as the appearances in the early stages are very similar. They are probably only different degrees of the same disease. But the necessity of admitting the existence of two forms depends upon the different course they each take; the superficial form being a relatively harmless disease, while the deep form entails serious consequences.

Periodic Transient Episcleritis (Fuchs), or **Hot Eye** (Hutchinson).—This affection has been long known by the name given to it by Mr. Hutchinson,[1] and since then it has been described by Fuchs[2] under the title Episcleritis Periodica Fugax. It is characterized by frequently recurring attacks of inflammation of the episcleral connective tissue, giving rise to a vascular injection of a violet hue, but without any catarrhal or other secretion, or any hard infiltration, as in episcleritis of the usual type. It rarely attacks the whole sclerotic at one time, but is commonly confined to a quadrant or more, and wanders from one place to another.

When the attack subsides there is no stain left behind. The attack may be confined to one eye, or both may be affected, or it may go from one eye to the other. Pain, watering of the eye, and photophobia are present in varying degrees. Sometimes there is swelling of the eyelids. Occasionally the iris and ciliary body become inflamed, and also the retro-bulbar tissue, with resulting exophthalmos. The attacks last from one or two days to several weeks, and may recur once or twice a year and with intervals of only two or three weeks. Patients are usually liable to the disease for several years of their life. It attacks adults of middle age for the most part. Mr. Hutchinson assigns gout as the cause; but Fuchs has not been able to find any symptoms of that diathesis in his patients. Rheumatism and malaria seem sometimes to produce it, and in many instances no cause can be ascertained.

Treatment.—The long continuance of most of the cases shows that treatment has but little influence over the disease. Quinin and salicylate of soda internally are the remedies likely to be of most use, with warm fomentations, or the Japanese warmer, and a protection bandage locally during an attack.

Episcleritis appears as a circumscribed purplish, rather than red, spot close to, or 2 to 3 mm. removed from, the corneal margin. It is often unattended by pain, unless when the eye is exposed to irritating causes, and need not be elevated above the level of the sclerotic; but in severe cases there is a decided node at the affected place with more or less pronounced pain, which is increased on pressure. All the symptoms disappear in the course of a few weeks, and reappear at an adjoining place; and in this way, in time, the whole circumference of the sclerotic will have been attacked. The duration of the affection is usually long; and in those instances where the entire sclerotic becomes affected by degrees the process may last for years, on and off. Both eyes are often affected. The disease is liable to leave behind it a dusky discoloration of the sclerotic where each node was seated, but otherwise no harm to the eye ensues. But the patient should be made acquainted with the tedious nature of the affection. Very mild attacks of episcleritis will be met with, which pass away in a few days, and do not recur.

Causes.—The affection is often of gouty or rheumatic origin. It occurs sometimes in persons of scrofulous or syphilitic constitution; and it is more frequent in senior adults than in children

or young people, and more commonly attacks women than men.

Treatment.—No irritant should be applied to the eye. Local treatment should be confined to protection with dressing and bandages and warm fomentations, and the Japanese warmer is very useful. In addition to these, massage should be used, if there be not too great tenderness on pressure. Leeching at the external canthus is of use when the pain is severe. As regards internal remedies, where a syphilitic taint is present, mercury should be employed; if struma, cod-liver oil, maltine, etc.; or if, as is most frequently the case, rheumatism be the source of the evil, large doses of salicylate of sodium (say 20 grains four times a day) will often be found to act well. Salicylate of lithium is recommended in preference to the sodium salt by some. Iodid of potassium in large doses (20 grains four times a day, or oftener) is a useful remedy in some cases of this obstinate disease.

Deep Scleritis.—Here the whole of that part of the sclerotic which forms the front of the eye is more likely to be affected at once than in the milder form; although cases often enough occur where only an isolated node is present at a time.

It is the progress of the case alone which can render the diagnosis between this and the milder forms certain, and hence the importance of a guarded prognosis in the early stages of every case of scleritis. In the deep form changes—thinning and softening—of the scleral tissue take place, which render the latter less resistant, and consequently expose it to distention by even the normal intra-ocular tension. The result of this is a bulging (staphyloma) of the anterior part of the eyeball. This bulging in itself produces myopia, and has a deleterious effect upon the sight; but at a later period vision is often wholly destroyed by secondary glaucoma. It may happen that the thinning, etc., of the sclerotic affects only a portion, and not the whole, of its anterior surface; and in such a case the resulting staphyloma will be confined to that part of the sclerotic. A staphyloma, whether total or partial, presents a bluish-gray appearance, due to the uveal tract shining through the thinned sclerotic.

Either with or without such staphylomatous changes, sclerotizing opacity of the cornea may come on, and iritis, chorioiditis, and opacity of the vitreous humor are not uncommon complications, especially in strumous subjects. Both eyes are usually affected.

Causes.—Young adults are the most common subjects of deep scleritis, and it attacks females more often than males. Congenital syphilis, rheumatism, gout, struma, and disturbances of menstruation are the most common assignable causes.

Treatment.—There are few diseases less amenable to treatment. When any of the above causes can be assumed to be present, the suitable remedies are of course indicated. Besides this, a bandage when only one eye is affected, warm fomentations, or the Japanese warmer, dry cupping on the temple, or the artificial leech, complete rest of the eyes, and protection with dark glasses are to be recommended.

When all acute inflammation has passed away, an iridectomy is sometimes indicated—either for optical purposes, when the pupil is obstructed by corneal opacity, or for the purpose of reducing glaucomatous tension, or of diminishing a staphyloma.

Injuries of the Sclerotic.—Ruptures and perforating wounds are those which have to be considered. Mere losses of substance may be said not to occur.

The primary danger of a rupture or perforating wound of the sclerotic—apart from the loss of the contents of the eyeball, which is often associated with it—consists in the possibility of infecting organisms being introduced into the interior of the eye, and there setting up serious inflammatory reaction.

Ruptures are caused by blows on the eye, and are often indirect; thus, if the blow be received below the cornea, the rupture takes place above the cornea. A common cause of sclerotic ruptures amongst the agricultural population is a blow from a cow's horn while the animal is being tied up or fed in the byre. The rupture is usually from 2 mm. to 5 mm. distant from the corneal margin, and runs concentrically with the latter; and often the conjunctiva is not ruptured, but bridges over the opening in the sclerotic. Some of the contents of the eyeball may have been forced out through the rupture—*e. g.*, portions of the uvea, iris, and ciliary body, the vitreous, and the lens; and it is sometimes difficult at first to ascertain the exact state of affairs by reason of extravasated blood in the anterior chamber, under the conjunctiva, and in the tissues of the eyelids. When the conjunctiva is not ruptured, it is often advisable to confine treatment to the application of a bandage, as the covering conjunctiva acts as a protection against infection of the wound. Where serious damage has not been done to the retina, fair or even good vision may be regained

in many of these cases, which at first sight seem almost hopeless; and should perception of light be present, one may reasonably conclude that the retina is not detached. When the lens has been dislocated under the conjunctiva,—from whence it can be removed, after the sclerotic opening has closed,—the patient will of course require a glass, as after cataract operation, to give him the best vision.

A large and gaping perforating wound is easily recognized. A portion of the chorioid, ciliary body, or iris, according to the position of the wound, probably lies in it, or part of the vitreous humor may be found in it; while the vitreous humor, as seen through the pupil, will be full of blood (hemophthalmos), and blood may be present in the anterior chamber (hyphemia, *ὑπό, under; αἷμα, blood*), especially if the wound be far forwards. Small wounds may be concealed by subconjunctival hemorrhage, and here reduced tension of the eyeball is sometimes a valuable diagnostic sign.

A clean-cut perforating wound of the sclerotic may heal without inflammatory reaction, even when portions of the uveal tract or vitreous humor are prolapsed into it, these prolapsed parts becoming incarcerated in the cicatrix.

When inflammatory reaction follows upon one of these injuries it may either be of the purulent or plastic form. In the former case all the contents of the eyeball take part in the suppuration, and we term it panophthalmitis, phthisis bulbi being its ultimate result. In the plastic form the iris and ciliary body alone are implicated, and sight is slowly lost; the eye here, too, becoming phthisical. Of the two, the latter process is the more serious, as it is prone to give rise to sympathetic ophthalmitis—a danger which is not associated with the eye lost through panophthalmitis.

Where the wound has been produced by a small foreign body, which has remained in the interior of the eye, the seriousness of the position is much aggravated. This matter will be discussed in chap. xiv., on Diseases of the Vitreous Humor.

Treatment.—In cases where the wound is small (say less than 3 mm.), no suture need be used: a carefully applied dressing and bandage will be sufficient to promote the natural tendency to healing. But where the wound is large and gaping, any prolapsed chorioid, etc., must be freely irrigated with sublimate lotion, 1 in 5000, and completely reduced—or if the prolapsed parts, or portion of them, cannot be reduced, they must be abscised

—and the margins of the wound drawn together by a few points of fine silk suture passed through part of the thickness of the sclerotic; or, the sutures may be passed through the conjunctiva at some distance from the edges of the wound, the traction on the conjunctiva being often sufficient to close the scleral wound. A bandage is applied to each eye, and the patient is confined to bed. But, if the injury be such (very gaping wound, much loss of contents of the eyeball, or extensive intra-ocular hemorrhage) as to render restoration of useful sight, or at least retention of the shape of the eyeball, beyond reasonable hope, it is wiser to remove the eyeball at once, rather than to run the risk of sympathetic ophthalmitis without compensating advantage.

Tumors of the Sclerotic, as primary growths, are exceedingly rare; but fibroma, sarcoma, and osteoma have been so observed.

Pigment Spots of a yellowish-brown color are often seen in the sclerotic close to the corneal margin. They are congenital, and of no importance. Occasionally a black pigmented patch may be associated with pigmented sarcoma of the ciliary region.

References.

[1] "Trans. Ophth. Society, U. K.," v. p. 6.
[2] "Von Graefe's Archiv," xli. 4. p. 229.

CHAPTER X.

DISEASES OF THE UVEAL TRACT.

Inflammations.

If it be remembered that the iris, ciliary body, and chorioid * closely resemble each other histologically, that their blood supply is identical, and that they form with each other a continuous membrane, it is a matter of surprise to learn that any one of these three divisions of the uveal tract can undergo inflammation while the other two remain healthy. Yet this is by no means uncommonly the case. But it is, perhaps, more common for at least two of them, and especially the iris and ciliary body (*irido-cyclitis*), to be simultaneously inflamed; and the entire uveal tract may of course be affected at one time. If all three portions be affected, one of them may be much more affected than either of the others. Or, commencing in one portion, the inflammatory process often spreads to one or both of the other portions.

Clinically we cannot always know whether only one or more than one division of the uveal tract is in a state of inflammation, and this uncertainty of diagnosis is particularly liable to arise when there is severe acute iritis; for then the symptoms present might all be derived from the iritis alone, while the contracted and obscured pupil, the opacity in the aqueous humor and cornea, and the irritability of the eye, render impossible a diagnosis of chorioiditis by the ophthalmoscope, and the position of the ciliary body puts it at all times out of reach of ophthalmoscopic examination. It may be taken for granted that in every rather severe case of iritis, particularly in those of syphilitic origin, more or less cyclitis is also present; while a deep anterior chamber, tenderness on pressure, or punctate deposits on the posterior surface of the cornea increase the suspicion. In most cases of slight iritis there is probably no cyclitis.

It is only after the acute inflammatory symptoms have subsided and the pupil has become clear that disseminated changes in the

* *χόριον, the chorion;* hence chorioid, *like the chorion.*

chorioid, opacities in the vitreous humor, and even retinitis and optic neuritis, which may lead to optic atrophy, can be discovered, with their corresponding depreciation of vision.

It is convenient, in a systematic consideration of inflammation of the uveal tract, to discuss it under the separate headings of iritis, cyclitis, and chorioiditis.

Inflammations of the Iris.—The *Signs of Iritis,* more or less marked, are:

Discoloration, loss of luster and of distinctness of pattern, and functional disturbances (impaired mobility) of the iris, with contraction of the pupil. The loss of luster and of distinctness of pattern is due to an alteration in the endothelium, which covers the surface of the iris, to the presence of lymph in it and on it, and to cloudiness of the aqueous humor through which it is seen, caused by inflammatory products held in suspension; and there is often, too, some cloudiness of the cornea. The change in color is due to hyperemia of the iris, as well as to the presence of the inflammatory products; a blue iris becomes greenish, a brown iris yellowish. The impaired mobility and the contracted pupil are due to engorgement of the blood-vessels of the iris, to spasm of the sphincter iridis, and to posterior synechia.

Exudation of inflammatory products is present in greater or less degree, and may be found on either surface of the iris, in the pupil, in the aqueous humor, on the posterior surface of the cornea, and in the tissue of the iris.

Posterior synechiæ *—*i. e.,* adhesions between the iris and the anterior capsule of the lens—occur as the result of inflammatory exudation on the posterior surface or on the pupillary margin of the iris. The presence of posterior synechiæ is ascertained by observing the play of the pupil when the eye is placed alternately in strong light and in deep shadow, or by observing the effect of a drop of atropin solution on the pupil, the latter dilating only at those places where there are no synechiæ. If the entire pupillary margin have become adherent, the condition is termed complete posterior synechia, circular posterior synechia, ring synechia, or exclusion (or seclusion) of the pupil; and in such cases, especially if of some standing, atropin has no effect on the pupil. If the area of the pupil be filled with exudation, circular synechiæ being usually also present, the condition is known as occlusion of the pupil.

* *συνέχειν, to bind together.*

Total posterior synechia is that condition in which the whole posterior surface of the iris is adherent to the capsule of the lens, and is rarely the result of ordinary iritis, but is seen frequently in sympathetic ophthalmitis.

In addition to the foregoing, circumcorneal injection of the ciliary vessels is a common symptom in most cases of iritis.

The subjective symptoms in iritis consist, in the first place, of pain, due to irritation of the ciliary nerves in the inflamed part. Yet this pain is not always referred to the eye itself, but often appears in the form of supra-orbital neuralgia, or affects the infra-orbital division of the fifth nerve. Dimness of vision is the second subjective symptom of iritis. It may be due to cloudiness of the aqueous humor, to deposits on the cornea, to exudation of lymph on the pupillary area of the anterior capsule of the lens, or, where the ciliary body is implicated, to opacities in the vitreous humor.

Cases of iritis in which there has been no pain and no circumcorneal injection, and in which the failure of sight alone it is which brings the patient to the surgeon, are not uncommon. Examination then discovers the presence of extensive posterior synechiæ, which have probably been gradually forming for a long time back. These cases of quiet iritis are, in my experience, usually due to rheumatism (*vide infra*).

A mistake into which beginners very often fall is to take a case of iritis to be conjunctivitis or scleritis (see pp. 95 and 244), the "redness of the white of the eye" being that which misleads. The appearance of the iris itself will chiefly assist in the diagnosis. Moreover, the pain in iritis is of neuralgic character, but in conjunctivitis it is similar to that caused by a foreign body in the conjunctival sac. In iritis there is no discharge, while in conjunctivitis the eyelids are gummed in the morning by mucopurulent secretion. Of course iritis and conjunctivitis may occur together.

Those cases of iritis in which the inflammatory exudation is mainly on either surface of the iris and in the pupil are the most common. Here the circumcorneal injection is generally well marked, sometimes causing elevation of the limbus of the conjunctiva, and even general, although slight, chemosis. In very mild cases, however, as also in chronic cases, the injection may be slight. The loss of luster and of distinctness of pattern of the

iris is well marked, and there is considerable change in the color of the iris. Posterior synechiæ are very apt to form. In some rare cases of this form of iritis an enormous quantity of gelatinous exudation is present in the anterior chamber. In Secondary Syphilis one often sees iritis of this kind.

Rheumatic Iritis, too, is sometimes of this kind, but accompanied by circumcorneal injection, which is great in proportion to the other signs of iritis present. The pain in rheumatic iritis is often peculiarly severe. Yet, as I have already stated, quiet iritis is most often due to rheumatism.

Gonorrheal Iritis is also of this kind, although with it there is often seen the punctate deposit on the posterior surface of the cornea. It does not attend on, nor immediately follow, a gonorrhea; but an attack of rheumatic arthritis, usually of the knees, always intervenes. Gonorrheal iritis is extremely rare.

Those cases of iritis which are chiefly characterized by the deposit of fibrinous elements, as very fine pale buff-colored dots on the posterior surface of the cornea, with more or less turbidity of the aqueous and some tendency to the formation of posterior synechiæ, used to be, and very often are still, although erroneously, called cases of Serous Iritis, while the fine dotted appearance on the back of the cornea was commonly known as "keratitis punctata," a term for which "unctate corneal deposits" should be substituted. We know now that the inflammatory product is fibrinous and not serous, and that in these cases the ciliary body may be quite as much, if not more, affected than the iris, and hence that these are to be regarded as cases of iridocyclitis of a sluggish or chronic form. The fibrin passes from the ciliary body into the aqueous humor, and from it is precipitated as fine buff-colored or pale yellowish dots on the cornea in its lower quadrant by force of gravity. The part of the cornea which is thus affected is often of a triangular shape, the base of the triangle corresponding with the lower margin of the cornea, the apex being directed towards the center of the cornea, with the finer dot near the apex. The triangular shape is a mechanical result of the motions of the eyeball. In many cases, however, nearly the whole cornea is more or less affected.

In cases where the punctate corneal deposits continue for a length of time, in consequence of the resulting degeneration of the posterior epithelium, permanent secondary changes in the true cornea take place, and a consequent peculiar triangular

opacity at the lower part of the cornea will ever afterward indicate the nature of the process which has begun before.

In this form of iritis the circumcorneal injection is slight, the anterior chamber is often deep, and the aqueous humor is sometimes cloudy. The intra-ocular tension is frequently increased, from the presence of exudation in the angle of the anterior chamber.

Where the inflammatory product is situated in the tissue of the iris, the consequent swelling may be present over its whole extent, or may be confined to a circumscribed part of it. In the latter case the swelling is sometimes called a condyloma. The color of the iris changes remarkably at the affected part to a yellowish or reddish-yellow hue, and new vessels are formed in it.

In Syphilis, late in the secondary stage, a form of iritis occurs which may be always recognized as syphilitic. It is characterized by the formation of circumscribed tumors or small condylomata of a yellowish color, the rest of the iris being apparently intact. These tumors vary in size from that of a hemp-seed to that of a small pea, and are situated usually at the pupillary margin, occasionally at the periphery of the iris, and very rarely in the body of the iris. There may be but one tumor present, and there are seldom more than three or four. This form is not common. But many authors hold that in most, if not in all, cases of syphilitic iritis condylomatous tumors are present, though of such small size as to escape detection with our ordinary clinical methods.

Hemorrhagic iritis is not a special form of iritis, but is merely a severe inflammation of the iris with hyphemia. It is chiefly seen in iritis due to operations and injuries, in some cases of sympathetic iritis, and in old people.

Symptoms of Iritis in General.—(1) Pain. This is situated not so much in the eye as in the brow over it, in the corresponding side of the nose, and in the malar bone, and may even extend to the whole side of the head. It varies in its intensity; it is usually more severe at night, and is often called neuralgia by the patients. That form of iritis with exudation on the surface of the iris and in the pupil is the one attended by the most severe pain; the form with punctate deposits on the posterior surface of the cornea as its main characteristic is generally unattended by pain; while the form with marked circumscribed deposits or condylomata in the stroma of the iris is in many cases excessively painful and in some completely painless. (2) Lacrimation and

photophobia are occasionally present, but never to such a degree as is often observed in some corneal affections. (3) Dimness of vision. This is usually complained of as soon as the inflammation is pronounced. Cloudiness of the aqueous humor and punctate corneal deposits affect sight in proportion to their degree; and exudation in the pupil may reduce vision to a quantitative amount.

The tension of the eye in iritis is usually normal, but in some violent cases it will be found to be high, as also in some cases where, while the inflammatory symptoms are but slight, the punctate corneal deposits are marked, and in cases, too, where the angle of the anterior chamber is blocked.

Prognosis.—The length of duration of an attack of iritis cannot be foretold at the outset. Cases which are in other respects mild—*i. e.,* where the pupil dilates well and rapidly to atropin, where the aqueous humor is clear, and where but little lymph is thrown out—often continue for weeks irritable and painful, with a marked tendency to relapse if treatment be at all relaxed. An attack of iritis may last from two to eight weeks. Recurrences of the inflammation are common, owing to continuance of the constitutional taint, which gave rise to the iritis in the first instance.

It is possible that an attack of any form of iritis, if carefully treated from the beginning, may leave the eye in as healthy a condition as before; but it is quite as common, in spite of every effort, to find posterior synechiæ, isolated or as a circular synechia, left behind.

The presence of a few isolated synechiæ, if the pupil be clear, is in itself harmless to sight; but if relapses take place and fresh adhesions be formed, a complete posterior synechia may ultimately be established. When this occurs, the aqueous humor being still secreted behind the iris, and being unable to escape into the anterior part of the chamber, the iris becomes bulged forward, like the sail of a ship, until it touches the peripheral part of the cornea; while the center of the anterior chamber retains its normal depth. This condition is very liable to induce glaucomatous tension (secondary glaucoma) and consequent loss of vision; or, if the eye escape this danger, chronic inflammation of the ciliary body and chorioid—so-called chronic irido-cyclitis, or irido-chorioiditis—may develop, and may lead to diminished tension and phthisis bulbi, with detachment of the retina and calcifi-

cation of the lens; or, the eye having been first blinded by high tension, may at a later period undergo phthisis bulbi.

Complete posterior synechia may of course result from the first and only attack of iritis, and not by means of repeated relapses.

In some cases of iritis the vitreous humor becomes more or less opaque, and this condition does not always disappear as the iritis gets well; indeed, it may not be possible to ascertain its presence until after the inflammatory process in the iris has subsided. Very great and permanent deterioration of vision may result in such instances; and they emphasize the importance of a cautious prognosis at the commencement. It is probable that in these cases the ciliary body participates in the inflammation, and is mainly responsible for the opacities in the vitreous.

Causes.—Iritis is not common in children, except as complicating a corneal process, or as a result of congenital syphilis or tuberculosis (see New Growths of the Iris, p. 289). Towards puberty slight iritis is sometimes found in girls. Youth and middle age are the times of life in which iritis is most often seen, while in old age it again becomes rare.

More than 50 per cent. of the cases depend on syphilis, and a large proportion of the remainder are due to rheumatism. During desquamation after smallpox iritis is sometimes observed. In metria and septicemia purulent iritis occurs, as also with typhoid fever, pneumonia, and recurrent fever. Diabetes sometimes causes iritis.

Treatment.—A mydriatic (see table, p. 316), is, above all others, the most important means. Most commonly a solution of atropin is used (Atrop. sulph. gr. iv, Aq. dest. ℥j) as drops. An atom of sulphate of atropin in substance, placed in the conjunctival sac, gives a very active reaction. It is also used in the form of an ointment (Atrop. sulph. gr. iv, vaselin ℥j), and gelatin discs containing atropin are manufactured. By paralyzing the sphincter iridis atropin provides rest for the inflamed iris; and if adhesions have already formed, the dilatation of the pupil may break them down, while if none are as yet present, the dilatation will greatly aid in preventing their formation. To produce a maximum effect, where it is desired to break down adhesions, six drops of the atropin solution should be instilled into the eye, with an interval of from five to ten minutes between each; and in this way the atropin from each drop has time to make its way into the anterior chamber, and finally the accumulated effect

of all six is obtained. More than one drop can hardly be retained in the conjunctival sac at a time. The use of cocain (2 per cent.) along with atropin insures a maximum dilatation. A drop in the eye from twice to four times a day is required, in order to maintain the desired dilatation of the pupil *ad maximum.*

Some individuals are peculiarly susceptible to Atropin Poisoning, of which the symptoms are dryness of the throat, fever, fullness in the head, headache, delirium, coma. The antidote is morphia, of which 1-4 grain used hypodermically neutralizes 1-30 grain of atropin in the system. Atropin poisoning can occur by the introduction of the solution into the stomach through the lacrimal canaliculi, nose, and fauces; and to prevent this the finger (of the patient) may be placed in the inner canthus, so as to occlude both canaliculi during, and for some moments after, the introduction of the drop into the eye.

After long use of atropin the skin of the lower eyelid, or of both eyelids, often becomes eczematous, red, swollen, and painful, from infiltration with the drug; and in other cases follicular conjunctivitis is induced. If these complications occur, sol. extr. belladonna (gr. viij ad ℥j) should be substituted for atropin, and suitable remedies used for skin or conjunctiva.

In old people tenesmus and retention of urine sometimes result for use of atropin.

Atropin, while it is so useful a means in the treatment of inflammations of the iris, ciliary body, and cornea, is of no benefit in many other diseases of the eye, and is positively harmful in some of them. It is necessary to make this statement very explicitly, for some—perhaps I should say many—medical men who have not devoted attention to the subject of eye-disease include atropin in every eye-lotion they prescribe. If the disease prescribed for be conjunctivitis, as it very often is, the atropin is calculated rather to increase than to relieve the conjunctival affection; while, if the patient be advanced in life, there is always the danger that a tendency to glaucoma may be present, and in such a case the dilatation of the pupil caused by the atropin will be sufficient to bring on an attack of acute glaucoma. In these days it falls to the lot of most ophthalmic surgeons to be called, at one time or another, to a case of acute glaucoma brought on by the gratuitous use of atropin in this manner. It is to be feared that the reason for this random prescribing of atropin is to be found in an ignorance of diagnosis, which leads practitioners to throw

atropin with a number of other drugs into their eye-lotions in the hope that some of the ammunition will hit the mark, wherever the latter may be.

Dark protection spectacles should be worn by patients suffering from iritis; and in severe cases, especially in cold weather, a bandage should be worn, and the patients should be confined to a dark room, and even to bed.

In that form of iritis where the inflammatory exudation is mainly on the surface of the iris and in the pupil, iodid of potassium or perchlorid of mercury may be given internally. If there be much irritation, pericorneal injection, or chemosis, leeching at the external canthus is of use. Intermittent warm fomentations (every two hours) promote healthy vascular reaction. Pain is to be relieved by hypodermic injections of morphia, and by chloral internally.

In rheumatic iritis and in iritis due to diabetes, salicylate of sodium in large doses (20 to 30 grains every three hours) has often a remarkably favorable effect.

In many of those cases in which punctate deposits on the cornea are the chief characteristic a small quantity of atropin will suffice, as there is often but little tendency to the formation of synechiæ, and, the irritation being slight, leeching is unnecessary. The skin (Turkish baths and dry rubbing), kidneys, and bowels should be acted on; and to the diuretics prescribed some iodid of potassium may be added. Turpentine in ʒj doses, as recommended by Carmichael, of Dublin, is often a useful remedy here.

Blistering on the temples or behind the ears is with many surgeons a favorite remedy. It adds to the annoyance of the patient, but as a remedy in this, or, indeed, in any other eye disease it is useless.

Great care is required in watching the tension of the eye in this form of iritis; and, if it be found to increase and to remain high for three or four days, paracentesis of the anterior chamber must be performed to reduce it temporarily while the iritis is still progressing towards cure. This little operation is exceedingly useful if there be much deposit on the posterior surface of the cornea, as by means of it the deposit, to a great extent, may be floated away. Indeed, in many cases of punctate corneal deposits, repeated tapping of the anterior chamber is a valuable curative measure. It should be performed, say, on three successive days,

and similarly again after an interval of four or five days, and so on, according as the condition of the eye permits. (For mode of performing paracentesis, see p. 147.)

Where the exudation is in the form of circumscribed tumors or condylomata in the stroma of the iris, it is important to obtain rapid absorption of the inflammatory products, which are so abundantly thrown out, and which, in an organ like the eye, would soon cause extensive destruction. Consequently, the system should be put under the influence of mercury as quickly as possible, by the use of inunctions of mercurial ointment or by small doses of calomel internally; and this treatment is indicated, even when the inflammation is not of syphilitic origin. Warm fomentations are useful. An after-treatment with iodid of potassium is to be employed.

In Syphilitic Iritis von Graefe gave the following:

℞ Hydrarg. biniodid. gr. vj; Potass. iodidi, ʒ iss.; Aq. destill. ℥ ss. Syr. Aurant. ℥ iiss. M. A teaspoonful to be taken once a day. The dose to be gradually increased.

In Purulent Iritis quinin and salicylate of sodium are the most suitable internal remedies.

Inflammation of the Ciliary Body; Cyclitis.—Cyclitis, as already stated, is often present with inflammatory affections of the iris or chorioid, although its presence in many of these cases cannot be clinically determined; but primary idiopathic cyclitis is rare.

The Symptoms of Cyclitis, although not always all of them present, consist in marked circumcorneal injection, ciliary neuralgia, pain on pressure of the ciliary region, deep anterior chamber, opacity in the anterior part of the vitreous humor, punctate deposits on the posterior surface of the cornea, and sometimes hypopyon in the anterior chamber. Edema of the margin of the upper lid is also frequently present.

There are three forms of cyclitis:

In the first the circumcorneal injection is but slight. The anterior chamber is often at first deeper than normal, owing to hypersecretion of aqueous humor from the ciliary body; there are punctate opacities on the posterior surface of the cornea; and the anterior part of the vitreous humor is filled with fine dust-like opacities. Iritis may come on, and the danger of glaucomatous increase of tension is great, owing to the tendency to blocking of

the anterior chamber with inflammatory exudation. Unless increase of tension gives rise to it, pain is not often present. Primary idiopathic cyclitis is usually of this form.

In the second form there is much plastic exudation, the circumcorneal injection is very marked, and there is venous congestion of the iris. The anterior chamber is deep, owing to retraction of the periphery of the iris by inflammatory exudation in the ciliary body, and owing to this retraction, too, the pupil is dilated. The inflammation may extend to the iris or to the chorioid, and in the latter case the vitreous may become very opaque. Violent ciliary pains attend the affection, and the eyeball is very tender on pressure of the ciliary region. The intra-ocular tension is reduced.

In purulent cyclitis, which is the third form, the circumcorneal injection is very well marked. The vitreous humor is filled with membranous opacities. There is hypopyon in the anterior chamber, which has the characteristic of appearing and disappearing at intervals of a few days. There is severe ciliary neuralgia; and purulent iritis, or chorioiditis, or both, are apt to supervene.

Prognosis.—In an early stage all these forms are capable of undergoing cure and of leaving the eye in a fairly useful condition. On the other hand, the first form, as already stated, is liable to produce secondary glaucoma; the purulent form leads to atrophy of the iris and chorioid, disorganization of the vitreous humor, detachment of the retina, cataract, and phthisis bulbi; and the form with much fibrinous exudation, when it is due to a perforating injury, in addition to serious damage to the affected eye, similar to that produced by purulent cyclitis, has the tendency to cause sympathetic uveitis of the other eye. The shrunken eyes resulting from the affection are often very liable to fresh attacks of inflammation, and frequently remain painful to the touch, circumstances which indicate that chronic cyclitis is still present, and, when the original cause of the inflammation has been a perforating injury, these stumps are a constant source of danger to the sound eye from sympathetic ophthalmitis.

Causes.—Traumata are the most common causes of the affection. Both the fibrinous and the purulent forms are liable to occur after cataract operations, as the result of infection of the wound, but this has now become very rare, owing to the careful aseptic measures employed.

The Treatment for cyclitis is similar to that for iritis. Leeching at the outer canthus is often of great benefit.

Inflammations of the Chorioid.—There are two great forms of inflammation of the chorioid (*χόριον, the chorion;* hence chorioid, *like the chorion*), the exudative and the purulent. Of the exudative form, again, there are several kinds, namely, disseminated chorioiditis, central chorioiditis, central senile chorioiditis, guttate chorioiditis, and syphilitic chorio-retinitis.

Disseminated or Exudative Chorioiditis.—The usual Ophthalmoscopic Appearances of this disease consist either in round white spots of different sizes with irregular black margins, or in small spots of pigment, these changes being surrounded by healthy chorioidal tissue; or there may be few or no white patches, but rather spots of pigment surrounded by a pale margin. The retinal vessels pass over, not under, the patches. The number of these patches or spots varies according to the intensity of the disease. Their position is at first at the periphery of the fundus only, but later on they appear also about the posterior pole of the eye.

These appearances represent a rather late stage of the disease, the early stage not usually coming under observation. It consists in small circumscribed plastic exudations into the tissue of the chorioid, which, seen with the ophthalmoscope, give the appearance of pale pinkish-yellow spots. The exudations may undergo absorption, leaving the chorioid in a fairly healthy state; but more usually they give rise to atrophic cicatrices, in which the retina becomes adherent, with proliferation of the pigment-epithelium layer in their neighborhood, and hence the white patches with black margins above described. This is the form of chorioiditis which is often associated with the inflammatory processes in the iris or ciliary body, either as a primary or secondary affection. But, again, in many instances the disease does not extend beyond the chorioid.

Sometimes, in addition to the above changes, the pigment-epithelium layer all over the fundus becomes atrophied, exposing to view the vascular network of the chorioid, while here and there small islands of pigment are present.

Opacities in the vitreous humor are sometimes found.

Symptoms.—Diminution in the visual acuity, especially if the macula be involved; there may also be subjective sensations of light or colors, positive scotoma (a dark area visible to the

patient), and distortion of objects (metamorphopsia), or alteration in their size (megalopsia and micropsia). Night-blindness is not uncommon.

Causes.—Disseminated chorioiditis is due to acquired syphilis in a considerable number of the cases, and possibly in some it may be tubercular. But in a very large proportion of cases no ascertainable cause exists; and those, there is reason to suspect, are congenital, and probably many of them are dependent on an inherited syphilitic taint. In eyes with congenital cataract, patches of chorioiditis are often found.

Prognosis.—Disseminated chorioiditis is always a serious disease, and complete recovery cannot be looked for. The degree of defect of sight it may cause depends much on the extent to which the region of the macula lutea has become involved.

Treatment.—In fresh cases due to acquired syphilis a prolonged, but mild, course of mercurial inunctions is the most suitable measure, to be followed by a lengthened course of treatment with iodid of potassium. Where an inherited syphilitic taint is suspected, iodid of iron or iodid of potassium internally may be of use; while, in the cases due to other causes, small doses of perchlorid of mercury may be given; and in all cases, from whatever cause, dry cupping on the temple, or even the artificial leech, should be employed. Dark protection spectacles should be worn, and absolute rest of the eyes from all near work insisted upon so long as the disease is active. Subconjunctival injections of 4 per cent. of common salt are also used in these cases, and I find this method very useful.

Syphilitic Chorioido-Retinitis.—See Syphilitic Retinitis, chap. xv.

Central Senile Guttate Chorioiditis.—Under this name an appearance has been described by Mr. Waren Tay and others which consists of fine white, pale yellow, or glistening dots, best seen in the upright image, and situated chiefly about the macula lutea, or between this and the optic papilla. These dots are due to colloid degeneration with chalky formations in the vitreous layer of the chorioid, which give rise to secondary retinal changes. The appearances must not be confounded, as they sometimes have been, with those of retinitis punctata albescens (chap. xv.), which is an entirely different disease. The functions of the retina usually suffer in a marked manner, so that a partial central

scotoma may be produced; but some cases have been observed in which vision was but little, or not at all, affected.

This disease attacks both eyes, either simultaneously or with an interval, and is most often seen in persons of advanced life, although also found in middle age, and even in youth. It should always be looked for in cases of incipient cataract; for when the lental opacity is more advanced it cannot be seen, while functional examination does not then detect it.

Treatment is of no avail.

Central Chorioiditis.—This is an exudation at the macula lutea, without any similar disease elsewhere in the fundus. Absolute central scotoma is its prominent symptom, and syphilis its usual cause.

Treatment.—Active mercurialization; and where this can be adopted early the prognosis for recovery of sight is fair.

Purulent Chorioiditis.—This consists at first in a purulent extravasation between the chorioid and retina, and into the vitreous humor, recognizable by the yellowish reflection obtained from the interior of the eye on illuminating it. The eyeball may become hard, the pupil dilated, and the anterior chamber shallow. Purulent iritis with hypopyon soon comes on, and the cornea may also become infiltrated and slough away. There is usually considerable chemosis. The eyeball is pushed forwards by inflammatory edema of the orbital connective tissue. The eyelids are swollen and congested. There is intense pulsating pain in the eye, and radiating pains through the head; and in this stage all the tissues of the eyeball are engaged in the purulent inflammation, and the condition is termed Panophthalmitis.

Purulent chorioiditis does not always reach this latter stage, but may remain confined chiefly to the chorioid, vitreous humor, and iris. The pain in such cases is not severe; and when the affection occurs in children it may be mistaken for glioma; indeed, the name pseudo-glioma has, unfortunately, been given to it. It is distinguished from the malignant disease by the muddy vitreous usually present with it, by the posterior synechiæ, and by the retraction of the periphery of the iris, with bulging forwards of its pupillary part.

Prognosis.—The ultimate result in the vast majority of cases is loss of sight, with phthisis bulbi. The severe cases go on to bursting of the eyeball through the cornea or sclerotic, after which the pain subsides. It would seem from the description of

authors who have seen much of epidemic cerebro-spinal meningitis, that a certain number of cases of irido-chorioiditis occurring in the course of that disease do recover with retention of good sight.

The shrunken eyeballs produced by panophthalmitis are not generally painful on pressure, and it is remarkable that they are not very liable to give rise to sympathetic ophthalmitis, which latter observation is also true of the acute purulent process itself. It is traumatic cases of plastic irido-chorioiditis which produce sympathetic ophthalmitis.

Causes.—The most common causes of purulent chorioiditis are wounds of the eyeball, whether accidental or operative; foreign bodies piercing and lodging in the eyeball; and purulent keratitis. It may also come on suddenly in eyes which are the subjects of incarceration of the iris in a corneal cicatrix through infection of the incarcerated iris.

It is seen as embolic or metastatic chorioiditis, in connection both with epidemic and sporadic cerebro-spinal meningitis (chap. xix.); in some cases of metria, similarly as purulent retinitis (chap. xv.); in pyemia of the ordinary type; and in endocarditis.

In infancy and childhood, besides its occurrence with cerebro-spinal meningitis, it has been known to be caused by, or associated with, inherited syphilis, measles, bronchitis, diarrhea, whooping-cough, and omphalo-phlebitis; and it is more than probable that in every idiopathic case some infective blood-disease is the fundamental cause of the process, although it may not always be possible to determine its existence.

Treatment may be said to be powerless in this disease. The utmost one can do is to endeavor to diminish the pain in the very severe cases by warm fomentations, poultices containing powdered conium leaves, hypodermic injections of morphium, or, finally, by giving exit to the pus by a free incision in the eyeball, followed by a copious irrigation with weak sublimate lotion, so as to wash out the whole contents of the scleral cavity. Quinin and chlorate of potash are suitable internal remedies.

I agree with those who think that enucleation of the eyeball should not be undertaken during purulent chorioiditis in the acute stage, as it is liable to lead to purulent meningitis and death; but some surgeons do not recognize any such danger, and practice enucleation in this condition.

Sympathetic Ophthalmitis, and Sympathetic Irritation.

Introductory.—By Sympathetic Ophthalmitis we understand a general plastic uveitis (plastic inflammation of iris, ciliary body, and chorioid) of one eye, which has been caused by a plastic uveitis of the other eye, the latter condition being due to a perforating trauma, or other perforation of the eyeball. Or, the sympathetic ophthalmitis occasionally consists in an optic neuritis. That sympathetic ophthalmitis can be caused without a perforation of the coats of the first eye is exceedingly unlikely.

There are no such diseases as sympathetic cataract, conjunctivitis, detachment of the retina, keratitis, scleritis, etc.

The term "sympathetic" in this connection is an old one, and, we may safely say, would not be employed had the disease to be named at the present time. It has indeed been proposed to substitute the title Ophthalmia Migratoria, but, although preferable, as indicating the prevailing theory of the causation of the disease, it has not come into general use, and the old name will probably always remain the current one. Nor is it a matter of importance what the name may be, save in so far as it may give an erroneous impression concerning the nature of the disease.

It is well, therefore, to state at once that sympathetic ophthalmitis is not a fifth-nerve reflex phenomenon. It is a uveitis, and often also an optic neuritis, which, according to the generally accepted theory, has arisen by direct transmission or migration of bacteria, by way of the optic nerves and optic commissure, from the injured eye, in which uveitis has been set up, to the sympathizing eye. No sympathetic uveitis can be developed in the second eye until after uveitis in the first eye has commenced. A perforating injury of the eyeball, or other perforation, which does not produce uveitis in that eye, does not give rise to sympathetic ophthalmitis in the fellow eye. Traumatic uveitis in the injured eye does not cause sympathetic ophthalmitis in every case.

But while sympathetic ophthalmitis is not a reflex condition, there is an affection known as sympathetic irritation, which is a true fifth-nerve reflex neurosis; and the two affections, although sometimes closely associated clinically, are quite distinct from each other.

We speak of the eye that has received the perforating injury as the exciting eye, and its fellow, which becomes the subject of sympathetic ophthalmitis, as the sympathizing eye. The eyes are also spoken of as the injured eye and the sympathizing eye; also as the first eye and the second eye.

Before treating of sympathetic ophthalmitis, it will be convenient to state what is necessary concerning sympathetic irritation.

Sympathetic Irritation.

This may be caused by almost anything which produces irritability of the ciliary nerves in the first eye—*e. g.*, foreign bodies on the cornea or under the upper lid, losses of substance of the corneal epithelium, anterior staphyloma, acute glaucoma, iritis, dislocation of the crystalline lens, etc., etc. Amongst the causes of sympathetic irritation is an irritable shrunken globe, whether the latter condition be the result of a uveitis from a perforating injury, or of an idiopathic uveitis; and a shrunken globe may rise to sympathetic irritation at any time, even after many years. Having remained quiet for so long, the shrunken eye begins to lacrimate, and becomes painful and injected. A fresh injury to the stump may be the cause of this, or it may be ossification of its chorioid, and the irritation, whatever its cause, may be transmitted to the sound eye.

The most common symptoms of sympathetic irritation are: photophobia, lacrimation, vascular injection of the front of the eyeball, and accommodative asthenopia, and, in a well-marked case, these symptoms may become intensely distressing to the patient. Neuralgia in the orbit and brow, and retinal asthenopia, sometimes occur. Amblyopia has also been recorded, but the existence of a sympathetic amblyopia has not, I think, as yet been satisfactorily established.

Now we know (*vide infra*) that sympathetic ophthalmitis, too, is often caused by a shrunken fellow eyeball in which uveitis, indicated by the presence of irritability, is present; and of great interest and importance is the question: What relation, if any, has sympathetic irritation to sympathetic ophthalmitis in such cases? Is sympathetic irritation to be regarded as a reliable and essential premonitory symptom of sympathetic ophthalmitis? The answer is in the negative. Sympathetic irritation may last an indefinitely long time without being followed by sympathetic

ophthalmitis. Further, although some sign or signs of sympathetic irritation often do precede the onset of sympathetic ophthalmitis, yet in many cases every such sign is wanting. It is, therefore, wrong to postpone a prophylactic enucleation until sympathetic irritation shows itself.

Treatment.—When sympathetic irritation is caused by an irritable shrunken globe on the opposite side, it can be immediately relieved by removal of the stump. The employment of rest in a dark room and sedative measures, while they may seem to cure, merely lead to disappointment, owing to the almost certain return of the symptoms, when the eye is brought into use again. Moreover, as sympathetic irritation does often precede sympathetic inflammation, it is wise to enucleate the exciting stump in order to assure the safety of the second eye.

Sympathetic Ophthalmitis.

Diagnosis.—The inflammation of the uveal tract in the sympathizing eye has no characteristics which enable us to make the diagnosis "Sympathetic Ophthalmitis," for precisely the same plastic or sero-plastic uveitis, as the case may be, is seen under other conditions; nor is the state of the first eye, taken alone, a certain guide. To arrive at a diagnosis it is necessary to weigh different circumstances, and to take them collectively into consideration—namely:

1. The condition of the exciting eye, and the nature of the injury to, or disease of, that eye.
2. The condition of the sympathizing eye.
3. The interval that has elapsed between the injury to the exciting eye, and the onset of the uveitis in the sympathizing eye.
4. The state of the general system.

1. *The Condition of the Exciting Eye.*—As already stated, perforating injuries, or perforating corneal ulcers, of the first or exciting eye, which are followed by plastic uveitis, are by far the most common, if not the only, causes of sympathetic ophthalmitis. If no inflammation follows the injury, there will be no sympathetic ophthalmitis, even though the wound be a large one. Nor has the position of the wound in the eye any influence in the production of sympathetic ophthalmitis. If uveitis comes on in the injured eye, it is due to infection of the wound by bacteria derived, most usually, we believe, from the foreign body or instrument

which has caused the wound, but possibly from the surface of the eye, or from the atmosphere.

Either a purulent uveitis or a plastic uveitis may result from the injury.

Purulent uveitis of a not very pronounced type (purulent infiltration of the vitreous humor, iritis, hypopyon) is very occasionally followed by sympathetic ophthalmitis, but, and it is a remarkable clinical fact, the marked purulent uveitis, which is called panophthalmitis (p. 264), may be said never to give rise to it. It is not a pyogenic micro-organism which causes sympathetic ophthalmitis, otherwise the latter would be a purulent process; but it is rather some specific and non-pyogenic micro-organism; and it can hardly be doubted that in those rare instances in which sympathetic ophthalmitis has followed on slight purulent uveitis, this specific organism has been present in the injured eye along with the staphylococcus.

When the infection of the injured eye is purulent, the inflammatory reaction comes on within the first thirty-six hours after the injury. The fibrinous or plastic inflammatory reaction, which is so dangerous in relation to sympathetic ophthalmitis, declares itself in the injured eye less quickly and more insidiously. The injection and irritation, the immediate result of the injury, disappear in a few days, only soon to return; then the pupil begins to dilate less well to atropin, the tissue of the iris becomes less distinctly seen, some posterior synechiæ form, and some opacities appear in the vitreous humor. At first there is little or no pain spontaneously or on pressure. Then fresh synechiæ form, the iris stroma becomes more indistinct and discolored, often of a dull greenish or yellowish gray, and the pupil becomes closed. The anterior chamber becomes shallower than normal, and the intra-ocular pressure is diminished. Sight is much affected by the exudation in the pupil and by the opacities in the vitreous, and, in case of detachment of the retina from shrinking of the inflammatory products in the vitreous humor, it may be reduced in a marked degree. There is often pain on pressure of the eyeball, and the latter soon begins to be diminished in size and becomes soft to the touch, while the pressure of the tendons of the orbital muscles on this soft eyeball gives rise to deep furrows on its surface. In short, the eye has now become phthisical, and sight is quite lost. This entire process may be completed in three or four weeks, or it may occupy a considerably longer time.

The danger of sympathetic ophthalmitis supervening on a perforating injury of the first eye commences with the onset of plastic uveitis in the first eye—although the inflammatory process in the second eye does not develop until after a certain interval (*vide infra*)—and is present not only all through the acute process in the injured eye, but also after this has subsided, and when the eye has become shrunken, and even for many years more.

Shrunken eyeballs, as just stated, are liable to cause sympathetic ophthalmitis. Pain on pressure of the ciliary region in them, showing, as it does, the presence of inflammation of the ciliary body, is an important danger-signal; but the absence of pain on pressure is not conclusive of the absence of cyclitis, for the latter may exist to only a slight and yet dangerous degree, or the ciliary body may be detached and out of reach of pressure.

The presence of a foreign body in the interior of the injured eye does not necessarily lead to sympathetic ophthalmitis by the inflammatory reaction which it may cause; for an aseptic foreign body in the eye will cause an active inflammatory reaction; yet this latter, not being of bacterial origin, will not in its turn give rise to sympathetic ophthalmitis. There are, however, few foreign bodies, except atoms of hot metal, which can be guaranteed as aseptic.

Similarly as in accidental perforating injuries, so also the wounds made in the sclerotic or cornea in operations, especially cataract extractions, may be followed by plastic uveitis, which will produce sympathetic ophthalmitis. In consequence of the thorough antiseptic measures now in use, inflammatory processes after cataract extractions are very much less common than they used to be.

Perforations caused by ulcers of the cornea sometimes give rise to plastic uveitis, which may be followed by sympathetic ophthalmitis; but this is an exceedingly rare event, although some iritis is present with almost every severe corneal ulcer, and especially with those which tend to perforate. It is not easy to assign a reason for the rare occurrence of plastic uveitis in these cases, unless it be the hypothetical one that there is some obstacle in the way of the passage of the bacteria into the deeper parts of the eye.

In how far plastic uveitis of the first eye, which is not due to perforating injuries, is capable of being the cause of sympathetic

ophthalmitis is an important question. It was stated in the beginning that such a mode of origin is exceedingly unlikely.

Intra-ocular tumors, which have not yet perforated the sclerotic, especially sarcoma of the chorioid, are said occasionally to set up a uveitis, which leads to sympathetic ophthalmitis. Schirmer [1] is disposed to regard the uveitis in these cases as being caused by an endogenous bacterial infection, and would refer the uveitis in the second eye to the same source, and not, therefore, to be looked upon as a sympathetic affection. But the matter may still be regarded as *sub judice.*

Ruptures of the eyeball from blows, which usually occur in the ciliary region, without rupture of the conjunctiva, come under our notice occasionally (p. 247). They almost invariably run a course free from inflammation, or even irritation of the injured eye, owing to the unbroken conjunctiva which covers the rupture and prevents the access of infecting bacteria, and, consequently, they do not cause sympathetic ophthalmitis. It is probable that in the few cases of this injury in which uveitis in the injured eye and sympathetic ophthalmitis in the second eye appeared, there existed some small opening in the apparently sound conjunctiva.

Cases of gonorrheal ophthalmia have been published in which sympathetic ophthalmitis came on. But these were all cases in which ulceration took place, followed by perforation of the cornea; and, hence, in which infection by bacteria other than the gonococcus was quite possible.

Whether idiopathic or spontaneous inflammation of the uvea, such as those caused by gout, diabetes, syphilis, tubercle, and rheumatism, when they occur in one eye, are capable of giving rise to sympathetic ophthalmitis in the second eye, is a question which cannot at present be quite definitely answered. But it may be taken as almost certain, that the toxic substances or micro-organisms which circulate in the blood or lymphatics, and which cause the uveitis in the first eye, in a similar manner give rise to the uveitis in the second eye when it occurs; and, consequently, that the uveitis in the latter is not to be regarded as sympathetic ophthalmitis.

2. *The Condition of the Sympathizing Eye.*—The diseased process in the second or sympathizing eye, as has already been stated, is, with certain rare exceptions, an inflammation of the uvea, of a plastic or sero-fibrinous type, but never purulent, and

the inflammation almost always begins in the uvea, or, at any rate, is first discovered there as iritis. Occasionally optic neuritis is the first sign of sympathetic ophthalmitis, uveitis coming on subsequently, and, in the rare exceptions referred to above, optic neuritis has been seen as the one and only sympathetic inflammation, the uvea remaining unaffected. It is, however, held by some, that optic neuritis is the first sign in the sympathizing eye in nearly all cases, if only they could be examined before opacities in the vitreous humor, and exudation in the pupil, interfere with an ophthalmoscopic diagnosis.

The appearance of the optic neuritis, or papillitis, as seen in these cases, consists in hyperemia of the disc, without much swelling of it, but with slight wooliness of its margin, the opacity spreading a short distance into the surrounding retina. The veins are distended, and the arteries are normal. The sight is considerably affected, and there is often rather severe headache.

The remedy for sympathetic papillitis is removal of the exciting eye, and a few days after the operation the beneficial effect on the optic nerve inflammation begins to show itself. This prompt response is evidence that the papillitis is not due to an invasion of bacteria, but that here it is rather their toxins which reach the sympathizing eye, and cause the papillitis.

There are no reliable premonitory symptoms of the attack of uveitis in the sympathizing eye. As already stated, in many cases sympathetic irritation does precede the first signs of sympathetic uveitis, but it does not always do so; and when sympathetic irritation does appear, it need not always indicate the approach of sympathetic uveitis.

The early signs of the actual presence of uveitis in the sympathizing eye are: some fine punctate deposits on the posterior surface of the cornea, and these are often the first symptom; slight pericorneal injection; slight opacity of the aqueous humor; some discoloration and indistinctness of the iris; contraction of the pupil, but as yet no synechiæ; some fine opacities in the vitreous humor; and slight loss of sight owing to these changes.

Posterior synechiæ soon begin to form, and the adhesions occur in the most serious cases, not merely between the margin of the pupil and the anterior capsule of the lens, but, after a little while, between the whole of the posterior surface of the iris and the capsule. The exudation which causes this extensive adhesion soon pushes the iris forward, and renders the anterior cham-

ber shallow; but after a time, when the fibrinous exudation begins to shrink, the anterior chamber becomes deep at its periphery, owing to retraction of the iris. The iris gradually becomes more altered, its tissue more dull, discolored, and indistinct, while large vessels form in it. Occasionally, in the anterior chamber a small hypopyon is seen, formed by the fibrin which floats in the aqueous humor, some of which has gravitated.

The intra-ocular tension may become high, often very high, owing to blocking of the angle of the anterior chamber with inflammatory products, and this glaucomatous tension is apt to be attended by great pain. In consequence of the presence of such extensive adhesions, eserin and pilocarpin have no influence on this high tension, and the temptation to perform an iridectomy is very great. Yet there is no more disastrous mistake in ophthalmic practice than any operative interference at this period. Far from doing good, an iridectomy is almost certain to do harm. It is impossible, owing to the disorganized state of the iris and its close adherence to the anterior capsule, to obtain anything like a satisfactory coloboma; and even if the tension be reduced after the operation for a day or two, it very soon becomes as high as ever, in consequence of the rapid filling up of the coloboma by fresh inflammatory products, while the traumatism of the operation only seems to lend additional violence to the inflammation.

The cornea gradually becomes more or less opaque, from derangement of its posterior epithelium by the punctate deposits of fibrin upon it, and the crystalline lens becomes cataractous. After a time the high tension disappears, and gradually low tension comes on. Vision, already very bad, sinks further. The eyeball becomes less in size and very soft to the touch, and phthisis bulbi, with complete blindness, is presented. This entire process may occupy many months, and is often interrupted by short periods of slight improvement in the symptoms.

It is rarely that cases occur in which the sympathetic uveitis comes on with violent pain, chemosis, and swelling of the eyelids, and ends rapidly in phthisis bulbi.

On the other hand, a less severe class of cases is met with, in which the whole posterior surface of the iris does not become adherent to the capsule of the lens, the pupillary margin alone becoming adherent, and these cases may run a comparatively favorable course. Should the pupillary margin become adherent all round in these less severe cases, then increase of tension ensues

from retention of aqueous humor in the posterior portion of the anterior chamber, with consequent bulging forward of the iris, and blocking of the angle of the chamber.

A yet milder, and not uncommon, form of sympathetic uveitis is that in which the signs are: punctate deposits on the posterior surface of the cornea, and increased depth of the anterior chamber, without any iritis. The punctate deposits are at first often so fine as to be undiscoverable unless by aid of a high convex lens behind the sight-hole of the ophthalmoscope, or with a corneal microscope. This form of sympathetic ophthalmitis is termed serous sympathetic uveitis, and its prognosis is favorable. Its one danger consists in the increased intra-ocular tension which is liable to come on, but which should not tempt the surgeon to employ an iridectomy.

More common than this typical serous uveitis are cases in which some fibrin is thrown out, with resulting posterior synechiæ at the pupillary margin. This form of sympathetic ophthalmitis is not attended by much irritation of the eye, nor need vision be much affected. The corneal deposits very gradually increase in number, and consequently vision becomes affected to some extent, and then, if the tension does not increase, the signs and symptoms after a time very slowly abate, and a normal state is re-established. But relapses are liable to occur even after some months, and they may assume the very dangerous fibrinous type. So that, even in these mildest cases, the utmost care in treatment and prognosis is needed.

Removal of the exciting eye does not cure the uveitis in the sympathizing eye (*vide infra*).

3. *The Interval which elapses between the Injury to the Exciting Eye and the Onset of Uveitis in the Sympathizing Eye.*— So far as our present knowledge based on reliable cases goes, the shortest interval is fourteen days, and very few cases with this shortest interval have been reported. The period between the sixth and twelfth week after the injury seems to be the most dangerous. In 170 of the 200 cases collected by the Committee on Sympathetic Ophthalmitis of the Ophthalmological Society [2a] the second eye was attacked within the first year after the injury to the exciting eye. In only 12 of the 200 cases was the interval more than one year, and the longest interval was twenty years.

4. *The State of the General System.*—As the subjects of trau-

matic plastic uveitis in one eye are not immune against plastic uveitis in the other eye due to syphilis, rheumatism, tubercle, diabetes, etc., it is necessary in each case to consider whether the attack in the second eye may not be a symptom of some systemic condition rather than a sympathetic uveitis.

From the foregoing it appears, then, that the diagnosis of sympathetic ophthalmitis depends on circumstantial evidence, namely: (1) As regards the exciting eye: an ectogenic infection. (2) As regards the sympathizing eye: an inflammatory process of a plastic type which attacks all three portions of the uveal tract, is very chronic in its course, often improves for a while, but relapses again. Excision of the first eye does not cure sympathetic uveitis in the second eye; yet, if the sympathetic affection be an optic neuritis, it is rapidly cured by removal of the first eye. (3) As regards the interval between the perforating injury in the first eye, and the appearance of sympathetic ophthalmitis: an interval of at least fourteen days is required. The period between the sixth and twelfth weeks is the most dangerous, and very few cases occur after the first year. (4) As regards the general system: when careful examination of it does not reveal any condition which might be the cause of uveitis in the second eye, the probability of this uveitis being sympathetic is increased.

Prognosis.—The prognosis of sympathetic uveitis is, in general, bad; yet it need not be quite hopeless, for even in some severe cases, and of course more frequently in the less severe cases, the sympathizing eye does recover after prolonged treatment, with a useful amount of vision. But in the rather rare cases which undergo cure the eyes are liable to occasional recurrences of the uveitis, and at least a year should elapse since the last recurrence before a definite end to the diseased process can be said to have been reached.

The prognosis of sympathetic papillitis is quite favorable when once the exciting eye has been removed.

Treatment.—Prophylactic measures are of the first importance. Where the traumatism is so extensive as to make all prospect of saving sight, or even the shape of the eyeball, hopeless, immediate excision of the globe is obviously indicated. In case some prospect of saving sight exists, our attention is claimed in the first instance by the wound in the injured eye, which, in those cases that come for surgical aid sufficiently early, is to be

protected from secondary infection by careful antiseptic cleansing, abscission of any prolapsed portions of the uvea, suturing of the wound in suitable cases, and an antiseptic dressing and bandage.

Should the wound be already infected, then excision of the injured eyeball, or such other operation (evisceration or neurectomy) as in the judgment of the surgeon will secure the second eye from the invasion of bacteria coming from the injured eye, is called for. No temporizing is admissible; even some useful vision being, for the time, retained by the injured eye is not a contra-indication to the operation.

Where sight in the injured eye is lost, it will not be difficult for the surgeon to recommend excision of the eyeball, and even to urge it on the patient; but when some useful sight is still retained, it is not so easy to press this advice, although that should be done. We know, indeed, that in some cases of traumatic uveitis sympathetic uveitis does not supervene; and, provided the first eye has not been too much disorganized by the injury, sight even in it may be ultimately obtained. But, unfortunately, we are unable to foretell whether any given case will run so favorable a course; and to temporize, in the hope that it will do so, involves serious danger to the second eye, and, it may be, ultimate loss of all sight in each eye.

In short, it cannot be doubted that there are cases, but probably not many of them, in which, in the present state of our knowledge, we recommend removal of the injured eye, and where, had we decided to run a fearful risk by allowing it to remain, not only would sight have been restored to it, but no sympathetic ophthalmitis would have come on.

It must be further stated that we cannot feel sure that our removal of the first eye has averted sympathetic ophthalmitis from the second eye, until four weeks after the operation has elapsed. Nearly every ophthalmic surgeon has seen cases in which sympathetic ophthalmitis has appeared subsequently to excision of the first eye, and in which, at the time of the operation, the second eye was perfectly sound. There are well-authenticated cases where sympathetic ophthalmitis appeared as long as four weeks after the enucleation of the injured eye. The assumption is that, in these cases, the infective micro-organisms had already started on their journey to the second eye, and had passed that point in the optic nerve of the injured eye where the

nerve was divided in the operation. Hence, in excising the eyeball, it is very desirable to take away as much as possible of the optic nerve, by severing it far back in the orbit. These cases are deplorable for the patient, and very trying for the surgeon, especially if the outbreak of sympathetic ophthalmitis should occur very soon after—perhaps the day after the operation. Yet where sympathetic ophthalmitis comes on after excision of the first eye, the operation need not be regarded as having been quite useless; for experience shows that the attack of uveitis in the second eye is then usually of a comparatively mild type, and fairly amenable to treatment.

In those cases in which the exciting eye has not yet been removed, and in which sympathetic ophthalmitis in the second eye has commenced, what are our duties? In the first instance, and at the earliest possible moment, removal of the exciting eye, always provided that it be quite and hopelessly blind. The immediate result on the second eye of removal of the first eye under these circumstances is not marked, for the inflammatory process in the former seems to proceed as actively as before. But statistics show that more sympathizing eyes are saved, or partially saved, when the injured eye has been removed soon after the outbreak of sympathetic ophthalmitis than when the injured eye is removed a considerable time after the outbreak, or not at all. Presumably when the source of supply of bacteria is withdrawn, the virulence of the sympathetic disease gradually subsides.

But no exciting eye, which possesses even a slight degree of sight, should be removed when once sympathetic ophthalmitis has appeared, for it might happen that the sympathizing eye would become entirely lost, while the exciting eye might ultimately retain some degree of useful sight. Great caution is therefore required in deciding whether the exciting eye is capable of recovering to a certain extent, and this frequently is a matter of considerable difficulty. Even a partially phthisical eyeball may sometimes ultimately come round sufficiently to gain useful vision. Schirmer[1] lays down the following rule: When sympathetic ophthalmitis has broken out, the exciting eye should not be removed unless it be absolutely blind; or unless, if it still possess perception of sight, it has been for several weeks very soft and reduced in size; or that, by reason of extensive corneal opacity, all hope of restoration to it of form-vision must be abandoned.

If sympathetic ophthalmitis has broken out, either before or after removal of the exciting eye, the treatment and care of the sympathizing eye to promote its recovery must be considered. This consists in the use of atropin, warm fomentations, and subconjunctival injections, which latter are held by some to be very beneficial for the high tension. With these local means is combined a general and prolonged course of mercurialization—mercurial inunctions or calomel internally, or both, care being taken to avoid any severe stomatitis. Diaphoresis, produced two or three times a week by means of pilocarpin (one-eighth to one-third of a grain according to the individual) hypodermically, and hot jars placed in the bed, is a useful adjunct. Salicylate of soda is now used in large doses (as much as one hundred and fifty grains in the day) by some surgeons. The patient is to be confined in one warm, but well-ventilated room, which should be kept almost dark. As this treatment must often be carried on for many weeks or even months, it is a trying one for the patient; but it should be remembered that the issue at stake is a fateful one.

No operation on the iris is to be performed so long as there is the slightest inflammation, or tendency to inflammation, and this rule holds good even if the tension of the eye becomes glaucomatous. Operative interference on the iris here has only the effect of lighting up fresh inflammation; and even if the tension be reduced by an iridectomy, which owing to the diseased and degenerated state of the iris cannot be satisfactorily carried out, it will soon again become high. In six months or a year after every slight sign of inflammation (*e. g.*, injection of the ciliary vessels on insertion of a spring speculum) has passed away—and a longer interval can only be of advantage—it may be allowable to perform an operation on the iris with the object of making an artificial pupil, always provided that there is good prospect of materially improving vision by this means. It must be remembered that, while every operation has its risks, the risks are unusually great in such disorganized eyes; while any loss of sight is felt all the more in a case in which the eye operated on is probably the only one possessing even a little vision. On the other hand, when success crowns an operation in these sad and perplexing cases, the gain is great.

If it be decided not to remove the exciting eye, after sympathetic ophthalmitis has broken out, then its treatment is conducted

on lines quite similar to those above recommended for the sympathizing eye, and the advice concerning operating on it is the same.

Prophylactic, Therapeutic, and Optical Operations used in Sympathetic Ophthalmitis.

Prophylactic Operations.—These are: Enucleation, Evisceration, Mules' Operation, Optico-Ciliary Neurectomy, and Optico-Ciliary Neurotomy.

Enucleation (or Excision).—For the performance of this operation there are two methods.

1. O'Farrall's Method.—The speculum having been inserted, an incision is made in the conjunctiva all round the cornea, and about 6 mm. removed from the latter. The bulbar conjunctiva is separated from the globe freely in all directions with a scissors. With a strabismus hook each orbital muscle is caught up, and its tendon divided close to the sclerotic. The optic nerve is then divided with a strong scissors passed into the orbit from the median side. It is important to divide the optic nerve as far back in the orbit as possible, in order to include infective bacteria which may have traveled thus far on their road to the second eye. Finally the edges of the conjunctiva are drawn together with a few points of suture.

2. The Vienna Method.—The only instruments used in this operation, in addition to the speculum, are a strong straight scissors and a strong toothed forceps. The tendon of the internal rectus at its insertion, with the overlying conjunctiva, is seized in one grasp with the forceps, and so held until the conclusion of the operation. Immediately behind the forceps the tendon is divided with the scissors; and now the forceps is holding merely the stump of the tendon adherent to the globe with the overlying conjunctiva. Through the opening necessarily made at the same time in the conjunctiva one blade of the scissors is passed, and pushed on under the tendon of the inferior rectus muscle, which is then divided along with the overlying conjunctiva. In the same way the superior rectus is divided. The globe is now drawn well forwards and rotated outwards, the scissors passed into the orbit, the optic nerve felt for and divided. With one or two strokes of the scissors the external rectus and the two obliques are divided close to the globe, and the operation is completed. This method is very rapid. It is

not suited to any globe of which the walls are weak (fresh perforating injury, extreme staphyloma, etc.), for a good deal of pressure is exercised on the eyeball during its performance.

Careful aseptic and antiseptic precautions are to be employed in connection with enucleation of the globe. Of these, next to thorough sterilization of the instruments, I regard irrigation of the cavity of the orbit as soon as the eyeball is removed. with a full stream of sublimate solution, 1 in 5000, or of sterilized normal salt solution as the most important. The interior of the orbit is then well covered with xeroform, or other fine antiseptic powder, and an antiseptic dressing is applied with a bandage. The orbit should be similarly dressed every twenty-four hours.

I have never seen the slightest trouble after enucleation of the eyeball; but some cases of meningitis following upon the operation, and which have proved fatal, are reported. There can be no reasonable doubt but that in these instances septic matter made its way along the lymphatics of the optic nerve to the meninges, and that this septic matter was introduced upon the instruments, or escaped, in purulent cases, from the interior of the eyeball. Hence the very great importance of the careful aseptic precautions above indicated.

An artificial eye can usually be inserted after a fortnight, but should not be constantly worn for a month at least, as, until that period elapses, it is liable to cause irritation and conjunctivitis.

Of prophylactic operations for sympathetic ophthalmitis, enucleation is the only one which is regarded by all as thoroughly reliable, when performed in good time.

Evisceration.—For mode of performing this operation, *vide* p. 171. Although, so far as published cases teach us, evisceration, as a preventive measure for sympathetic ophthalmitis, is as effectual as enucleation, yet there is the theoretical objection to it that no portion of the optic nerve is removed; and that, if uveitis has commenced in the injured eye, the bacterial infection may have reached the head of the optic nerve prior to the operation. Consequently, while the source of the infection—the uveal tract of the injured eye—has been removed, what is contained of the infection in the nerve may continue on its way to the second eye, and set up sympathetic ophthalmitis. There can be no such objection to the employment of evisceration where it is performed quite soon after the injury, or at any rate before

uveitis has commenced. The advantage of evisceration over enucleation lies in the better stump provided by it for a prothesis, and the consequent better cosmetic effect.

Mules' Operation.—For the description of this operation, see p. 172. The objections to and advantages of this operation are the same as in evisceration,[2] but it gives a better stump than the latter.

Optic-Ciliary Neurectomy.—An opening is made into the conjunctiva about 3 mm. behind the insertion of the internal rectus muscle; this muscle is laid bare, and two curved blunt strabismus hooks are inserted beneath it. The hooks are drawn in opposite directions, so that one is caught in the angle of insertion of the tendon with a tendency to roll the eye outwards, while the other will draw the muscle forwards out of the orbit. Near the latter hook a catgut thread is passed through muscle and conjunctiva, first from within outwards, and then the opposite way. The muscle is now divided at a distance of at least 5 mm. from its insertion into the sclerotic, and the ends of the catgut thread are tied in a knot. A second thread is passed through the terminal stump of the muscle, and similarly tied in a knot. The wound is now extended both towards the superior and inferior recti muscles, and a small pointed double hook is inserted into the sclerotic far back, in order to draw the globe forwards and outwards. A scissors curved on the flat is inserted alongside the globe, and the optic nerve is cut through as near the optic foramen as possible. The posterior aspect of the globe can now be exposed to view by means of the double hook. The stump of the optic nerve remaining on the eyeball is then cut off near its insertion into the sclerotic, the insertion of the oblique muscles divided, and the whole of the posterior circumference of the sclerotic laid bare by dissection. The eyeball is replaced, the wound closed by means of the catgut threads previously introduced, and, as a precaution against sanguineous exophthalmos, the eyelids are united by three sutures.

The advantage of this operation is that the eyeball is retained.

It is now rarely performed, for it is not regarded as a very reliable protection against sympathetic ophthalmitis, even when done sufficiently early; and there is, moreover, a suspicion that, in some cases, the operation itself, or rather infection taking place during it, has excited sympathetic ophthalmitis.

Optic-Ciliary Neurotomy.—The proceeding here is the same as for resection, except that only the first division of the optic nerve is made. For the prevention of sympathetic ophthalmitis this operation is open to the same objections as resection of the nerve, but in a greater degree. For the relief of sympathetic irritation, either operation may be employed, with at least temporary benefit, in suitable cases.

Therapeutic Operations.— The field for these operations, if it exist at all, is exceedingly limited. Practically the only indication for operative interference, in the active period of sympathetic ophthalmitis, is long-continued high tension; and in the foregoing pages the warning has been repeatedly uttered that any operative meddling with the iris in this period is more apt to aggravate the process than to alleviate it, and that even if tension be relieved by an iridectomy, it soon becomes high again, owing to fresh plastic exudation.

Should it seem imperatively necessary to endeavor to reduce a long-continued high tension, sclerotomy is to be preferred to iridectomy. It may have a beneficial effect, and is not likely to do harm. It can be repeated more than once, should it be deemed necessary.

Paracentesis of the cornea is a measure which can be used as a temporary means of relief for high tension, and it, too, may be repeated.

Optical Operations.— The object of these operations is to provide an artificial pupil in the sympathizing eye, in order to improve, or to restore, vision which is interfered with by closure of the pupil. Similar operations may be indicated occasionally in the exciting eye, in cases where it has not been excised.

The cardinal point to be borne in mind is that these operations may never be performed until six months at least have elapsed—and a longer period is preferable, since all and every tendency to inflammation, or irritation, has subsided. Inattention to this rule will result in a relighting of the inflammation, reclosure of the pupil which may have been made, and a long period of waiting before any further operation can be undertaken.

Moreover, as even under the most favorable conditions, and with the most skillful operation, inflammation may return, or intra-ocular hemorrhage may occur, or the eye may become phthisical, no operation should be done unless the advantage to be gained from it, if successful, promises to be considerable.

The two chief operations, one or other of which may be applicable, are: Iridectomy, and extraction of the clear or cataractous lens; for the lens is often cataractous from interference with its nutrition by the iridocyclitis.

Iridectomy.—It is only exceptionally that iridectomy can be of use in those eyes which have been the subjects of the severer plastic uveitis, resulting in adhesion of the whole posterior surface of the iris to the anterior capsule. In these cases the tissue of the iris has undergone such extreme degeneration that it is impossible to obtain more than mere shreds of the membrane with the forceps, so that a satisfactory coloboma can rarely be made. Or, if a fairly good coloboma be procured, it will probably be found that the pigmentary layer of the iris is left behind; and that this, with organized inflammatory products, lies on the anterior capsule, and obviates any gain that might have been derived from the coloboma.

Iridectomy is indicated rather in those cases where a less severe form of iritis has existed, resulting in a complete ring synechia of the pupillary margin only. Here a wide coloboma may often be made satisfactorily. The iris should be seized with the forceps at about the lesser circle. If seized at the pupillary margin, the intimate adhesion between the latter and the lens capsule may cause injury to the capsule, and consequent traumatic cataract.

Extraction of the Lens.—This is indicated, if, on the formation of a coloboma, the lens be found to be cataractous; in those cases of ring synechiæ where iridectomy has been performed and the coloboma has closed again; and in practically all cases of total adhesion of the iris to the capsule, be the lens clear or opaque. In the former class of cases the ordinary combined method of cataract extraction answers the purpose, or a preliminary iridectomy may be made some weeks previously.

Cases of total adhesion of the iris require a different procedure. A very wide incision is made in the cornea. The knife, having been entered at the outer side, is passed through iris and lens; the latter is then delivered as completely as possible, and out of the membrane composed of degenerated iris, retro-iridic connective tissue, and capsule, a V-shaped piece is cut with the forceps-scissors.

Discission.—This operation was employed by the late Mr. George Critchett with much success in some cases where cataract

was the main obstruction to sight. A discission needle is passed, by a boring motion, through the lenticular capsule; another needle is then passed in close to the first, and by separating one point from the other a rent is made in the center. This is followed generally by the escape into the anterior chamber of a small quantity of cheesy lens matter, which becomes gradually absorbed, and in the course of some weeks the capsule closes again. The operation has to be repeated several times before a clear pupil is obtained, care being taken that all irritation from the previous operation has subsided before another is undertaken. The danger in this method would be irritation and high tension from swelling of the lens in a disorganized eye, and I do not know that the method has been much used by other surgeons.

Pathogenesis.— The pathogenesis of sympathetic ophthalmitis is still surrounded by obscurity. An old view was that irritation of the ciliary nerves in the exciting eye caused an irritation of the ciliary nerves in the opposite eye, which irritation, in its turn, gave rise to inflammation in this second eye. But modern pathology teaches that there is no such thing as reflex inflammation.

There are, nowadays, two chief theories, each of which relies on bacterial infection of the sympathizing eye by specific organisms of the process coming from the exciting eye. But these theories differ from each other in respect of the path by which they would respectively make these organisms pass from the first to the second eye. Neither school, it must be stated, has been able, by present-day methods, to identify the bacteria which set up the inflammation. Yet this cannot be held to be a supreme objection to either theory, for there are many bacterial diseases—*e. g.*, syphilis, rheumatism, smallpox—the micro-organisms of which our present methods have failed to demonstrate.

According to the theory which numbers most adherents, the path which the infective organisms take in passing from the exciting to the sympathizing eye is along the lymphatics of the optic nerve, or its sheaths, of the first eye, then by the optic commissure, and down the optic nerve of the sympathizing eye—in short, by direct continuity, as erysipelas extends over the skin. This view, which was first put forward by Leber,[3] is maintained by many trustworthy authorities. Deutschmann,[4] indeed, semed to have definitely proved the case by experiments on animals. He injected the staphylococcus pyogenes into one eye,

and thus produced a uveitis in the other eye. Not only this, but he discovered the coccus in the optic nerve sheath of the exciting eye, and also in that of the sympathizing eye. Many others have repeated his experiments, but no one has obtained the same results. Moreover, the inflammatory process, which the staphylococcus pyogenes produces, is purulent, while sympathetic ophthalmitis is a sero-plastic or fibrinous process. The theory has yet to be shown to be correct by experimental research, or, what would be preferable, by pathological demonstration in the human subject. For the present it relies on clinical evidence, and it is, moreover, in accord with accepted pathological principles.

The other view, propounded by Schmidt-Rimpler,[5] explains the infection of the sympathizing eye as a manifestation of a bacterial infection of the general system from the exciting eye. The germs, it is suggested, enter the general circulation, and travel through all parts of the body, but find a suitable home only in the fellow-eye.

Injuries.

Injuries of the Iris.—*Punctured Wounds* of the cornea, or of the corneo-scleral margin, frequently implicate the iris, but rarely do so without also injuring the crystalline lens or ciliary body, on which then the chief interest centers, as being the organs from which serious reaction is most likely to emanate. A small, simple incised wound of the iris is not of great importance, for inflammatory reaction is not common, and any extravasation of blood at the seat of the iris wound, or into the anterior chamber (hyphemia) becomes absorbed, while, in most cases, the functions of the iris will probably not be affected nor sight endangered. Nevertheless, as iritis does sometimes occur, it is desirable to use such measures as are calculated to prevent it, such as atropin, a bandage, and rest of the eye and general system. Even extensive wounds of the iris are not often, as such, associated with serious danger to the eye, although the loss of continuity in the iris never closes up. Where, for instance, the iris is cut in its entire width from ciliary margin to pupillary margin, the permanent result is a wide coloboma, the margins of which may be adherent to the corneal wound. When the iris is prolapsed in the corneal wound, every effort should be made to reduce it completely; and, if this cannot be affected, it is necessary to abscise

the prolapsed portion. Incarceration of the iris in the corneal cicatrix may lead to secondary glaucoma, cystoid cicatrix, secondary septic infection of the iris, etc.

Foreign Bodies of small size, such as bits of steel or iron, may perforate the cornea and fasten in the iris, the puncture in the cornea closing rapidly, and possibly no aqueous humor being lost. It is necessary always to remove such a foreign body without delay, although for some time it may cause no reaction. An incision should be made with a Graefe's knife at the margin of the cornea corresponding to the position of the foreign body, and the portion of iris containing the foreign body is then removed with forceps and scissors.

Blows on the Eye are apt to cause, in addition to hemorrhage into the anterior chamber from the iris or from the canal of Schlemm, one of several remarkable lesions of the iris, namely:




Fig. 92.

1. *Iridodialysis* *—*i. e.,* separation of the iris from its attachment to the ciliary body, which is usually accompanied by considerable hyphemia. As much as one-half of the circumference of the iris may be involved in the lesion, or the latter may be so small as to be diagnosed only by the presence of the resulting small fresh hemorrhage near the ciliary margin of the iris; or, after this has become absorbed, by aid of light transmitted to the eye by the ophthalmoscope, and then not only the physiological pupil, but also the minute marginal traumatic pupil, will be illuminated. It is rarely that there is more than one dialysis. In certain degrees of the detachment, by reason of the sphincter of the iris having lost its fixed point, it becomes stretched in a straight line (Fig. 92) at the part corresponding to the dialysis, and assumes a D-shape, or if the detachment be more extensive, the pupil becomes kidney-shaped, or the de-

* ἶρις ; διάλυσις, *a separating.*

tached portion may entirely cover the pupil. The detached portion, too, may be turned on itself (anteflexion of the iris), the uveal surface being to the front. The functions of the eye after such an injury, even when extensive, may be but little disturbed, or there may be monocular diplopia.

It is stated that iridodialysis does not become re-attached; but I have observed one case in which a very minute one was healed, and there is one other such case recorded. The lengthened use of atropin is the most likely way in which to promote such a result, but it can only be hoped for if the iridodialysis be not extensive and the case be seen early.

Iridodialysis does not increase in extent in the course of time, nor lead to further mischief in the eye.

2. *Retroflexion of the Iris.*—The whole, or more commonly a portion, of the iris in its entire width becomes folded back on the ciliary processes, giving the appearance of a very dilated pupil or of a coloboma produced by a wide and peripheral iridectomy. In a true coloboma the ciliary processes would be easily seen, but not so in retroflection, for the processes, being covered by the retroflexed iris, present a smooth surface. A slight dislocation of the lens in the direction away from the iris lesion is often observed. Retroflexion of the iris cannot be cured, but useful vision is retained, if the injury be uncomplicated.

3. *Rupture of the Sphincter Iridis.*—There are not many cases of this lesion recorded; although, according to Hirschberg, in all cases of permanent traumatic mydriasis the margin of the pupil is torn. My observations do not agree with his, nor do I think that rupture of the sphincter would be sufficient to account for the traumatic mydriasis which is usually associated with it. There may be but one rupture, or there may be a number of small ruptures distributed round the pupil. They show themselves as small triangular gaps in the pupillary margin, their bases directed towards the latter. This condition is also incurable, and some permanent disturbance of vision due to the mydriasis results.

4. *Dehiscence of the Iris* between the pupillary and ciliary margins. This is a slit-like rupture of the iris which runs in a radial direction through the whole width of the iris with the exception of the sphincter. The diagnosis sometimes cannot be made with certainty until, after a few days, the blood-clot covering the dehiscence is absorbed. The opening may be caused

to close by the use of a myotic, which, by contracting the sphincter, brings the edges of the dehiscence together.

5. *Traumatic Aniridia.*—The whole iris is torn from its ciliary insertion, and may be found lying in the anterior chamber or under the conjunctiva, having in the latter case passed through a rent at the corneo-scleral margin. Not only does the anterior chamber contain blood, but the vitreous humor is often infiltrated with hemorrhage. When the extravasated blood has become sufficiently absorbed, the absence of the iris will be noted, and in many instances the ciliary processes will be visible. If these latter are visible the diagnosis "aniridia" can be definitely made, but cases do occur in which, notwithstanding the absence of the iris, the ciliary processes are not visible, owing probably to changes in the processes which cause them to shrink. Such cases then are difficult to distinguish from retroflexion of the iris, but the importance of the diagnosis is not great.

6. *Traumatic Mydriasis, and Myosis.*—Of these, mydriasis is the more common. The dilatation is of medium degree, and the pupil is usually of irregular shape—oval, pear-shaped, or more dilated at one part than elsewhere—and contracts but slightly, or not at all, to light. Paralysis of accommodation usually accompanies traumatic paralysis of the sphincter iridis. The mydriasis is probably the result of concussion of the delicate nerve-endings in the sphincter of the iris. (See above, under Rupture of the Sphincter Iridis.)

With traumatic myosis there is apt to be spasm of accommodation, which may produce apparent myopia. The prognosis is fairly good. Traumatic mydriasis may recover after a long interval, but in most instances it remains as a permanent defect, with some defect of vision due to it, and to the paralysis of accommodation.

Treatment.—For mydriasis, protection spectacles, galvanism, and eserin. For myosis, atropin.

Injuries of the Ciliary Body.—*Punctured Wounds, and Foreign Bodies,* perforating the sclerotic at a distance of about 5 mm. around the cornea, are almost certain to implicate the ciliary body. If there be no prolapse of the ciliary body, nor any foreign body in the interior of the eye, the sclerotic wound may heal by aid of a bandage without further ill results. If a prolapse of the ciliary body or iris be present, it is to be abcised, with careful antiseptic measures; and if the sclerotic wound be

large, it may be thought desirable to unite its margins with sutures.

Wounds of the ciliary body are apt to cause cyclitis, especially if the former be incarcerated in the sclerotic wound in healing, for the incarcerated portion is liable to become infected.

Injuries of the Chorioid.

Small Foreign Bodies may pierce the sclerotic, or the cornea and lens, and lodge in the chorioid, and, if favorably situated, can be detected with the ophthalmoscope, but in any case by the Röntgen rays if of metal. They require operative removal by the magnet, if of steel or iron; or if the foreign body cannot be extracted, the eyeball must be removed, to avert sympathetic ophthalmitis.

Incised Wounds of the sclerotic very frequently involve the chorioid (see p. 248).

Rupture of the Chorioid is often produced by blows on the eye, and is seen with the ophthalmoscope as a whitish-yellow (the color of the sclerotic) crescent some two or three papilla-diameters in length, and one papilla-diameter or so distant from the optic entrance, the concavity of the crescent being directed towards the papilla. Immediately after the accident extravasated blood sometimes prevents a view of the rupture. Some chorioiditis may result; but, when this passes away, good vision is frequently restored and maintained, provided detachment of the retina does not ultimately supervene from cicatricial contraction at the seat of the rupture. On the other hand, a scotoma in the field may be produced, and if the rupture be in the region of the macula lutea, serious loss of sight may be caused.

Treatment.—Careful protection of the eye, and abstinence from use of it, with dry cupping at the temple for three weeks, or until it may be assumed that all inflammatory tendency has subsided.

Blows upon the eye may cause *Extravasation of Blood in the Chorioid.* If small, these extravasations do not extend beyond the chorioid. But in the case of copious extravasation the hemorrhage is poured out from the chorioidal vessels between that coat and the sclerotic, lifting and bulging forward the chorioid; or between the chorioid and retina, giving rise to a detachment of the latter; and if the retina gives way, the blood is poured out

into the vitreous humor. Should there be no vitreous humor opacity, the extravasations in the chorioid can be seen with the ophthalmoscope as somewhat indistinct (owing to resulting opacity in the overlying retina) small red spots, or large round red spots, darker in the center than at the margin. That these hemorrhages are in the chorioid can be recognized from the fact that they lie behind the retinal vessels. The hemorrhages become slowly absorbed, and after a time, provided the retina has not burst, useful vision may be restored.

Treatment.—Complete rest in bed. Atropin. Bandage.

New Growths.

New Growths of the Iris.— *Cysts.*—These vary from a very small size to that which would fill the anterior chamber. They may have either serous or solid contents. The serous kind was said to result always from a trauma causing an anterior synechia, or otherwise shutting off a fold of the iris, which became distended into a cyst by accumulation of aqueous humor. A case, however, which was not preceded by a trauma has come under my notice. The cysts with solid contents (epidermoid elements) are believed to have their origin in an eyelash or morsel of epidermis, which may have made its way into the anterior chamber by occasion of a perforating corneal wound. All these cysts are sources of serious danger to the eye (irido-chorioditis, glaucoma, etc.), and, it has been stated, may even be the cause of sympathetic ophthalmitis, and hence their removal is called for. This can be effected without much difficulty if the tumor be small, but if it have attained a large size the attempt may be unsuccessful. A long incision should be made in the corneo-scleral margin, and the cyst, along with the portion of iris to which it is attached, drawn out and cut off.

Granuloma is the name given to a benign neoplasm of the iris, of which the structure resembles granulation tissue. Clinically it is a small pale tumor, or there may be several such tumors, which gradually grow to fill the anterior chamber, rupture the cornea, and finally induce phthisis bulbi. It is held by some that these growths depend on a syphilitic taint, and by others that they are tubercular.

Tubercle (*Tubercular Iritis*).—This disease occurs generally in children or young adults, who may or may not present evidence

of general tuberculosis, such as enlarged or caseating glands, or diseases of joints, etc. It is met with in two forms—viz., disseminated or miliary tubercle, and conglomerate or solitary tubercle.

Miliary Tuberculosis of the iris is a relatively mild form, which presents the clinical appearances of a chronic iritis, sometimes with punctate corneal deposits; but it is chiefly characterized by the formation of a number of grayish or cinnamon-colored, semi-translucent nodules on the surface of the iris and at the angle of the anterior chamber. Occasionally they are not very numerous. The disease may either run its course, and finally cause shrinking of the eye from plastic iridocyclitis, or it may subside even without treatment. It is to this form of iritis that Leber [6] has given the name attenuated tuberculosis of the iris; but it is not due to any attenuation of the virus, for inoculation in the anterior chamber of a rabbit's eye of an excised portion of such an iris produces severe local and general tuberculosis.

Solitary tubercle may be accompanied by a few smaller growths, but it generally begins as a single yellowish-white tumor, often without iritis, which gradually increases in size until it may fill the anterior chamber. It finally involves the cornea, which it perforates, forming a fungating mass, and this, subsequently, breaks down, leaving only a small stump in the socket. Microscopically both varieties present the usual structure of tubercle, but bacilli are very difficult to detect in either of them.

Treatment in the miliary forms consists of the usual local and constitutional means. Internal administration of creasote has been recommended. If the disease continue to progress, enucleation may be necessary.

Should a solitary tubercle be seen in an early stage, it may be removed by an iridectomy; but if the disease have advanced too far, or the iridectomy have failed, the eye must be extirpated. Operative treatment will depend very much on the view which the surgeon takes of the origin of the disease. It has until recently been generally believed that, while tubercle of the chorioid was a disease secondary to tuberculosis elsewhere, tubercle of the iris was a primary affection, and as such necessitated immediate enucleation of the eye, in order to prevent it from becoming a source of general infection. The impression, however, seems to be growing that tubercle of the iris is also a

secondary affection, and that, consequently, enucleation is not always indicated, with the object of averting tubercular disease of the general system. Of course in those cases, which are not uncommon, where both eyes are affected, the question of enucleation cannot be entertained.

Sarcoma.—Of the uveal tract, the iris is the portion most rarely affected with primary sarcoma. It arises usually from a congenital pigmented nevus of the iris, and is commonly a melano-sarcoma; but leuko-sarcoma has also been recorded. As the tumor increases in size, it fills the anterior chamber, and grows backwards into the ciliary body and into the canal of Schlemm. It is not usual for the tumor to become extra-ocular by growing through at the corneo-scleral margin, and in this respect it is unlike tubercle of the iris. Irritation or inflammatory symptoms are not often present, and secondary glaucoma does not come on until a later stage, when the growth has filled the anterior chamber, or involved the ciliary body extensively.

Treatment.—Enucleation of the eye should be advised as soon as the diagnosis of sarcoma of the iris has been made. There is naturally a desire on the part of the surgeon, when the tumor is small, to save the eye, which probably has full vision, by excising the portion of iris in which the growth is seated, and there are some cases on record in which this was done, and where no recurrence of the tumor took place. But in adopting this conservative method, the surgeon accepts grave responsibility; for it is not possible to determine clinically whether the sarcomatous growth is truly, or only apparently, confined to the limited region of the iris, where it can be seen. Even in the early stages of many cases of sarcoma of the iris the neoplasm invades the ligamentum pectinatum, the canal of Schlemm, or the ciliary body; so that, although the iris tumor be thoroughly removed, the growth reappears in the eye before long, while in the meantime risk of infection of the general system has been run.

Ophthalmia Nodosa is a very rare affection, of which about a dozen cases have been recorded. It is caused by the irritating secretion contained in the hollow hairs of certain caterpillars. In nearly all cases there was a history of caterpillars having accidentally come into forcible contact with the eye. The disease, which is very chronic, is characterized by the presence of small hard nodules in the conjunctiva and iris, and may lead to severe iridocylitis. The diagnosis is confirmed by the presence of

brownish hairs; or by the examination of an excised nodule, which shows the hair in sections as a yellow oval body with a central cavity.

New Growths of the Ciliary Body.—*Sarcoma* of the ciliary body is generally pigmented, and often passes unobserved, until it attains considerable size as a brown mass, which was at first concealed from view by the iris. Occasionally it is first noticed when it makes its appearance at the angle of the anterior chamber. It usually also grows backwards into the chorioid, and runs the same course as sarcoma of the chorioid. Removal of the eyeball should be urged, and may for a time be declined by the patient, as sight is but slightly affected in the early stages.

Myosarcoma originating in the ciliary muscle has been observed a few times.

Carcinoma.—Primary carcinoma of the ciliary body is an extremely rare disease. Its occurrence in this situation is easily explained if the ciliary body, which secretes the intra-ocular fluid, has a glandular structure; and, from the researches of Collins,[7] there seems to be every reason to believe that it does contain tubular glands.

Secondary carcinoma may occur in the ciliary body as in the chorioid (*vide infra*), but is very rare.

New Growths of the Chorioid.—*Sarcoma* is by far the most common neoplasm of the chorioid, and the chorioid is the most common seat of ocular sarcoma. It is seen at all times of life, but most frequently between the ages of forty and sixty. Both melano-sarcoma and leuko-sarcoma occur, and may originate in any part of the chorioid.

If seen in a very early stage, it is easily recognized from its projecting over the general surface of the fundus, the retina lying closely applied to it; but, unless it be in the region of the macula lutea, it may not cause any serious disturbance of vision, and hence may not at that period be brought under the notice of the surgeon. The diagnosis from detachment of the chorioid at this stage is made by the presence in the latter condition of the characteristic chorioidal vessels, and by the peculiar color of the chorioid. Detachment of the chorioid, too, is much rarer than sarcoma.

The new growth soon gives rise to detachment of the retina by reason of serous exudation from the chorioid; and this may be accompanied by opacity in the vitreous humor, which contributes

to render the diagnosis with the ophthalmoscope difficult or impossible. If the detachment be shallow and the retina translucent, the tumor may still sometimes be seen through the subretinal fluid by aid of strong illumination; and even direct sunlight may be employed in some such cases. Owing to the great, and often sudden, defect of vision which comes on in this stage, we very commonly see these cases then for the first time. The history of the case may aid us; while the absence of the more usual causes of detachment of the retina should make us suspicious of an intra-ocular tumor, and the fundus should be carefully examined, with dilated pupil, in all such cases.

Soon the intra-ocular tension increases. This makes the diagnosis again more easy in many cases, for the combination of detached retina and increased tension exists only with intra-ocular tumors. The increased tension may come on very slowly, and without ciliary neuralgia; or more rapidly, and with all the signs and symptoms of acute glaucoma. Still, if the case come now under observation for the first time, the diagnosis may be by no means easy, should the refracting media be opaque (as always in acute glaucoma), and consequently the detachment of the retina concealed from view. Here, again, the history of the case is all we have to depend on, especially the fact of the patient having noticed a defect at one side of his field of vision previous to the onset of glaucoma.

In the next stage of the growth it perforates the cornea or sclerotic, and, increasing rapidly in size, although still covered with conjunctiva, it pushes the eyeball to one side, the upper lid being stretched tightly over the whole. On raising the lid the tumor is seen as a bluish-gray mass with irregular surface. The conjunctiva is now soon perforated, and the surface of the tumor becomes ulcerated, with a foul-smelling discharge and occasional hemorrhages. The tumor gradually invades the surrounding skin and the bones of the orbit, and by extending through the sphenoidal fissure and optic foramen reaches the base of the brain.

Another, and less usual, course of chorioidal sarcoma is that in which, without first perforating the cornea or sclerotic, the tumor sets up irido-cyclitis, leading to phthisis bulbi.

It is usually upon the neighboring tissues of the eyeball becoming involved that secondary growths begin to form in other organs, the one most prone to be affected being the liver. The

lungs, stomach, peritoneum, spleen, and kidneys may all be attacked.

Chorioidal sarcoma is almost always primary, but it has been seen a few times as a metastatic disease.

The entire progress of such a growth varies considerably. It may occupy but a few months, or it may extend over many years.

Cases in which sarcoma of the chorioid was present in shrunken eyeballs have given rise to the view that such eyeballs are prone to develop sarcoma. While it is quite possible that sarcoma may develop in a shrunken eyeball, it is tolerably certain that in the majority of the cases in which both diseases are present the sarcoma is the primary disease.[8]

Carcinoma.—This is extremely rare, and the cases of it on record were all of metastatic origin, the primary disease being in the breast. It is not possible to distinguish chorioidal sarcoma from chorioidal carcinoma by the ophthalmoscope.

Tubercle is sometimes seen in cases of acute miliary tuberculosis as round, slightly prominent, pale yellowish spots, of sizes varying from 0.5 to 2.5 mm. in diameter, situated always in the neighborhood of the optic papilla and macula lutea, and unaccompanied by pigmentary or other chorioidal changes. There may be but one of these tubercles, or there may be many of them. They occur as a rule in a late stage of the general disease, but have occasionally been noted long before its appearance. In obscure cases of the general disease, the ophthalmoscope has sometimes rendered valuable diagnostic aid by discovering these minute tubercles in the chorioid.

Very rarely a tubercular tumor grows in the chorioid in cases of general chronic tuberculosis, and attains a large size, the growth destroying the eye similarly as sarcoma or carcinoma. In young children it may be impossible to diagnose between a tubercular tumor of the chorioid and a glioma of the retina (chap. xv.). Yet, as in either case enucleation is indicated, the diagnosis is not of much practical importance.

Other, but rare, forms of tumor of the chorioid are:

Sarcoma Carcinomatosum and *Osteo-Sarcoma.*

Treatment.—So long as, in cases of sarcoma and carcinoma, the tumor is wholly intra-ocular, enucleation of the eyeball should be performed, and may be done with fair hopes of saving the patient's life, if the disease be primary. When the orbital tissues

have become involved, extirpation of all the contents of the orbit, and even, if necessary, removal of portions of its bony walls, ought to be undertaken, should the general health permit, in order to rid the patient of his loathsome disease; although the probable presence of secondary growths elsewhere renders but small the prospect of saving the patient's life.

Cases of miliary chorioidal tubercle do not call for direct treatment. In cases of tubercular tumor the question of removal of the eyeball must depend upon the general state of the patient; but if it seem probable that life will be prolonged until after the ocular growth would become extra-ocular, removal of the eye should be recommended.

Other Diseases of the Chorioid.—*Posterior Sclero-Chorioiditis, or Posterior Staphyloma.*—This condition is described in connection with myopia (p. 46), which is its almost constant cause.

Detachment of the Chorioid.—As the result of copious loss of vitreous, during operations, or from injury, detachment of the chorioid is not uncommon, but it does not require to be specially diagnosed in these instances, and, therefore, it is not important to consider it further here.

But idiopathic detachment of the chorioid, although extremely rare, is of importance, as forming a well-defined diseased condition in itself.

The ophthalmoscopic appearances here are apt to be taken at first sight for a simple detachment of the retina, or for leuko-sarcoma; but on closer inspection the chorioidal stroma is observed to lie immediately behind the detached retina, and its vessels, etc., are seen in the upright image by aid of the same lens as are the retinal vessels. The chorioid is not everywhere detached, but is separated from the sclerotic in several different places, and these detachments are seen in the form of apparently solid hemispherical protuberances rising abruptly from the fundus into the vitreous humor. In other places the chorioid is in contact with the sclerotic, although in some of these positions there may be detachment of the retina alone. The vitreous humor is more or less opaque. Needless to say, vision is greatly lowered or quite destroyed.

It is probable that a chronic chorioido-retinitis has been an antecedent condition in all of these cases. Indeed, there often are signs of old retinitis present, such as perivasculitis and con-

nective tissue striation; and in one case a retinitis was actually observed long before the detachment of the chorioid came on. Adhesions between the chorioid and sclerotic are formed in consequence of this inflammation; and then inflammatory exudation takes place behind the chorioid, and separates it from the sclerotic, where it happens not to be adherent to the latter.

The process ends either in phthisis bulbi, in consequence of vascular changes and disturbances of nutrition, or in cure of a certain degree, in so far as by absorption of some of the exudation, and by alteration of the remainder of it into connective tissue, a return of the chorioid and retina to their normal position is rendered possible; but even then restoration of sight, with tissues so disorganized, cannot be looked for.

Treatment hitherto seems to have been of no avail. Probably active mercurialization might afford the best chance of doing good, should a case come under notice.

Fuchs has pointed out that detachment of the chorioid occurs in a good many cases of cataract extraction some days after the operation, and also in some cases of iridectomy. It can often be found with the ophthalmoscope, and even sometimes with the oblique illumination, in those cases of cataract extraction in which the anterior chamber has not formed, or in which, having formed, it has become empty again. It is mainly after iridectomy for chronic simple glaucoma that chorioidal detachment has been noticed. The probable explanation is that a slight aperture of communication has been made between the anterior chamber and the subchorioidal space, through which the aqueous humor passes behind the chorioid. With the re-establishment of the anterior chamber, the chorioidal detachment goes back, and the prognosis is in all cases good as regards vision.

Central Senile Areolar Atrophy of the Chorioid.—This is not a very rare disease, and presents the appearance of a white patch, often of considerable extent, at and around the macular region. In some cases a hemorrhage in the chorioid and posterior layers of the retina form the starting-point of the disease. The retinal functions always suffer much; for an absolute central scotoma is produced, which renders reading and writing impossible, although locomotion is not much impeded, as the periphery of the field remains intact. The discovery of the presence of this disease, after a cataract has been successfully removed, is sometimes a source

of intense disappointment both to patient and surgeon, which cannot be guarded against unless it has been noted while the cataract was still incipient.

Treatment is of no avail, but absolute rest of the eyes from all attempts at near work, and the use of dark protection spectacles are important, so that, at the least, the advance of the disease may not be encouraged.

Malformations.

Malformations of the Iris.—*Heterophthalmos* (*ἕτερος, different; ὀφθαλμός*).—This term indicates that the color of the iris in one eye is different from that in the other.

Corectopia (*κόρη, the pupil; ἔκτοπος, out of position*), or malposition of the pupil. The pupil sometimes occupies a position farther from the center of the iris than normally.

Polycoria (*πολύς, many; κόρη, the pupil*).—Where there is more than one pupil. The supernumerary pupil may be separated by only a small bridge from the normal pupil, or it may be situated very near the periphery of the iris. In neither case has it a special sphincter.

Persistent Pupillary Membrane appears in the form of very fine threads stretched across the pupil. They cannot be mistaken for posterior synechiæ, as they are attached to the anterior surface of the iris some distance from the margin of the pupil. They do not interfere with the motions of the pupil, nor with vision.

Coloboma (*κολοβός, maimed*) and *Irideremia* (*ἶρις, ἐρημία, want of*).—These two defects have been shown by Treacher Collins [9] to be due to a similar cause—in short, that they are different degrees of one and the same condition. Before the iris is formed in the fetus there exists, between the posterior surface of the cornea and the anterior capsule of the lens, the anterior portion of the fibro-vascular sheath. This receives its blood-supply partly from the ciliary arteries, and partly from those in the posterior fibro-vascular sheath, prolonged round the sides of the lens to join it. The cornea, anterior fibro-vascular sheath, and lens lie in close contact with each other. The iris is developed by growing forwards from the margin of the anterior chamber, and in doing so has to insinuate itself between the cornea and anterior fibro-vascular sheath on the one side and the lens on the other, pushing the prolongation from

the posterior fibro-vascular sheath in front of it. The anterior fibro-vascular sheath subsequently becomes the pupillary membrane, of which portions sometimes persist (see above). If we suppose some abnormal adhesion to occur between the cornea, anterior fibro-vascular sheath, and lens-capsule, or some delay in their separation at the whole circumference of the future anterior chamber, we can understand how a mechanical obstruction to any growth of the iris forwards would be introduced, resulting in complete absence of the iris, or irideremia. If the obstruction be confined to a portion only of the anterior chamber, the corresponding portion only of the iris will be prevented from growing forwards, and the result will be one or more congenital colobomata. Irideremia may be complete or partial. In the latter case it may be the inner circle only which is wanting, giving the pupil the appearance of dilatation with atropin. Where the entire iris is absent the ciliary processes can be seen all round. The condition may be binocular. The patients suffer chiefly from dazzling by light, from which either protection or stenopeic spectacles afford some relief.

Malformations of the Chorioid.— *Coloboma.*—This is a solution of continuity occurring always in the lower part of the chorioid, and usually associated with a similar defect in the iris. It may commence at the optic papilla, and involve the ciliary body also, and even the crystalline lens may have a corresponding notch; or it may not extend so far in either direction. The condition is recognized ophthalmoscopically by the white patch, due to exposure of the sclerotic where the chorioid is deficient. Sometimes the retina is absent over the defect in the chorioid, a circumstance which may be ascertained by the arrangement of the retinal vessels; but, even if it be present, its functions at that place are wanting, and a defect in the field of vision exists. Central vision is often normal.

Albinismus, or the want of pigment in the chorioid and iris. This is usually accompanied by defective pigmentation of the hair of the body. The iris has a pink appearance, due to reflection of light from its blood-vessels, and from those of the chorioid, and with the ophthalmoscope the latter vessels can be seen down to their finest branchings. The light which enters the eye, not being partially absorbed by pigment, causes the patient much dazzling, and high degrees of the condition are usually accompanied by nystagmus. In childhood the albinismus and at-

tendant symptoms are more marked than later on, when some degree of pigmentation usually takes place.

Much advantage may be derived in many of these cases by



FIG. 93.



FIG. 94.



FIG. 95.




FIG. 96.

the use of stenopeic spectacles, at least for near work. Any defect of refraction should be carefully corrected, in order to give the patients the best possible use of their eyes.

Operations on the Iris.

Iridectomy.—This is performed for optical purposes, as in zonular cataract, corneal opacities, or closed pupil; to reduce abnormally high intra-ocular tension, in primary and secondary glaucoma; and for the removal of tumors or foreign bodies in the iris.

The instruments required are a spring speculum; a fixation forceps, with spring catch (Fig. 97); a lance-shaped iridectomy




Fig. 97.

knife (keratome) (Fig. 93), or a Graefe's cataract knife; a bent iris forceps (Fig. 94), or a Tyrrell's hook (Fig. 95); an iris scissors curved on the flat (Fig. 96), or a de Wecker's forceps-scissors; and a small spatula.

The width of the coloboma depends a good deal on the length of the corneal incision, for it cannot be wider than the incision is long. *Its depth* depends on the proximity of this incision to the corneo-scleral margin. If a wide and very peripheral coloboma



Fig. 98.



Fig. 99.



Fig. 100.

be desired, the incision must be long, and must lie actually in the corneo-scleral margin; the iris forceps being then introduced, a portion of the iris corresponding to the length of the incision may be seized, drawn out, and cut off, the blades of the scissors being applied parallel, and close to the incision, and by this means a coloboma, as at Fig. 98, is produced. An incision somewhat inside the corneal margin will give a pupil, as in Fig. 99. A narrow coloboma (Fig. 100) is obtained by a short corneal incision, which may be more or less peripheral as circumstances

require; by taking up as little as possible of the iris in the forceps, or by using a Tyrrell's hook, instead of an iris forceps, for catching and drawing out the iris; and by applying the blades of the scissors at right angles to the incision in the corneal margin.

In glaucoma a wide and very peripheral coloboma is required. For optical purposes a narrow iridectomy is required, because with a wide coloboma the diffusion of light may be very troublesome to the patient.

The best position for an iridectomy for glaucoma is in the upper quadrant of the iris, as there the subsequent dazzling by light and the disfigurement are least. But the position, by preference, for an optical pupil is below and to the inside, being that most nearly in the direction of the axis of vision. If, however, this position be occupied by a corneal opacity, the coloboma should be made directly downwards; or, if that place be ineligible, then downwards and outwards, or directly downwards, or directly inwards. The upward positions are not satisfactory for optical pupils, owing to the overhanging of the upper lid; yet it often happens that we have no other choice.

In the Performance of an Iridectomy the eye should be fixed with a forceps at a position on the same meridian as that in which the coloboma is to lie, but at the opposite side of the cornea, and close to the latter. The point of the lance-shaped knife is then to be entered almost perpendicularly to the surface of the cornea, and made to penetrate the latter. As soon as the point of the blade has entered the anterior chamber the handle of the knife is lowered, and the blade is passed on into the anterior chamber in a plane parallel to the surface of the iris, until the incision has attained the required length. The handle of the knife is now lowered still more, so as to bring the point of the blade almost in contact with the posterior surface of the cornea, in order to prevent any injury to the lens in the next motion. The knife is then very slowly withdrawn from the anterior chamber. At the same time the aqueous humor flows off, and the crystalline lens and iris come forwards. The fixation forceps is now taken over by the assistant, and the bent iris-forceps is passed closed into the anterior chamber, its points directed towards the posterior surface of the cornea, so as to avoid engaging them in the iris. When the pupillary margin has been reached, the forceps is opened as widely as the corneal incision will permit,

and the corresponding portion of the iris is seized and drawn out to its full extent through the corneal incision. With the scissors held in the other hand the exposed bit of iris is snipped off quite close to the corneal incision. Care should now be taken that the angles of the coloboma do not remain in the wound; and if they are seen to do so, they must be reposed by stroking the region of the incision with a hard rubber spoon, or by pushing them into their places gently with the spatula.

Iridotomy.—For description and use of this operation see chap xiii.

References.

[1] "Von Graefe's Archiv," xxxviii. 4. p. 93. "Centralblatt für Augenheilkunde," 1899, p. 40. "Graefe-Saemisch Handbook" (2d Ed.), vi.

[2] "Trans. Ophth. Society, U. K.," vi., p. 170.

[2a] Report of Committee of the Ophthal. Society of the United Kingdom on the "Relative Value of Simple Excision of the Eyeball, and the Operations which have been Substituted for it."—"Trans. Ophth. Soc., U. K.," 1898, p. 233.

[3] "Von Graefe's Archiv," xxvii. 1. p. 331.

[4] *Ibidem*, xxx. 3. p. 277.

[5] *Ibidem*, xxxviii. 1. p. 199.

[6] "Bericht der ophthalmologischen Gesellschaft." Heidelberg, 1891, p. 44.

[7] "Trans. Ophth. Society, U. K.," xi. p. 55.

[8] "Von Graefe's Archiv," xliv. 1. p. 164.

[9] "Trans. Ophth. Society, U. K.," xiii. p. 128.

CHAPTER XI.

THE MOTIONS OF THE PUPIL IN HEALTH AND DISEASE.

The Size of the Pupil in Health depends chiefly on the intensity of the light to which the eye is exposed, contracting when light falls into the eye, and dilating in the shade. However defective vision may be, if quantitative perception of light remains, the reaction of the pupil as a rule takes place.

There is no absolute *Standard for the Physiological Size of the Pupil.* The latter varies in different healthy individuals, being in general smaller in elderly people than in youthful subjects; for with increasing age the energy of the sympathetic—the dilating nerve of the iris—is reduced, while there is sclerosis of the walls of the vessels of the iris and rigidity of its stroma. Hypermetropic eyes are apt to have small pupils, owing to the constant effort of accommodation; while in myopia, for the converse reason, the pupils are wide. The diameter of the pupil when the accommodation is at rest has been found to vary between 2.44 and 5.82 mm., giving an average diameter of 4.14 mm.

Tange's [1] very careful investigations showed that the size of the pupil varies much in individuals; that women have larger pupils than men; that the physiological dimension of the pupil diminishes with the age; that the pupil is smaller in hypermetropia than in emmetropia; that the size of the pupil becomes less as the hypermetropia increases; that the physiological dimension of the pupil in myopia up to the twentieth year is greater than in emmetropia; that with increasing age the differences in the dimensions of the physiological pupil in the different kinds of refraction becomes less, until beyond the fortieth year they are insignificant; that the color of the iris has no influence on the size of the pupil.

Contraction of the Pupil.—Contraction to light is a reflex motion, the optic nerve being the afferent nerve, and the third nerve the efferent nerve innervating the sphincter pupillæ. It has been shown by a high authority [2] that there are special afferent fibers in the optic nerve for the pupil-reflex, distinct from those for vision, and that it is possible to distinguish with the microscope these two kinds of nerve fibers from each other.

The anatomical investigations of Meynert [3] have shown that between the corpora quadrigemina and the center for the third nerve run communicating fibers (2 and 2, Fig. 101), which probably enable this reflex to take place. Owing to the semi-decussation of the fibers in the optic chiasma, the stimulus of light, when applied to one eye alone, passes up

each tract with equal power to the corpora quadrigemina, and thence, by Meynert's fibers, to the nucleus of the third nerve (or rather to that portion of it which acts as a special center for the sphincter pupillæ), and from that point down the myotic, or short ciliary, branches of this nerve to each ciliary ganglion, the ciliary nerves, and each sphincter iridis, causing as active a contraction of the pupil in the non-illuminated eye (consensual contraction) as in its fellow. It is probable, however,[4] that, in addition to this method of bringing about consensual contraction of the pupil, there is a communication, direct or indirect, between the nuclei of the third nerve of each side capable of effecting it. In no other way can the fact be explained that consensual contraction of the pupil is maintained in cases of homonymous hemianopsia. If, for instance (Fig. 101), there be a lesion of the right tractus opticus giving rise to left hemianopsia, the nucleus of the left third nerve alone can be primarily




FIG. 101.—3*N*. Nucleus of third nerve. 1. Connection between nuclei of third nerves. 2. Meynert's fibers. *Q*. Corpora quadrigemina. *C*. Chiasma. *O*. Optic nerve. *P*. Myotic fibers of third nerve. *L*. Seat of Lesion. Arrows show path of impulse in lesion of right tract at *L*.

stimulated; but, as both pupils act, a communication between the nuclei of the third nerves must exist. Merkel[5] believes that there is a direct anastomosis between these nuclei.

But it must be stated that there is a good deal of divergence of opinion as to the path by which the pupil reflex is brought about. Bechterew is of opinion that the centripetal pupillary fibers pass uncrossed from the chiasma directly to the gray matter surrounding the third ventricle, and thence backward to the pupillary nucleus of the oculo-motor nerve of their own sides respectively. Gudden made experiments which seemed

to him to prove that the corpora quadrigemina had nothing to do with this path, and ascribed to the external geniculate body the part usually assigned to the corpora quadrigemina. Mendel's experiments [6] would lead to the view that it is the ganglion habenulæ which is the center for the pupillary reflex in animals, and in this he is largely supported by Darkschewitz, who holds that the pupillary fibers from the optic tract pass both into the pineal gland and the ganglion habenulæ. According to Mendel, the reflex path would be: Optic nerve, optic tract, to the ganglion habenulæ of the same side, thence by the posterior commissure to the nucleus of the third nerve, and thence to the ciliary nerves.

Henschen [7] inclines to think that the centripetal pupil-fibers are to be found in connection with the mesial root of the optic tract, and he excludes altogether the external geniculate body from their path. They would appear, he says, to pierce the posterior section of the cerebral peduncle, and, with the other fibers of the mesial tract, to enter into relations rather with the internal geniculate body.

Bernheimer's [8] researches prove to him that the pupil-fibers pass into the external geniculate body, from thence under the internal geniculate body, and into the substance of the anterior quadrigeminal body, and thence to the third-nerve nucleus in the aqueduct of Sylvius. Crossed and uncrossed fibers are contained in this course, and hence each sphincter nucleus is related to uncrossed fibers from the eye on the same side, and crossed fibers from the eye on the opposite side. He holds also that there is a more or less direct connection of the sphincter nuclei with each other, although he has not succeeded in demonstrating it anatomically.

The reflex mobility of the pupil to light is tested most commonly for the purpose of deciding the existence or otherwise of posterior synechiæ. The next most common object of the test, and the one with which we are here concerned, is to determine the sensitiveness to light of the retina or of the visual center. It affords generally a sufficient test of the presence or absence of quantitative perception of light; but it must be remembered that the latter function may be wanting in certain diseased states, and yet the pupil-reflex take place; or the pupil-reflex may be wanting, and still perception of light be present. The test is best performed in diffuse daylight, with the patient's face directed towards the window, a distant object being looked at, and the eye which is not under examination being carefully excluded from the light. The surgeon then, having observed the size of the pupil to be examined, excludes the eye from light with his hand for some moments. On removing the excluding hand, a normally reacting pupil will be found to have become dilated; and this dilatation, after an interval of about half a second, will be observed to give way to an extreme contraction, which is maintained only for a moment, and is then succeeded by a moderate dilatation, and the pupil then again contracts somewhat, and so on, until, after some further minute oscillations, it comes to a standstill. The explanation for this phenomenon—which is termed hippus—is that each contraction of the pupil, by diminishing the supply of light to the retina, contains in itself the cause of the succeeding dilatation; and, for the converse reason, each dilatation sets a-going the succeeding contraction, until at last equilib-

rium is attained. A comparison between the maximum of dilatation and maximum of contraction, along with the promptness and rapidity with which the contraction takes place, enables the observer to form an estimate of the activity of the pupil-reflex. In performing this test it is important that the patient's gaze should be fixed all through on a distant object—hence, unless where a mere trace of perception of light remains, the test used with the artificial light is not so reliable as that with daylight—so that the pupil-contraction which is associated with convergence or accommodation (*vide infra*) may not vitiate the experiment. The danger of a vitiation of the experiment by the reflex dilatation from the skin (*vide infra*) caused by the excluding hand is insignificant in practice. The consensual reflex of the pupil, as well as the direct, should always be tested—one eye being alternately excluded and exposed, the motions of the pupil of the other eye are observed and compared with those of its fellow.

In examining the pupils we have also to decide whether they are of equal size; and, in order to avoid error through posterior synechiæ, the comparison should be made, with both eyes open, successively in two very different brightnesses of light. Under normal conditions equality in size of the pupils will exist, not only with both eyes open, but also if one eye be shaded; for the normal consensual pupil-reflex is equal to the direct reflex. If the pupils be found of different sizes, the least movable one is usually the pathological pupil, but this is a question often difficult to decide. Finally, it should be noted whether the direct pupil-reflex is similar in all respects in each eye.

In addition to the stimulus of light, the pupil-contracting center is excited by, or simultaneously with, the effort of accommodation for near vision. The object of this contraction is to cut off rays falling on the peripheral portions of the lens, which latter are not curved in the change for accommodation to the same degree as is the center of the lens. This contraction, however, is much more intimately connected with convergence of the visual lines than with the effort of accommodation. It has been shown [9] that the contraction increases with the effort of accommodation, but not proportionately to the distance of the fixation point from the eye; and [10] that the pupils do not contract if accommodation be effected without convergence, but that in convergence without accommodation contraction is observed. It has also been found that the contraction is proportional to the degree of convergence, and that in myopes of high degree contraction of the pupil takes place at the other side of the far point, where, of course, the accommodation does not come into play.

Aubert [11] thinks there is probably a common center for the three actions, convergence, accommodation, and pupil-contraction—a view supported by Priestley Smith [12]; and Hensen and Völckers [13] have found that in dogs, in the posterior part of the floor of the third ventricle, the nuclear centers for the branches to the ciliary muscle, the sphincter pupillæ, and the rectus internus occur in close succession, and they think that this region may be regarded as the center assumed by Aubert. The existence of such a nuclear center has been placed beyond controversy by Eales' case [14] of paralysis of convergence and accommodation, and of the

associated pupillary contraction. These three motions, then, are not dependent on each other, but are coeffects of one and the same cause—*i. e.*, a stimulus applied to the center for convergence, accommodation, and pupil-contraction.

In examining the mobility of the pupils in a given case the contraction on convergence should not be omitted. If the patient be blind of both eyes, the observation can be made by calling on him to direct his eyes towards his own hand at about 12 inches' distance. If both accommodative contraction and light reflex are wanting, a lesion in the course of the centrifugal pupil fibers is indicated; while, if the light reaction alone is wanting, the lesion is in the course of the centripetal fibers.

Dilatation of the Pupil.—The most reliable investigations [15] have distinctly proved that there is no such muscle as the dilator pupillæ. The dilatation of the pupil is in all probability largely the result of an inhibitory action of the sympathetic, a view maintained also by Gaskell [16] and Jessop.[17] The posterior limiting membrane of the iris is its only structure which is not thrown into folds when the pupil dilates (Fuchs); and therefore there can be little doubt that it takes an active part in dilating the pupil, probably by reason of its elasticity. Yet, inasmuch as when the pupil is dilated from paralysis of the third nerve a further dilatation can be produced by atropin, it is probable that some other, as yet unascertained, dilating power resides in the iris. The mydriatic, or long ciliary, nerves originating (Hensen and Völckers) in the front part of the floor of the aqueduct of Sylvius, pass to a region in the lower cervical and upper dorsal portion of the cord, called by Budge [18] the ciliospinal center, and from thence pass out with the two first dorsal nerves, and by way of the rami communicate to the sympathetic in the neck, and thence to the cavernous plexus, Gasserian ganglion, ophthalmic division of the fifth nerve, nasal branch of this division, ganglionic branch of this nerve, ciliary ganglion, there joined by more branches from the cavernous plexus, and from thence by the short ciliary nerves reach the eye.

The dilating nerve fibers are probably of twofold nature, muscular and vasomotor. The experiments of Grünhagen,[19] Salkowski,[20] Donders and Hamer,[21] Stellwag,[22] and J. Arlt, jun.,[23] indicate this; and that the center for each kind of fiber is different, though both are situated in the medulla oblongata, and their fibers probably run the same course to the eye. The center for the muscular fibers is called the oculo-pupillary center. That the vasomotor fibers have a decided and independent influence in dilating the pupil has been shown by Rouget,[24] Schoeler,[25] and others. It is not certain what the mechanism of this influence may be, but it probably consists in a diminution in volume of the iris from anemia caused by contraction of the muscular coat of the vessels.

Langley and Anderson [26] find that stimulation of the cervical sympathetic causes dilatation of the pupil before the vessels of the iris contract, and that stimulation of the portion of the iris can produce a displacement of the pupil towards the side stimulated without relaxation of the sphincter. They assume, therefore, that there must be some radial contractile substance in the iris, but in what form they do not say.

While light is the only stimulus capable of bringing about a reflex contraction of the pupil, the pupil-dilating center reacts to every sensitive stimulus—*e. g.*, the prick of a pin or a pinch on the neck, galvanism applied to the leg,[27] the tickling of a sensitive place in the region of the fifth nerve on the face,[28] etc., and Westphal[29] observed dilatation on shouting loudly into the ear of a person under chloroform. Schiff and Foa[30] found that in curarized dogs and cats a dilatation took place on the application of every stimulus, not necessarily painful, applied to the nerves of common sensation in any part of the body. Indeed, it is not necessary in the human subject that the stimulation should produce any sensation, for stimulation of the skin of the affected side in hemianesthesia, as also in sleep and in coma, will find response in dilatation of the pupil. The center for this reflex is probably in the medulla oblongata,[31] but, inasmuch as it takes place if the cervical sympathetic be divided,[32] it is evident that all the dilating fibers do not run to the eye by way of the cervical sympathetic. Schiff[33] thinks it probable that the Gasserian ganglion receives pupil-dilating fibers from the sympathetic traversing the cavum tympani.

Some psychical emotions produce dilatation of the pupil. The pupils of a cat in anger dilate, and those of a frightened child. In sleep, or when under the complete influence of an anesthetic, the pupils are contracted, for then all psychical and sensitive stimuli are reduced to a minimum. Facts authorize the conclusion that the medium dilatation of the pupil in the healthy state depends chiefly on the intensity of these stimuli, habitually transmitted through the sympathetic. If in any individual they be slight, his pupil is contracted; if intense, it is dilated. Arndt[34] asserts that in delicate, nervous, excitable people the pupils are often much, and habitually, dilated.

In addition to those already mentioned, there are causes for the dilatation of the pupil which can hardly be referred to simple reflex action, but which seem to be, like the contraction of the pupil on convergence of the visual lines, associated with those of other centers in the medulla oblongata, especially with those for respiration and uterine action. With every *deep* inspiration or expiration a considerable pupillary dilatation takes place, not identical with that slight dilatation occurring on each ordinary inspiration and depending on variation of blood pressure, but due[35] to simultaneous stimulation of the respiratory and pupil-dilating centers by retention of carbonic acid gas in the blood. Raehlmann and Witowski[36] have observed marked dilatation at the beginning of each labor pain, to be explained as an associated action of the neighboring centers for uterine movements and pupil-dilatation.

Besides the normal pupillary motions described in the foregoing, and visible for the most part to the naked eye of the observer, there is a phenomenon of pupillary motion which is discoverable only by aid of a corneal microscope or loup, consisting in perpetual, but very minute and irregular, fluctuations in size of the pupil. This hippus has been aptly termed by Laqueur[37] the Unrest of the Pupil, and is due to the ever-varying sensitive and psychical reflexes which are thus constantly manifesting their influences on the pupil.

The Fifth Nerve has been held by some to have an influence over the

motions of the iris similar to that of the sympathetic. This is, according to Leeser, a mistaken view [38]; the effect on the pupil following section of the fifth within the cranium being due to paralysis of the sympathetic fibers contained in it, and not to the lesion of the proper fibers of the fifth nerve. But Spallita and Consiglio [39] found, after removal of the superior cervical ganglion of the sympathetic, and when sufficient time for degeneration had been allowed to elapse, that stimulation of the fifth nerve caused myosis. Others, [40] again, have ascribed to the fifth nerve a direct influence over the contraction of the pupil; but this is to be regarded as a reflex action merely, Merkel indeed having demonstrated [41] the existence of a direct fibrillar connection between the centers of the fifth and third nerves.

Action of the Mydriatics on the Pupil. *Atropin.*—Inasmuch as a maximum mydriasis can only result from paralysis of the pupillary branches of the third nerve, combined with excitation of the pupillary branches of the sympathetic, and as atropin effects such a mydriasis, it is evident that it acts in the way indicated on these nerves.[42] A. von Graefe proved [43] that the aqueous humor of an eye into which atropin has been instilled acts as a mydriatic when applied to another eye. *Duboisin, Hyoscyamin, Scopolamin,* and *Daturin* act similarly to atropin. *Cocain* mydriasis seems [44] to be induced merely by a local irritation of the endings of the sympathetic in the iris, both of the vaso-constrictor fibers and of the pupil-inhibitory fibers. Strychnin and curare are not, strictly speaking, mydriatics, as they only indirectly affect the pupil; the mydriasis observed in poisoning by these drugs being, according to Schiff [45] and others, the result of the retention in the blood of carbonic acid gas.

Action of the Myotics on the Pupil. *Eserin* (or *Physostigmin*).—This drug is in all respects a complete antagonist of atropin,[46] paralyzing the peripheral endings of the sympathetic in the iris, and stimulating the endings of the branch of the third nerve in the sphincter pupillæ. *Pilocarpin* and *Muscarin* act similarly, but not with the same energy. *Nicotin* applied to the eye is found to act like eserin.[47] Morphium has an antagonistic effect to atropin, both as regards the pupil and the general nervous system, and is employed in cases of poisoning by atropin (*vide* p. 257).

Chloroform in the first or excitation stage of anesthesia, according to the investigations of Westphal,[48] Budin,[49] and Hirschberg,[50] stimulates the pupil-dilating center, and in the second stage gradually reduces the excitability of this center, until, finally, it is completely paralyzed, so that no form of stimulation causes any dilatation. Following on this is a still further contraction to a pinhole pupil, due to stimulation of the pupil-contracting center. Should the inhalation of the anesthetic be continued longer, a dilatation of the pupil, often sudden, takes place, and this indicates paralysis of the nuclear pupil-contracting center, and the most serious consequences for the life of the patient.

The Size of the Pupil in Disease.—*Myosis* may be caused by a diseased process irritating the pupil-contracting center or nerve-fibers (the Irritation Myosis of Leeser), or by one causing paralysis of the pupil-dilating center or nerve-fibers (the Paralytic Myosis of Leeser), or by a com-

bination of both. Either cause alone would produce a medium myosis; a combination of the two would give a maximum myosis.

Irritation Myosis, according to Leeser, is not usually increased by the stimulus of light, nor on convergence of the visual axes, nor does it diminish in the shade. Mydriatics dilate such a pupil widely; myotics contract it *ad maximum*. In paralytic myosis the pupil reacts well to light and on convergence, but does not dilate on application of sensitive or psychical stimuli, or with co-ordinated motions. Mydriatics dilate such a pupil only partially, while myotics contract it *ad maximum*. In maximum myosis every reaction is wanting, strong mydriatics alone producing a medium dilatation.

Irritation myosis is found—*a*. In the early stages, at least, of all inflammatory affections of the brain and its meninges; in simple, tubercular, and cerebrospinal meningitis. When in these diseases the medium myosis gives place to mydriasis, the change is a serious prognostic sign,[51] indicating the stage of depression with paralysis of the third nerve. *b*. In cerebral apoplexy the pupil is at first contracted, according to Berthold,[52] who points out that this contraction is a diagnostic sign between apoplexy and embolism, in which latter the pupil is unaltered. *c*. In the early stages of intracranial tumors situated at the origin of the third nerve or in its course. *d*. At the beginning of an hysterical or of an epileptic attack.[53] *e*. In tobacco amblyopia,[54] probably from stimulation of the pupil-contracting center by the nicotin. *f*. In persons following certain trades, as the result of long-maintained effort of accommodation [55] (watchmakers, jewelers, etc.), the pupil-contracting center being subject to an almost constant stimulus. g. As a reflex action in ciliary neurosis; consequently, in many diseased conditions of those parts of the eye supplied by the fifth nerve.

Paralytic myosis occurs in spinal lesions above the dorsal vertebræ—*e. g.*, injuries and inflammations, especially of the chronic form. The contracted pupil occurring in gray degeneration of the posterior columns of the spinal cord has been long known as Spinal Myosis. In the simple form of this myosis the pupil has but a medium contraction, and reacts both to light and on convergence. This condition is found in the early stages alone, when the disease has attacked merely the cilio-spinal center, or higher up, as far as the medulla oblongata; later on, when Meynert's fibers become engaged, we have the Argyll Robertson pupil. The very minute pupil, often seen in tabes dorsalis, is probably due to secondary contraction of the sphincter pupillæ.[56]

Argyll Robertson was the first to point out [57] that in tabes dorsalis the pupil, although contracted, and responding to light by further contraction but slightly, or not at all, does become more contracted on convergence of the visual axes (or accommodation). He explained this phenomenon as being due to paralysis of the ciliospinal nerves, which he therefore regarded as the nerves supplying the sphincter iridis. But Raehlmann points out [58] that the myosis and the motor phenomenon are not directly connected; for it sometimes happens that pupils which do not react to light and do contract on convergence are not habitually contracted, and may even be somewhat dilated. The two symptoms are no doubt often present together in tabes. The myosis is a sign, and an important

one, of disease of the posterior columns; while the defective reaction to light with retained contraction on convergence indicates disease at some distance from the spinal cord. It has been held by some that the seat of the disease, causing the Argyll Robertson pupil, is in Meynert's fibers (2 and 2, Fig. 100) connecting the corpora quadrigemina and the third nerve nuclei. But, as has been pointed out by Bevan Lewis,[59] while this explanation would answer were all instances of this symptom binocular, it cannot be the true one when, as we know, the symptom is sometimes unilateral; for the internuclear path (1, Fig. 101) between the nuclei of the two third nerves must exist in order to enable the consensual action of the pupils which takes place in lesions of one optic tract to be brought about. In the same way lesion of Meynert's fibers on one side would still permit of the pupillary reaction to light of each pupil. Bevan Lewis therefore concludes that the Argyll Robertson pupil is due to a nuclear lesion. Disease in Meynert's fibers (as also disease of the optic nerve) may be in direct connection with disease of the cord, Stilling having found[60] fibers passing directly from the optic tract into the crus cerebri.

Some authorities regard myosis as one of the earliest symptoms of tabes, while others do not. Raehlmann also thinks that, perception of light being present, if the pupils do not react to light, while they do contract on convergence, the symptom is usually one of serious central disease.

Paralytic myosis is also found in general paralysis of the insane. In acute mania the pupil is usually much dilated; and when this mydriasis is changed for myosis, approaching general paralysis may be prognosticated.[61] Myosis following on irritation mydriasis is also found in myelitis of the cervical portion of the cord. In bulbar paralysis, if paralytic myosis occurs, the disease is probably complicated with progressive muscular atrophy or with sclerosis of the brain and spinal cord.[62]

Hirschler states[63] that he has frequently noticed a contracted pupil in alcoholic amblyopia, due, probably, to an affection of the medulla oblongata, possibly fatty degeneration. Myosis may also be due to paralysis of the cervical sympathetic, resulting from injury, from pressure of an aneurysm of the carotid, innominate, or aorta, or from pressure of enlarged lymphatic glands. In apoplexy of the pons Varolii myosis is present, but it is not yet certain whether it is an irritation myosis[64] or a paralytic myosis.[65]

Mydriasis may be caused by a diseased process giving rise to irritation of the pupil-dilating center or fibers, or by paralysis of the pupil-contracting center or fibers.

The former is termed Irritation (or Spasmodic) Mydriasis, and, according to Leeser, is characterized by a moderately dilated pupil, contracting somewhat to light and on convergence, but not dilating on sensitive or psychical stimuli; easily dilated *ad maximum* by mydriatics, but with difficulty contracted *ad maximum* by myotics. The latter is called Paralytic Mydriasis, and in it there is a moderately dilated pupil, reacting to sensitive and psychical stimuli. The reaction to light and on convergence varies according to the seat of the lesion. If the lesion lie between the

iris and the pupil-contracting center, the direct and consensual reaction to light is wanting, as also the associated motion on convergence of the visual lines. But if the lesion lie between the retina and the pupil-contracting center, the direct contraction to light is wanting, while the consensual contraction and that on convergence are retained.[66] In either case the pupil can be dilated *ad maximum* by mydriatics, but not contracted more than to medium size by myotics.

Irritation of the pupil-dilating center and paralysis of the pupil-contracting center existing simultaneously give rise to maximum mydriasis. In it there is absolute immobility to stimuli of all kinds, except to strong myotics, which may bring the pupil back to the normal size.

Irritation Mydriasis occurs—*a*. In hyperemia of the cervical portion of the spinal cord and in spinal meningitis. *b*. In the early stages of new growths in the cervical portion of the cord. *c*. In cases of intracranial tumor and other diseases causing high intracranial pressure, according to Raehlmann, although Leeser points out that these may also give rise to paralytic mydriasis. *d*. In the spinal irritation of chlorotic or anemic people after severe illness, etc. *e*. As a premonitory sign of tabes dorsalis. *f*. in cases of intestinal worms, owing to the stimulation of the sensitive nerves of the bowel; and sometimes in other forms of intestinal irritation. *g*. In psychical excitement—*e. g.*, acute mania, melancholia, progressive paralysis of the insane (often then unilateral, with myosis in the other eye).

Unilateral mydriasis occurring at short intervals, now in one eye and now in the other, is, according to Von Graefe,[67] a premonitory sign of mental derangement. Von Graefe observes madness, in the form of *manie des grandeurs,* to come on some months after the occurrence of this symptom.

Paralytic Mydriasis (Iridoplegia) may be due either to a paralysis of the pupil-contracting center or as the result of the stimulus not being conducted from the retina to that center. It may be found under the following circumstances: *a*. Sometimes in progressive paralysis, where at first there was myosis. *b*. In various diseased processes at the base of the brain affecting the nuclear center of the third nerve. *c*. In a late stage of thrombosis of the cavernous sinus.[68] *d*. In orbital processes which cause pressure on the ciliary nerves. *e*. In glaucoma. *f*. In cases of intra-ocular tumors which have attained a certain size.

In paralytic mydriasis, due to non-transmission of the stimulus of light to a healthy pupil-contracting center and nerves, contraction of the pupil will take place only on convergence of the visual lines. The same condition of pupil will be found if the lesion lie in the course of Meynert's fibers, although vision may be normal. If the lesion lie in the center of vision, or in the course of the fibers connecting this center with the corpora quadrigemina, although absolute amaurosis exists, the reaction of the pupil to light will be perfect. Paralytic mydriasis, due to non-conduction of light stimulus, is found in most cases of optic atrophy.

Bevan Lewis has pointed out[69] that the reflex dilatation on stimulating the skin is wanting in cases of general paralysis and of epilepsy to the extent of about 36 per cent. in women and 43 per cent. in men.

Damsch has noticed [70] a marked increase of the hippus of the pupil in certain diseased states—namely, multiple sclerosis, acute meningitis, apoplectic attacks followed by secondary tremor and spasms of the paralyzed muscles, and in neurasthenia. He is inclined to liken the hippus in these cases to the increase of the tendon reflexes, while immobility of the pupil would be the homologue of loss of tendon-reflex. Yet he does not think an exclusively reflex origin, for the exaggerated hippus can be adopted in these cases, as it continues to an abnormal degree even when all reflex irritation is avoided; and consequently he concludes that an increase of the physiological hippus must be included as a cause.

Förster [71] finds that in tabes dorsalis the oscillations of the pupil diminish in intensity, while the rhythm remains unaltered; but that in progressive paralysis the rhythm is lost. When the pupil has lost its power of reaction to light, the hippus still continues for a while.

References.

[1] "Archiv f. Augenheilkunde," xlvi. 1. p. 49.

[2] B. von Gudden, "Stitzungsber. d. Münch. Ges. f. Morphol. u. Physiol.," 1886, i. p. 1.

[3] Vom Gehirn der Säugethiere, "Stricker's Handbuch." Leipzig, 1870.

[4] Leeser, "Die pupillarbewegung in Physiologischer und Pathologischer Beziehung," p. 14. Wiesbaden, 1881.

[5] "Graefe-Saemisch Handbuch," vol. i.

[6] "Neurolog. Centralbl.," 1890, p. 184.

[7] "Klin. u. Anatom. Beiträge z. Patholog. d. Gehirns," iii. p. 111.

[8] "Von Graefe's Archiv," xlvii.," 1899, pt. 1. p. 1. and "Graefe-Saemisch Handbuch" (2d. ed.), vol. i.

[9] Adamük and Woinow, "Archiv für Ophthalmologie," xvii. pt. 1.

[10] E. H. Weber, "De Motu iridis." Lipsiæ, 1851.

[11] "Graefe und Saemisch Handbuch," ii. p. 669.

[12] "Ophthal. Hosp. Rep.," vol. ix. p. 32.

[13] "Arch. f. Ophthal.," xxiv. pt. p. 23.

[14] "Trans. Ophthal. Soc.," January 10, 1884.

[15] Schwalbe, "Handbuch der Sinnesorgane"; Eversbusch, "Bericht d. Ophthal. Gesellsch.," 1884; Fuchs, "Graefe's Archiv," xxxi. pt. 3. p. 39; Jessop. "Proceed. Roy. Soc.," 1886, p. 478.

[16] "Jour of Phys.," viii. 1. p. 38.

[17] "Proceed. Roy. Soc.," 1886, p. 484.

[18] "Ueber die Bewegungen der Iris," 1855.

[19] "Zeitschrift f. rat. Med.," xxviii., "Archiv. f. d. Gesam. Physiol.," Bd. liii.

[20] *Ibid.*, xxix. p. 167.

[21] "Nederl. Tijdschr. v. Geneesk.," 1864.

[22] "Ueber Atropin," "All. Wiener Med. Zeitung," 1872, p. 146.

[23] "Archiv für Ophthal.," xv. i.

[24] "Comptes rendus et Mém. de la Soc. de Biologie," 1856.

[25] "Experimentelle Beiträge zur Irisbegung": Inaug. Dis., Dorpat, 1869.

[26] "Journal of Physiology," 1892, vol. xiii. No. 6.
[27] Arndt "Griesenger's Archiv f. Psych.," ii.
[28] Hecker, "Tageblatt der 45 Versam. deutscher Naturforscher in Leipsig," 1872.
[29] "Virchow's Archiv," xxvii. p. 409.
[30] "La pupilla come estesiometro," "L'Imparziale," 1874.
[31] Salkowski, *loc. cit.*
[32] Vulpian, "Archiv de physiol., etc., de Brown-Séquard," Janvier, 1874.
[33] "Untersuchungen zur Naturlehre," x 1867, p. 423.
[34] "Archiv f. Psychiatrie," ii. p. 589.
[35] Schiff, *loc. cit.*
[36] "Archiv f. Physiologie," 1878, p. 110.
[37] "Klin. Monatsbl. f. Augenheilk.," December, 1887.
[38] Leeser, *loc cit.* pp. 46-48.
[39] "Archivio de Ottalmologia," vol. i. 183.
[40] Grünhagen, "Berl. Klin. Wochenschr.," 1866, No. 24; Rogow, "Zeitschr. f. rat. Med.," vol. xxix. p. 289.
[41] "Graefe und Saemisch's Handbuch," i. p. 140.
[42] Hermann, "Lehrb. der exp. Toxicologie."
[43] "Archiv f. Ophthal.," i. pt. 1. p. 462, footnote.
[44] Jessop, "Proceed. Roy. Soc.," p. 441, 1885.
[45] "Pflüger's Archiv," 1871, p. 229.
[46] Harnack, "Arch f. exp. Pathol.," ii. p. 307; A. Weber, "Archiv f. Ophthal.," xxii. pt. 2.
[47] Rogow, "Zeitschrift f. rat. Med.," xxix. p. 1; Schur, "Zeitschrift f. rat. Med.," xxxi. p. 402.
[48] "Virchow's Archiv," xxvii. p. 409.
[49] "Gazette des Hôpitaux," 1874, p. 910.
[50] "Berl. Klin. Wochenschr.," 1876, p. 652.
[51] Leeser, *loc. cit.*, p. 82.
[52] "Bery. Klin. Wochenschr.," 1869, No. 39.
[53] Wecker, "Graefe und Saemisch's Handbuch," iv.
[54] Hirschler, "Arch. f. Ophthal.," xvii. pt. 1.
[55] Seiffert, "Allgem. Zeitschrift für Psychiatrie," x. p. 544.
[56] Hempel, "Archiv f. Ophthal.," xxii. pt. 1.
[57] "Edin. Med. Journal," xiv., 1869, p. 669, and xv., 1870, p. 487.
[58] *Loc cit.*, p. 7.
[59] "Brit. Med. Journal," April 25 and May 2, 1896.
[60] "Beilageheft zu Zehender's Monatsblätter," xvii. pp. 203-207.
[61] Seiffert, *loc. cit.*
[62] Leeser, *loc. cit.*, p. 94.
[63] "Archiv f. Ophthal.," xvii. pt. 1. p. 229.
[64] Larcher, "Pathol. de la protub. Annulaire," deux. tirage, p. 54.
[65] Jüdel, "Berl. Klin. Wochensch.," 1872, No. 24.
[66] Heddæus ("Archiv f. Ophthal." xxvii. p. 38) and Turner ("Royal London Ophthal. Hosp. Rep.." December, 1892) assume that the sphincter derives its nerve fibers from two nuclear centers—viz., from the special sphincter center, and also from the center for convergence (or accommodation).

[67] "Archiv f. Ophthal.," iii. pt. 3. p. 350.
[68] Knapp, "Archiv f. Ophthal.," xiv. pt. 1. p. 220.
[69] "Brit. Med. Jour.," April 25 and May 3, 1896.
[70] Neurolog. Centralbl.," 1890, p. 258; also Zeminski, abstract in "Annales d'Oculist.," March, 1894, p. 239.
[71] Versammlung deutscher Naturf. u. Aerzte, Nürnberg, 1893 ("Deutsch. Med. Wochenschr.").

The three following tables, showing the action of the various
been prepared for this

TABLE

THE MYDRIATICS

	ATROPIN.	SCOPOLAMIN.
Solutions commonly used	Atropin sulphate ½ to 2 per cent. Most commonly 1 per cent.	Scopolamin hydrobromate $\frac{1}{10}$ to ¼ per cent, (According to some it is identical with hyoscin.)
Effect on pupil	Almost maximum mydriasis; light reaction lost; increased by cocain.	Same as atropin.
Effect on accommodation	Complete cycloplegia; begins later than mydriasis.	Same as atropin.
Action— begins in reaches maximum in lasts from	 10 to 15 minutes 15 to 20 minutes. 6 to 10 days.	 7 to 10 minutes. 25 minutes. 4 to 7 days.
Effect on tension	Doubtful in normal eyes; increases tension in eyes predisposed to glaucoma.	Doubtful; tension not increased, according to Raehlmann.
Remarks	Atropin possesses some *disadvantages*, viz. (*a*) Absorption through lacrymal passages, causing poisonous symptoms (dryness and redness of throat and face. faintness, staggering, delirium). (*b*) Atropin infiltration, redness and swelling of eyelids and cheek. (*c*) Follicular conjunctivitis from frequent application. On account of its *strong* and *lasting* action it is the *best mydriatic for protracted use, as in iritis.*	Scopolamin is five times as powerful as atropin, but its effect is of shorter duration. In ¼ per cent. solution it is not more poisonous than atropin, and less so than duboisin. It is better borne by the conjunctiva than atropin. It should be used, therefore— (*a*) *Where atropin is not strong enough to break down posterior synechiæ.* (*b*) *Where atropin infiltration occurs.*

A few drugs not included in this table have been and are still used. Of these and are not to be recommended. Cocain as a mydriatic is not very useful when Holocain acts in the same way, without, however, affecting the cornea or circu-

* These tables have been drawn up chiefly by the aid of a paper by Dr. H. Schultz. *Archiv für Augen-*

mydriatics, myotics, and local anesthetics on the pupil, etc., have book by Dr. Louis Werner:

I.*

(Pupil Dilators).

Homatropin.	Ephedrin.	Mydrin.	Euphthalmin.
Homatropin hydrobromate 1 per cent.	10 per cent.	A mixture of homatropin and ephedrin: Homatropin 0.01 Ephedrin . 1 Water . . 10	Euphthalmin hydrochlorate 5 per cent.
Good mydriasis, but less than atropin.	Good mydriasis; light reaction retained.	Mydriasis greater than either constituent; light reaction feeble.	Maximum mydriasis; light reaction lost.
Marked, but not complete cycloplegia.	Little or no effect.	None.	Less than homatropin.
12 to 15 minutes. 40 minutes. 12 to 24 hours.	8½ minutes. 30 to 60 minutes. 5 to 20 hours.	8½ minutes. 30 to 40 minutes. 4 to 6 hours.	10 to 15 minutes. 60 to 80 minutes. 5 to 7 hours.
Not so liable to raise tension as atropin	Little or none.		None.
Homatropin is less powerful and less poisonous than atropin. On account of its action on accommodation, and the short duration of its effect, it is the *best mydriatic for estimating errors of refraction*. Its effect is increased by the addition of cocain.	It does not act quickly enough for use in practice as an aid to ophthalmoscopic diagnosis and is more useful when combined with homatropin.	The mydriasis being greater and of shorter duration than with either constituent, and having no action on the accommodation, it is well *suited for ophthalmoscopic diagnosis*	Although a little slower than homatropin, it is as good a mydriatic, but has the advantage of acting on the accommodation only in a slight degree, and its effects pass off much more quickly. It is also more powerful than mydrin, and therefore *the best mydriatic for ophthalmoscopic diagnosis*. It possesses no irritating or toxic effects, and does not injure the corneal epithelium.

daturin is the same as atropin. Hyoscyanin and duboisin are very active poisons employed alone, but it facilitates absorption, and increases the effect of other mydriatics lation, like cocain.

"Die älteren und neuern Mydriatica, Miotica und Anästhetica in der Augenheilkunde," *heilk.* Bd. xl., S. 125.

TABLE II.

Myotics (Pupil Contractors).

	Eserin Sulphate.	Pilocarpin Hydrochlorate.	Arecolin Hydrobromate.
Strength of solution commonly used	½ to 1 per cent. (the former less irritating).	1 to 5 per cent.	1 per cent.
Effect on pupil	Strong contraction of sphincter pupillæ, reducing pupil to pinhole size.	Same as eserin, but weaker.	The same as eserin.
Effect on accommodation	Spasm of ciliary muscle adapting eye for near point.	The same properties as eserine, but weaker.	The same as eserin.
Action—			
begins in	1 to 1½ minutes.	The same as eserin.	It acts more quickly and more powerfully than eserin, but its effects pass off very rapidly (1½ hours).
reaches maximum in	30 to 40 minutes.		
lasts from	24 to 36 hours.		
Effect on tension	It reduces the tension in glaucomatous conditions.	The same as eserin.	The same as eserin.
Remarks	When instilled into the eye it at first causes twitching of the eyelids and sometimes severe supra-orbital pain, due to spasm of accommodation. It is chiefly used for the reduction of abnormally high tension. It is also used in peripheral corneal ulcers, and after cataract extractions, to prevent prolapse of the iris. In some eyes, especially if hyperemic, it is liable to cause slight iritis. The fresh solution is white, but soon becomes red, without, however, losing its efficacy.	Is not so liable as eserin to cause iritis.	Is obtained from the areca nut. It may be of use in some cases in which eserin fails to act.

TABLE III.

Local Anesthetics used in Ophthalmology.

	Cocain Hydrochlorate.	Tropa-cocain.	Eucain B.	Holocain.
1. *Strength* solution in use	2 to 4 per cent.	3% (in 0.6% NaCl, to prevent irritation).	2 per cent.	1 per cent.
2. *Anesthesia* begins in	2 to 3 minutes.	½ to 2 minutes.	2 to 5 minutes.	½ to 1 minute.
3. Effect on *pupil* and *accommodation.*	Transient mydriasis and cycloplegia.	None.	None	None.
4. Effect on *cornea*	Tends to produce dryness and exfoliation of epithelium.	None.	Same as cocain, but in a less degree.	Same as cocain, in a slight degree.
5. Effect on *intra-ocular tension*	In normal eyes sometimes lowers tension, but raises it in eyes predisposed to glaucoma.	None.	None.	None.
6. Other effects	Causes slight smarting, followed by anemia and cold sensation, and occasionally marked retraction of eyelids.	Slight burning; no anemia.	Smarts more than cocain, and causes hyperemia. (Eucain A is too irritating for use)	Slight smarting and slight hyperemia
7. Remarks	Cocain as a local anesthetic possesses some *disadvantages.* It does not act so well in inflamed eyes. Its effects under heads 3, 4, and 5 are undesirable. Instillations rarely cause toxic symptoms, but they may occur after subconjunctival injection even of three-quarters of a grain. It is stated that cocain decomposes on boiling, and loses its effect; but we have repeatedly sterilized the same solution without any marked diminution in its action. It seems, however, to render the solution more irritating.	Acts more rapidly and more powerfully than cocain; is very stable and only half as poisonous.	Less poisonous and more stable than cocain, but is more irritating, especially in inflamed eyes.	More powerful and efficient than the other anesthetics. Can be sterilized without decomposition. Is an antiseptic. *It should not be used for subconjunctival injection or for lacrimal passages,* as it is more poisonous than cocain, resembling strychnia in its action.

It will be seen from the above that holocain is the best anesthetic for all purposes, except subconjunctival injection and treatment of the lacrimal passages. Some surgeons have derived benefit from it in the treatment of corneal ulcers, owing to its antiseptic properties. Several other local anesthetics (acoin, strophantin, etc.) have been tried, but they have proved either inferior or too irritating.

CHAPTER XII.

GLAUCOMA.*

The chief and essential symptom of this disease is Increased Intra-ocular Tension—increased hardness of the eyeball—due to overfullness of the globe.

There is Primary Glaucoma and Secondary Glaucoma.

In primary glaucoma, the increased tension comes on without any previous recognizable disease of the eye; and it is with it we have mainly to do in this chapter.

In secondary glaucoma, the increased tension comes on in consequence of obvious antecedent disease in the eye.

Primary Glaucoma.

Primary glaucoma is almost invariably a binocular disease, although it does not always attack each eye simultaneously, indeed it is more common for the disease to appear in the eyes with an interval of months or longer.

Of primary glaucoma there are two great kinds—the Non-inflammatory, Non-congestive, or Chronic Glaucoma; and the Inflammatory, Congestive, or more or less Acute Glaucoma. In using the term inflammatory here it is not implied that acute glaucoma is an inflammation in the strict pathological sense of the term, or, if so, to but a slight extent. The term is employed rather on account of some symptoms which are present (pain, redness of the eyeball, lacrimation), and which we are wont to see with inflammations of the eye—symptoms which are wanting in chronic glaucoma.

Increased intra-ocular tension, then, is the chief and essential symptom of glaucoma, whatever form of it may come before us; although this increased tension may not be present in the same degree, or, indeed, at all, at every time.

* From *γλαῦκος*, *sea-green*. The name was given to the disease by the old writers, on account of the greenish reflection obtained from the pupil in some cases. But this greenish reflection is seen in other diseased conditions, and therefore is not characteristic of glaucoma.

If the surgeon place the tips of his index fingers close together on a normal eyeball, and make gentle pressure with them alternately, he will observe that the eyeball pits slightly on this pressure, and that a sensation of fluctuation is given to the fingers. The amount of this pitting or fluctuation varies according to the degree to which the eyeball is filled with its humors, and also, to some extent, according to the thickness of the sclerotic coat, and is not precisely the same in every normal eye. The glaucomatous eyeball is felt to be more resistant—to be harder —than the normal globe.

But there are normal eyes which have a tension below the average normal tension, and others which have a tension somewhat above the average normal tension, and in eyes of the latter class it is occasionally difficult to decide whether or not the tension is abnormally high, especially if there happen to be symptoms which might be due to high tension. If it be a question of one eye only, then a comparison of its tension with that of its fellow decides the matter, for the physiological tension is always the same in each eye.

Some clinical experience is necessary before the surgeon can appreciate by palpation those degrees of tension which are just above or just below the normal; and no other method is equally satisfactory. Tonometers have indeed been invented for the purpose, but for ordinary use the educated fingers are to be preferred.

For the purposes of clinical notation Sir W. Bowman suggested some signs, which have been very generally adopted. Normal tension he indicated by the letter T, slight increase of tension by T + 1, still higher tension by T + 2, while T + 3 indicates stony hardness of the eyeball. In the same way diminished tension is T — 1, T — 2, and T — 3. T + ? and T — ? indicate that it is doubtful whether the tension be slightly above or below the normal. But the application of these symbols to the varying degrees of tension depends very much upon the observer. "T + 2," for instance, will not always convey precisely the same idea to every surgeon.

The other symptoms of glaucoma are largely due to the increased tension; but in chronic glaucoma there are by no means so many symptoms as in acute glaucoma. Let us now discuss these two great forms of primary glaucoma separately. And first as to **Chronic,** or **Non-Inflammatory, Glaucoma** (also

known as Simple Glaucoma, as Simple Chronic Glaucoma, and as Chronic Non-Congestive Glaucoma).

Symptoms.—The tension is raised. Sometimes the eye is very hard (T + 2, or more), and again it may be but slightly raised (T + 1). Even in one and the same eye the tension usually varies, and may be at one time too high and at another almost, or quite, normal.

The external appearance of the eye is usually quite normal, and the pupil reacts well to light. The anterior chamber is sometimes a little shallow.

On examination with the ophthalmoscope the optic papilla is found to be cupped. The optic disc, being the weakest part of the ocular wall, is the first place to give way to the high tension;




FIG. 102.—(*Ed. Jaeger*), sc, Sclerotic; ch, Chorioid; r, Retina; of, Optic nerve; ca, Intervaginal space; v, External sheath of the optic nerve; E, Excavation of the papilla; M, Margin of the excavation; lc, Lamina cribrosa.

and after a time it becomes depressed or cupped, the excavation being often deeper than the outer surface of the sclerotic, and the lamina cribrosa being pushed back (Fig. 102). This cupping of the papilla is a most important sign of glaucoma, and differs essentially in appearance from the physiological cupping (*vide* p. 88), inasmuch as it occupies the entire area of the papilla, and has steep, not shelving, sides. As shown in Fig. 102, the walls of the excavation are often hollowed out, and the ophthalmoscopic effect of this is to give to the retinal vessels the appearance of being broken off at the margin of the papilla (Fig. 103), where they pass round the overhanging edge of the excavation, and become hidden by it, while on the floor of the excavation they reappear.

The presence of an excavation may be recognized ophthalmoscopically, in the examination by the indirect method, by means of lateral motions of the convex lens. It will be then seen that, while the whole fundus seems to move along with the motion of the lens, the floor of the excavation apparently moves in the same direction, but at a slower rate. This parallax is the more marked the deeper the excavation. The phenomenon is explained by the accompanying figure (Fig. 104). If *o* be the optical center of the lens being used in the examination, and *b* and *a* two points lying one behind the other, the inverted




FIG. 103.—(*Ed. Jaeger*), a, Arteries; v, Veins; M, Margin of disc; K, Bending of vessels at margin of the disc; Vp, Vessels on the floor of the excavation; z, Glaucomatous ring.

images of these points will be situated at b' and a'. The line a' b' lies in the visual line of the observer; and if the lens be moved upward a very little, so that the optical center comes to o', the inverted images of *b* and *a* will be removed to b^2 and a^2. If the observer has not altered his point of view, it will seem to him that the point *b* has made a more extensive motion than the point *a;* or that it has moved more rapidly than *a*, and has glided between *a* and the observer. Short and rapid motions of the lens from side to side, or from above downwards, will best show the parallax.

In the upright image, the existence of an excavation may be ascertained by observing that a lens of a different power is required in order to obtain a clear image of the margin of the papilla and of its floor. The depth of the excavation may be estimated by noting the difference between these two lenses—*e. g.*, if the general fundus of the patient be emmetropic, and the emmetropic observer require 3 D to see the floor of the excavation, the depth of the latter is about 1 mm., and in the same proportion up to 10 D.

Besides being cupped, the optic papilla becomes atrophied from the pressure, and its consequent pallor serves to aid the diagnosis between this and a physiological excavation.

But we meet with cases in which the optic disc is cupped and




Fig. 104.

pale, and in which increased tension is either not present at the time of the examination, or is so slight as to be doubtful. And here the diagnosis between glaucoma and primary atrophy of the optic nerve with cupping of the disc is one of the most difficult to be met with. The examination of the field of vision may not always assist, for in each of these diseases it is liable to be contracted; but if the contraction occur with re-entering angles, it points to primary atrophy, for this form of field is not found with glaucoma. The effect of a myotic on the intra-ocular pressure may aid the diagnosis, for it would not materially influence normal tension, while it would reduce abnormally high tension.

The color sense usually remains normal in glaucoma until

a late stage, while in primary optic atrophy it is defective at an early stage. Again, if slight pressure with the tip of the surgeon's finger produces arterial pulsation at the optic papilla, it is suggestive of glaucoma, for in a normal eye considerable pressure is needed to produce this effect.

Spontaneous pulsation of the arteries on the optic papilla may be often noted in glaucoma, because blood can only be forced into these vessels by a pressure greater than that opposed to it. In the eye with normal tension there is no arterial pulsation,—and slight pressure with the tip of the finger would not bring it on,—for the tension of the coats of the vessels is greater than the intra-ocular tension; and, therefore, the blood passes on in a continuous stream. But in the decidedly glaucomatous eye the intra-ocular tension opposes so great an obstacle to the arterial flow that at the systole alone can it make its way through.

Arterial pulsation also occurs, although rarely, in exophthalmic goiter (see chap. xviii.); and it occurs where the pressure in the arteries themselves is low (weak heart's action, aortic regurgitation, etc.), although that in the vitreous chamber be normal.

Around the margin of the glaucomatous excavation, especially in chronic glaucoma, one usually sees the whitish appearance, termed the glaucomatous ring (Fig. 103), which is said to be due to atrophy of the chorioid from pressure.

The acuteness of central vision is diminished, and increasing dimness of sight is the only symptom of which the patient complains in chronic simple glaucoma.

Besides this, the field of vision becomes contracted in consequence of interruption to conduction in the retinal nerve-fibers from pressure on them at the margin of the depressed optic papilla. This contraction of the field must always be carefully examined for with the perimeter in eight or ten meridians. A hurried or careless examination of the field is worse than useless, for it may lead astray in the diagnosis and prognosis. The contraction commences at the nasal side, as a rule, while at the same time central vision is lowered, and later on the temporal portion of the field becomes contracted, and gradually absolute blindness is brought about.

The progress of the disease is extremely slow and insidious, extending often over several years, and ends in total blindness if untreated. It usually attacks both eyes, but generally one of

them long before its fellow. Sometimes chronic simple glaucoma, after a time, takes on the acute or the subacute form.

Acute, or Inflammatory, Glaucoma (also called Acute Congestive Glaucoma).—In this form the increase of tension is always very marked. In addition to this there are the following symptoms:

Diminished Depth of the Anterior Chamber, from pushing forwards of the lens and iris.

Diminution of the Refracting Power of the Eye, by reason of the nearer approach of the latter to a globular shape.

Diminution of the Amplitude of Accommodation, and Anesthesia of the Cornea, owing to pressure on the ciliary nerves as they pass along the inner surface of the sclerotic.

Opacity of the Cornea, giving its surface a peculiar steamy or breathed-on appearance, due to edema of the corneal tissue and epithelium, by infiltration into them of the intra-ocular fluids from high tension. A similar opacity of the cornea is sometimes seen in iritis and iridochorioiditis, and in interstitial keratitis.

Indistinctness of the Pattern of the Iris, similarly due to edema.

Opacity of the Aqueous and Vitreous Humors.

Dilatation and Immobility of the Pupil, the result, according to some, of paralysis of the ciliary nerves, but, according to others, of anemia of the iris from pressure on its vessels. The pupil is oval, with its long axis vertical.

The Episcleral Veins are large and tortuous, owing to the pressure on the vasæ vorticosæ preventing the discharge by those channels of the chorioidal venous blood, which must then pass off by the anterior ciliary veins.

Subjective Appearances of Light and Color, and colored halos or rainbows around lamps and candles (iridescent vision), are complained of. Similar appearances are sometimes experienced by persons suffering from chronic conjunctivitis.

Pain is a very marked symptom of acute glaucoma, both in the eye, and radiating over the corresponding side of the head. This pain is often very violent.

Vision is greatly affected, and the field of vision will be found contracted, in cases of some standing.

The Optic Papilla, when the media are sufficiently clear to admit of its being examined, is seen to be cupped if the disease have continued sufficiently long to bring about this change.

In acute glaucoma we recognize certain *Premonitory Symp-*

toms—viz., sudden diminution of the amplitude of accommodation, evidenced by the rapid onset or increase of presbyopia, and the consequent necessity for higher + glasses for near work; and the occasional appearance of colored halos around the flames of lamps or candles, with attacks of fogginess of the general vision. The duration of one of these foggy attacks may be from a few minutes to several hours. Such attacks are apt to occur after a sleepless night, or after a meal, and are sometimes accompanied by peri-orbital pains. Slight opacity of the aqueous humor, and sluggishness of the pupil, with some dilatation, are present during an attack; but the eye afterwards returns to its normal condition, and remains so for weeks or months, until another similar attack comes on. Such a premonitory stage may last a year or longer, but cases also occur in which there is no premonitory stage.

The most favorable time for operative interference is during this prodromal stage. It can then be performed with technical accuracy in an eye free from congestion and with a normally deep anterior chamber. There is, too, as yet no loss of sight, nor any degeneration of the tissues of the eyeball. The difficulty is to induce patients to consent to operation at this period.

The onset of the *True Glaucomatous Attack* is usually at night. It is accompanied by violent pain radiating through the head from the eye, by pericorneal injection, chemosis, and lacrimation. The aqueous humor is cloudy, the anterior chamber shallow, the iris discolored, and the pupil dilated to medium size and of oval shape, the cornea steamy and anesthetic. The patient frequently complains of subjective sensations of light, and vision is very defective, or may be quite wanting. Vomiting very frequently accompanies acute glaucoma, and has often led to errors of diagnosis, the patient's ailment having been taken to be a gastric disease, while the ocular symptoms were regarded as mere coincidences, such as a cold in the eye, neuralgia, etc.

An attack like that just described may, to a great extent, pass away in the course of a few days, but a complete remission of all the symptoms does not take place. Some defect of central vision is left, or, it may be, some slight peripheral defect in the field of vision; the tension does not become quite normal again, and the pupillary motions remain slightly sluggish. Another acute attack of glaucoma comes on in the course of some weeks or months, and it, too, may pass away, leaving the eye in a still worse

condition than it found it. The attacks gradually become more frequent; and if in the intervals the eye be examined the cornea and vitreous humor will be found more or less opaque, the optic papilla cupped, and an arterial pulsation may be discovered. Finally, there is no remission from the attack, the violent glaucomatous symptoms become permanent, and all vision is forever destroyed.

Even when vision has been destroyed the high tension continues, and gradually produces disorganization of the tissues of the eyeball (glaucomatous degeneration). The iris becomes atrophied, the lens becomes opaque, and the cornea frequently ulcerates, while hemorrhages are apt to occur in the anterior chamber. In time the excessive intra-ocular tension causes staphylomatous bulging of the sclerotic in the ciliary region, or further back; and, finally, such eyes may become the subjects of acute purulent chorioiditis, and end in phthisis bulbi.

Acute glaucoma almost always comes on in both eyes, either at the same time, or with an interval, it may be of weeks, or of months.

The reason why there is so marked a difference between the symptoms and course of chronic and of acute glaucoma is probably that in the former the increase of tension is very gradual, and therefore the eye gradually becomes accustomed to it; while in acute glaucoma the increase is rapid or sudden, and the circulation of the eye has not time to accommodate itself to the new state of things.

Glaucoma Fulminans is the name given by von Graefe to a form of the disease which is more acute than the ordinary acute glaucoma just described. It has no premonitory stage, and, coming on with all the symptoms of acute glaucoma greatly exaggerated, does not remit, and causes complete permanent destruction of vision in the course of a few hours. It is a rare form.

Subacute Glaucoma.—This form differs from acute glaucoma in that its premonitory stage merges gradually into the actual disease without the occurrence of an acute attack. The eye gradually becomes hard, the pupil dilated, the anterior chamber shallow, the aqueous humor opaque; while the cornea is "steamy" and anesthetic, and the episcleral veins are distended. Ophthalmoscopically the cupped disc and pulsating arteries may be seen, when the opacities of the media permit. Vision sinks, and the field is contracted towards its nasal side. The progress

of the disease is very slow; and in its course attacks of ciliary neuralgia, with greater increase of the tension, greater opacity of the aqueous humor, increase of the corneal opacity and anesthesia, and further dimness of vision, are experienced. These




FIG. 105.—Diagrammatic representation of normal condition. *l.* Angle of anterior chamber, and ligamentum pectinatum. *s.* Canal of Schlemm. *p.* Venous plexus of Leber.

attacks pass off again in the course of a few days or hours, leaving the eye harder and blinder than before. The subacute glaucoma sometimes takes on the acute form. It is liable to bring




FIG. 106.—Diagrammatic representation of glaucomatous condition. *l'.* Obliterated angle of anterior chamber.

about the same glaucomatous degeneration of the eye as does the latter.

Etiology of Glaucoma.—Glaucoma is a disease of advanced life, occurring most usually after fifty years of age, and rarely

under the thirtieth year. It is not peculiar, or more common to any one constitution or temperament. Anxiety, sorrow, and influences in general which depress the spirits have often been noticed to precede the onset of acute glaucoma.

As regards the *Pathology of Glaucoma* the theory which of late years has obtained most acceptation owes its origin to Max Knies [1] and Adolf Weber [2] and is known as the Retention Theory. These observers ascertained that in glaucomatous eyes the periphery of the iris lies in contact with the periphery of the cornea (Figs. 105 and 106) in the region of the canal of Schlemm, venous plexus, and ligamentum pectinatum. But this region and these tissues had previously been proved by Leber [3] to be the ways of exit of the effete intra-ocular fluids, which flow to that point from the posterior part of the aqueous chamber through the pupil. Weber and Knies concluded, therefore, that the blocking of these passages, from the close application of the iris, caused glaucoma by preventing the effete fluids from escaping: and that thus the disease was one of retention rather than of hypersecretion, as it had previously been considered to be. Weber believed that swelling of the ciliary processes, from one cause or another, pushes the periphery of the iris forwards, and gives the starting point for glaucoma.

Brailey [4] to a certain extent adopts this view of Weber, but regards [5] a chronic inflammation of the ciliary processes and periphery of the iris, with distention of the blood-vessels of these parts, to be the chief factor in the earliest history of the disease.

Max Knies [6] also regarded glaucoma as an iridocyclitis, which, owing to varying intensity, produces the different forms of the disease.

Priestley Smith [7] adopts the retention theory, and holds that the main predisposing cause of primary glaucoma is an insufficient space between the margin of the lens and the structures which surround it; and he attributes the greater liability of elderly people to the progressive increase in the size of the lens which he has proved [8] to occur as life advances. In eyes in which the circumlental space is insufficient, by reason either of the original structure of the eye—and small eyeballs, as Priestley Smith has shown, are specially liable to primary glaucoma, a fact often demonstrated by the small size of the cornea in the eyes attacked—or of the enlargement of the lens, any condition which tends to overfill the veins of the head and uveal tract may initiate

an attack of acute glaucoma as follows: An increase in the amount of blood in the uveal tract must be compensated by the expulsion of some other fluid from the eye; consequently, the aqueous humor filters out more rapidly than is normal at the angle of the anterior chamber. As the contents of the chamber diminish, the lens and iris move forwards towards the cornea. Now in the normal eye, and especially in the youthful eye, this compensation is effected without danger to the angle of the anterior chamber, because the lens is comparatively small, the circumlental space large, and the anterior chamber deep. But when the lens and ciliary processes are already in close relation to each other, and the anterior chamber already shallow, then any increased fullness of the uveal tract involves danger to the angle of the chamber. The turgid ciliary processes find insufficient space for their expansion; they are carried forwards together with the lens, and, pressing upon the base of the iris, lock up the angle of the anterior chamber. Thereupon, the further escape of fluid being impossible, high tension of the eyeball is established. According to this explanation, then, the high tension is due to impeded escape of the intra-ocular fluid, and depends, primarily, rather upon an increase in the amount of blood in the eye, than on an excess of the intra-ocular fluid. Priestley Smith considers that, in chronic simple glaucoma, the predisposing causes are the same as in acute glaucoma; but that in the former, the vascular disturbance being gradual and slight, the vessels adapt themselves to the slowly increasing pressure, and the angle of the anterior chamber is more or less compressed, but not tightly closed.

Von Graefe [9] believed that a serous chorioiditis lay at the root of the disease, which he thought was caused by exudation of serous fluid into the vitreous humor; while Donders,[10] von Hippel and Grünhagen,[11] and others held that irritation of the fifth pair of nerves, governing the secretion of the intra-ocular fluids, gave rise to hypersecretion of those fluids.

Others, again, held that changes in the sclerotic, rendering it rigid, and leading to some shrinking of it, caused the increased intra-ocular tension.

Laqueur [12] believes that some such sclerotic changes produce obstruction of the posterior ways of exit of the intra-ocular lymphatics—namely, those which pass out with the four vasæ vorticosæ—and that glaucoma depends largely upon this obstruction.

But these theories are now obsolete, and I mention them merely as being of historical interest.

Treatment.—The performance of an iridectomy is the means discovered by von Graefe,[13] in the year 1857, for the cure of glaucoma, a disease which had hitherto been incurable. This measure held an undisputed position as the sovereign remedy for the disease until a few years ago, and even yet has not suffered much from the competition of the operation of sclerotomy.

To insure the success of an iridectomy for glaucoma, so far as possible, it is necessary (1) that the incision should be peripheral—*i. e.,* as far back in the corneo-sclerotic margin as is compatible with the introduction of the knife into the anterior chamber, and with the avoidance of injury to the ciliary body; (2) that the portion of iris removed should be wide—*i. e.*, involving about one-fifth of the circumference of the iris (see Fig. 98).

It is, moreover, important to withdraw the knife very slowly from the anterior chamber, when the corneo-scleral section is complete, in order that the aqueous humor may flow off gradually, lest an intra-ocular hemorrhage from the sudden reduction of tension should occur. The portion of iris should be most carefully abscised, so that no tag of it may remain in the wound, and become caught in the cicatrix in the course of healing. Such an occurrence is apt to produce a cystoid cicatrix, which may at a later period become the starting-point of irritation, and even of serious inflammation. Some operators prefer von Graefe's cataract knife for the performance of the operation, but the ordinary lance-shaped iridectomy knife is the instrument usually employed. For the purpose of reducing the intra-ocular tension it matters nothing what region of the iris be abscised; but, as a rule, the upper quadrant is to be preferred, for there the resulting coloboma, being covered to a great extent by the upper lid, will give rise to less diffusion of light than in any other position.

Immediately after the operation, palpation of the eyeball should show a marked diminution of tension. When this is not so the prognosis is unfavorable. Should an increase of tension occur on the day after the operation, it is of no consequence, as it passes off again in the course of the next few succeeding days. Until then the anterior chamber will not be restored, and we see cases where the anterior chamber does not appear for a week or more. The bandage should be worn until the anterior chamber is completely restored. It is desirable to perform the operation

with general anesthesia, to secure technical accuracy. The pain for some time after the operation is considerable, and should be relieved by a hypodermic injection of morphia in the corresponding temple.

Very occasionally, immediately after iridectomy, although the operation may have been faultlessly performed, the case takes what we call a malignant course. In these cases the lens seems to be violently pushed forwards, blocking the wound, obliterating the angle of the anterior chamber, and preventing any fluid from escaping from the eye, so that very soon it is as hard, or harder, than before. This complication seems to be caused by the retention of fluid behind the lens, and is more likely to occur in cases of chronic simple glaucoma than in the acute forms of the disease.

In chronic simple glaucoma, if any cataract be present as well, it very often develops rapidly after the iridectomy; and in nearly all cases a certain amount of disturbance of vision results from astigmatism produced by the operation. All these risks and disadvantages must be faced by the patient, and put against the certainty of complete and hopeless blindness if the operation be not performed.

Unless the *Operation of Posterior Sclerotomy* be employed with success, all eyes which take on the malignant course just mentioned are inevitably lost, are apt to become very painful, and must often be excised. A broad needle, or a Graefe's knife, is entered through the sclerotic, 8 or 10 mm. behind the outer margin of the cornea, and the blade is given a quarter turn on its axis, so as to make the wound to gape, or the latter may even be somewhat enlarged in a meridional direction. At the same time gentle pressure is applied, by means of the upper lid, on the center of the cornea. This causes fluid to escape through the scleral wound by the side of the knife, and it also causes the lens to go back into its place, with restoration of the anterior chamber. The pressure on the cornea may be maintained with advantage for a minute or somewhat longer. This proceeding has also been suggested as a cure for glaucoma, but has not yet been put into practice. It is probable that it would only temporarily reduce the intra-ocular tension.

As a rule the more acute the form of glaucoma, and the earlier in the disease the iridectomy be performed, the more favorable is the prognosis in respect of the result which may be expected. The saving of normal vision can only be looked for in those

cases where it has as yet fallen but little, or not at all, below the normal, and where the contraction of the field has barely commenced. When the disease has interfered seriously with vision (of course I do not refer here to the enormous loss of sight immediately attendant upon an attack of acute glaucoma, for this is usually restored) we may not expect more than the retention of the *status in quo*. But even in this respect our prognosis should be most guarded, especially in chronic simple glaucoma, when the disease has advanced so far that the contraction of the field is found to have approached close to the fixation-point, although central vision may still be fairly good. Because in such cases, while the iridectomy may prove successful in so far as reduction of tension is concerned, yet the contraction of the field—*i. e.*, the progress of the atrophy of the optic nerve—is often not arrested, and shortly afterwards may be found to engulf the center of vision. Indeed, where the contraction is near the fixation-point at one side, and is advanced in other directions, it is better not to operate. In general, while the result obtained from iridectomy in acute and subacute glaucoma, on the bases above laid down, can be regarded as amongst the most satisfactory in the whole range of ophthalmology, in chronic simple glaucoma iridectomy does not act with the same degree of certainty. Nevertheless, iridectomy, as holding out the only prospect of retention of any vision, should be recommended to the patient—except in every advanced cases—and that at as early a period in the disease as possible, before the field has become much contracted. When, as is often the case, the patient comes with one eye far advanced in the disease, and the other in an early stage of it, it is a question which eye should be first operated on. Probably most surgeons operate on the worst eye first, with the result that the measure having been applied too late, the result is unsatisfactory, and the patient declines to allow the better eye to be operated on. Hence some operators, with whom I agree, advise operation on the better eye first.

In cases of acute or subacute glaucoma it has frequently been observed that shortly, even within a few hours, after the performance of the iridectomy, the other eye, previously healthy, or, at most, affected with but slight premonitory symptoms, is attacked with glaucoma. It is probable that this is due to dilatation of the pupil, with crowding of the iris into the angle of the anterior chamber, in consequence of confinement in the dark

room, and eserin should be put into the sound eye as a precaution.

It may here again (*vide* p. 257) be stated that the use of atropin, or of any other mydriatic, in an eye with a tendency to glaucoma is liable to bring on an acute attack of the disease, and must be carefully avoided in such cases.

If the tension be not relieved by the iridectomy, a supplemental iridectomy may be performed after a time, and von Graefe recommended that it should be placed at the opposite side of the pupil from the first coloboma. Dr. Gruening,[14] of New York, performed five iridectomies in one eye for glaucoma. At the end the patient had no iris, but enjoyed a vision of 6-6.

The Mode of Action of the Operation is not clearly known. Von Graefe at one time believed it to act by diminution of the secreting surface of the intra-ocular fluids. De Wecker [15] and Stellwag [16]—even previously to the formulation by Knies and Weber of the retention theory of glaucoma already referred to—held that the cure depended, not on the removal of the portion of iris, but on the incision in the corneo-sclerotic margin, or, rather, on the nature of the cicatrix resulting from that incision. They maintained that this cicatrix was formed of tissue, which admitted of a certain amount of filtration through it of the intra-ocular fluids, and that in this way the intra-ocular tension was kept down to the normal standard. This theory subsequently gained support from that of Knies and Weber.

Priestley Smith has satisfied himself that in a large number of successful iridectomies the success is due to a permanent corneo-scleral fistula—not merely a filtration cicatrix—having been formed. The same view is held by Treacher Collins,[17] who finds that this permanent gap is maintained by a prolapse of a fold of iris into the wound. The latter author, indeed, entirely and definitely discards the filtration-cicatrix theory, for which he considers there is no evidence. In those cases where a fistula, as described, is not formed by the operation, Treacher Collins considers that the obstruction becomes freed, either by the iris being torn away at its thinnest part,—that is, its extreme root,—thus leaving a large portion of the filtration angle open for drainage; or by the escape of the aqueous and the drag on the iris, incidental to the iridectomy, being sufficient to dislodge the periphery of an iris which has only recently come into apposition with the cornea.

De Wecker, Stellwag,[18] and Quaglino[19] sought to produce the corneo-scleral filtration cicatrix without the removal of a portion of iris. The peripheral position of the wound, however, rendered the proceeding difficult or impossible, owing to the tendency to prolapse of the iris which necessarily existed. The introduction of eserin into ophthalmic practice at last enabled de Wecker to place the operation on a surer footing, as the myosis produced by instillation of a solution of this drug into the eye insured the operator, to a great extent, against the danger of prolapse of the iris; and hence.

Sclerotomy, as the operation is called, came to be cultivated as a method for the relief of glaucoma. It has hitherto been employed mainly in chronic simple glaucoma—a form in which, as I have stated, iridectomy is less satisfactory than in acute or subacute glaucoma. Care must be taken that the pupil is contracted to pinhole size, or nearly so, when the operation is about



FIG. 107.

to be performed, as otherwise the danger of prolapse of the iris is very great. In those cases where eserin will not produce a sufficient myosis, sclerotomy should not be performed.

The instrument used for performing the operation is von Graefe's cataract knife. A speculum having been applied, and the eyeball fixed, the point of the knife is entered into the anterior chamber, through the corneo-sclerotic margin, at a point of its circumference corresponding to that selected for the puncture in cataract extraction, but 1 mm. removed from the corneal margin, as represented at *a* in Fig. 107. The counter-puncture is made at a point (*b*) corresponding to this, at the other side of the anterior chamber. With a slow sawing motion of the knife the section is enlarged upwards, until only a bridge of tissue, about 3 mm. broad, remains at *c,* and this is left undivided, the better to guard against prolapse of the iris. The knife is now slowly withdrawn from the eye, care having been first taken that the aqueous humor is thoroughly evacuated, which

can be effected by tilting the edge of the knife slightly forwards, so as to make the lips of the wound gape somewhat. If the pupil be quite round at the conclusion of the operation, the bandage may be applied, a drop of solution of eserin having been first instilled; but if the pupil be oval, or of other irregular shape, a tendency to prolapse of the iris is indicated, and the hard rubber or silver spatula should be introduced into the anterior chamber, to restore the pupil to its normal shape by gentle pushing of the iris. If there be an actual prolapse of the iris, an attempt may be made to repose it with the spatula; but should this not prove satisfactory the prolapse is to be abscised with scissors, thus turning the sclerotomy into an iridectomy.

The operation has not gained many adherents, but Mr. Cross, of Bristol,[20] regards it as of great value in the earlier stages of chronic glaucoma, before the sight is much reduced. The sclerotomy he performs is that first proposed by Snellen. The pupil, having been contracted with eserin, a broad keratome is passed —as though an iridectomy were about to be made—from a puncture about 2 mm. outside the corneal margin into the anterior chamber. A wide wound is made, and the knife is carefully withdrawn to avoid prolapse of iris. Mr. Cross recommends this sclerotomy, too, when high tension recurs after having been temporarily relieved by an earlier iridectomy. The operation may be repeated again and again without danger, should the case demand it.

Priestley Smith and Treacher Collins explain the cure by sclerotomy as they do that by iridectomy.

Sympathectomy.—Excision of the superior cervical ganglion of the sympathetic nerve is a new method for the treatment of glaucoma, which I shall merely mention; for, although of interest, it is not sufficiently established to bring it within the scope of this work. Those who desire information concerning the method will find it by aid of the references [21] given at the end of this chapter. Sympathectomy is not recommended as a cure in itself, but rather as a supplement to iridectomy in cases of chronic glaucoma. It is said to act by contraction of the pupil, and reduction of the intra-ocular blood pressure.

*The Treatment of Glaucoma by Myotics.**—Eserin and pilocarpin as eye-drops in 1 per cent. solutions often have the power

* The action of the Myotics which are most in use will be found in Table II., p. 318.

of reducing glaucomatous tension. This power depends on the contraction of the pupil, and consequent drawing away of the base of the iris from the angle of the anterior chamber; and, if the myopic does not contract the pupil greatly, it will not reduce the tension. Cases of acute glaucoma, brought on by the injudicious use of atropin, may frequently be completely and permanently relieved by a myotic instilled a few times. In acute glaucoma of the ordinary type, the use of a myotic in the premonitory stage will often postpone the true glaucomatous attack, and even sometimes relieve the latter for the time; but the myotic treatment cannot produce a radical cure, and it should only be used to preserve the health of the eye, until the operation is performed. In chronic simple glaucoma, also, myotics bring down the tension if they contract the pupil, and may be used in those cases where the patient positively declines an operation, or where an operation in the fellow eye has not resulted satisfactorily, or where an operation is contra-indicated by a very contracted field. The anti-glaucomatous action of the myotic only lasts so long as the pupil is contracted, and it must consequently be applied once or twice in the twenty-four hours.

In the myotic treatment of glaucoma, Priestley Smith recommends the combination of cocain with the myotic in such proportions (say, about 1-4 per cent. of cocain to 1 per cent. of the myotic) that the myotic will have the mastery over the pupil. For although, like every dilator of the pupil when used alone, cocain may promote high tension, yet it has the powers, invaluable in glaucoma, of contracting the ciliary blood-vessels, and of diminishing the sensibility of the ciliary nerves; and, when used in the foregoing manner, the advantage of each drug may be obtained, without any of the disadvantages of either.

It may here be once more stated that, while myotics possess the power of reducing glaucomatous tension, atropin, and all mydriatics, bring on glaucoma, where there is already a tendency to it. In all old people, therefore, before atropin is used, it is well to ascertain that the tension is not too high.

Treatment of Painful Blind Glaucomatous Eyes.—Eyes blind of acute glaucoma may, as I have stated, continue to be painful; and may in this way render the patient's life very miserable. Iridectomy is very commonly performed to relieve the pain, although all hope of restoration of sight is lost; but the operation sometimes fails in its object. Neurectomy (p. 280) seems to

offer a more certain result, and of course excision, or evisceration, would have the same effect. Mules' operation should not be performed here.

Secondary Glaucoma.

In addition to the different forms of primary glaucoma above described, we find, as already stated, that high tension occurs as a sequel of diseased conditions previously existing in the eye. There are several diseased states which are liable to become complicated with glaucomatous tension; but it should be clearly understood that, in almost every instance, the immediate cause of the high tension is the same as in primary glaucoma—namely, a closure of the angle of the anterior chamber.

The following are the chief conditions which are liable to lead to secondary glaucoma:

a. Complete Posterior, or Ring Synechia (*vide* p. 252). The iris, being pushed forwards by the aqueous humor pent up behind it in the posterior part of the aqueous chamber, is pressed tightly against the cornea, and obliterates the angle of the anterior chamber and the ways of exit. An iridectomy relieves the high tension here.

b. Perforating Wounds or Ulcers of the Cornea, followed by incarceration of the iris in the resulting cicatrix. The iris, being drawn tautly towards the cornea, a large portion, or the whole, of the filtration angle may be closed by it. An iridectomy is indicated. Lang divides anterior synechiæ by means of his twin knives.

c. Dislocation of the Crystalline Lens into the Anterior Chamber. Here the normal flow of the intra-ocular fluids through the pupil, on its way to the filtration angle, is arrested by reason of the presence of the lens in the anterior chamber. The onward current then presses the iris against the posterior surface of the lens, and the root of the iris, which is unsupported by the lens, against the periphery of the cornea, and in this way the angle of the anterior chamber is closed. In these cases the lens must be removed from the eye.

d. Lateral (traumatic) Displacement of the Crystalline Lens. The lens, being pushed in between the ciliary processes and the vitreous humor, drives the root of the iris forwards against the cornea at that place, while in other parts of the circumference

the displaced vitreous acts in the same way. In these cases, too, the lens must be removed.

e. Injury of the Crystalline Lens (*vide* chap. xiii.). The swelling lens pushes the iris forwards against the angle of the anterior chamber. Evacuation of the lens should be performed.

f. After Cataract Extraction. For explanation of this see chap. xiii.

g. Intra-ocular Tumors (*vide* p. 293). The growth of the tumor gives rise to a transudation of serum from the chorioid which detaches the retina, and after a time pushes the lens, the ciliary processes, and the iris forwards, and thus closes the filtration angle.

h. Serous-Cyclitis, or Iritis. Here the filtration angle is not closed. Priestley Smith thinks that the increased tension is due to diminished filtration-power of the eye, and perhaps by tissue changes around the filtration angle, and by deposit of exudation in the angle of the anterior chamber.

Another, and very peculiar, form of secondary glaucoma is **Hemorrhagic Glaucoma.**—Retinal hemorrhages of the ordinary type are sometimes followed, a few weeks later, by increased intra-ocular tension, which generally assumes the symptoms of acute or subacute glaucoma, and, more rarely, those of chronic simple glaucoma. A satisfactory explanation for these cases has not, so far as I am aware, been offered. When such a glaucoma has become pronounced, it is not usually possible to distinguish it from a primary form of the disease.

Treatment.—The disease is practically hopeless. Iridectomy is more likely to do harm than good, the operation being almost invariably followed by fresh intra-ocular hemorrhages, and by a further increase of tension. Sclerotomy is said by some to act with fairly good results in hemorrhagic glaucoma. The myotic treatment is powerless.

Congenital Hydrophthalmos.

Also known as Buphthalmos, and as Cornea Globosa. It is a glaucomatous disease of early childhood, of which the incipient stages are believed to be intra-uterine. The cornea becomes enormously enlarged in diameter, the anterior chamber deep, the iris trembling, and the sclerotic thinned. Increase of tension,

often attended by severe pain and cupping of the optic papilla, are usually present.

The Pathology of the disease is obscure. Treacher Collins [22] holds that it is a failure in the separation of the iris from the back of the cornea at its extreme periphery, in course of the development of the eye, whereby the angle of the anterior chamber is blocked; while E. von Hippel [23] believes it to be the result of an intra-uterine inflammation.

Treatment.—Iridectomy and sclerotomy are alike followed by disastrous results in this disease. The myotic treatment is the only one applicable, and in a few cases it arrests the disease.

References.

[1] "Von Graefe's Archiv," xxii. pt. 3. p. 163, and xxiii. pt. 2. p. 62.
[2] *Ibid.*, xxiii. pt. 1. p. 1.
[3] "Von Graefe's Archiv," xix. pt. 2. pp. 87-185.
[4] "Ophth. Hosp. Rep.," x. p. 282.
[5] *Ibid.*, ix. p. 199, and x. pp. 14, 89, 93.
[6] "Archiv. f. Augenheilkunde," xxxviii. p. 193.
[7] "Ophth. Hosp. Rep.," x.; "Trans. Internat. Med. Congress," 1881; "Ophthalmic Review," vol. vi. p. 191; "Pathology and Treatment of Glaucoma." London, 1891.
[8] "Trans. Ophth. Soc. U. K.," iii. p. 79.
[9] "Archiv f. Ophthal.," xv. pt. 3. p. 108.
[10] *Ibid.*, ix. pt. 2. p. 205.
[11] *Ibid.*, xiv. pt. 3, xv. pt. 1, and xvi. pt. 1.
[12] "Von Graefe's Archiv," xxvi. pt. 2.
[13] "Archiv f. Ophthal.," iii. pt. 2. p. 456.
[14] "Trans. American Ophthalmological Society," 1901, p. 314.
[15] "Bericht der Ophthal. Gesellsch." Heidelberg, 1869.
[16] "Der Intraoculare Druck," etc. Vienna, 1868.
[17] "Roy. Lond. Ophthal. Hosp. Rep.," December, 1891, and "Researches into the Anatomy and Pathology of the Eye," p. 111. London, 1896.
[18] "Bericht der Ophthal. Gesellsch." Heidelberg, 1871. "Chirurgie Oculaire," p. 212.
[19] "Annali di Ophthalmologia," i. pt. 2. p. 200.
[20] "Brit. Med. Jour.," August, 1900.
[21] Abadie, "Arch. d'Ophthal.," xix. p. 94; Axenfeld, "Sammlung swangloser Abhandungen a. d. Gebeiete der Augenheilkunde," iv. 1. p. 1.; Ball, "Internat. Ophthal. Congress." Utrecht, 1899; Demicherie, "Ann. d'Oculistique," cxxi. p. 188; Work Dodd, "Brit. Med. Jour.", August, 1900; Grunert, "Bericht der Ophthalmologischen Gesellschaft." Heidelberg, 1900, p. 15; Jonnesco, "Internat. Med. Congress," Paris, 1900.
[22] "Researches into the Anatomy and Pathology of the Eye," p. 104. London, 1896.
[23] "Bericht d Ophthal. Gesellsch." Heidelberg, p. 225, 1897.

CHAPTER XIII.

DISEASES OF THE CRYSTALLINE LENS.

Cataract, by which is meant an opacity of the lens, may be said to be the only disease of this part of the eye. Cataract may be complete—*i. e.,* occupying, in its final stage, the whole, or nearly the whole, of the lens; or partial—*i. e.,* occupying only part of the lens, and with little or no tendency to extend to other parts of it.

Complete Cataracts.

Of these, the most common is **Senile Cataract.** It occurs in persons of over fifty years of age, rarely in those under forty-five years of age.

Progress, Pathogenesis, and Etiology of Senile Cataract.—In commencing or incipient senile cataract the opacity is found in the cortical layers of the lens, especially at its equator, and in the latter position can often only be detected with transmitted light from the ophthalmoscope mirror, or with focal illumination, even when the pupil is dilated with atropin. This opacity takes the form of lines, or of triangular sectors, of which the bases are towards the equator of the lens, while the apices are towards its center. These lines and sectors look black with transmitted light, but gray with focal illumination, and between them clear lens-substance is present. Or, incipient cataract may first appear as a diffuse opacity in the layers surrounding the nucleus of the lens. Or, the opacity may commence both near the equator and around the nucleus at about the same time. Or, again, the opacity may in the beginning be disseminated through the cortex, in the form of flocculi, dots, and lines. In some cataracts, in a very incipient stage, there are no absolute opacities; but with weak transmitted light—*i. e.,* from a plane mirror—numbers of fine dark lines will be seen in the lens, which vanish and reappear according as the incidence of the light is altered; while a little later true opacities make their appearance. Gradu-

ally the cataract extends to other parts of the lens, until the whole cortical portion is opaque.

In senile cataract the very nucleus itself does not become cataractous, although it is usually sclerosed (harder and drier). Sclerosis of the nucleus of the lens is a physiological condition of advanced life, and will be found in many an eye where there is no cataract. It gives to the senile non-cataractous lens, as seen with a dilated pupil or with focal illumination, a peculiar smoky appearance, which is often mistaken by inexperience persons for cataract; but examination with transmitted light will show that there is no opacity. When a senile cataract has become complete, the sclerosed nucleus imparts to its center a brownish or yellowish hue, while the other parts of the lens are of a grayish-white. As a rule the most peripheral layers of the cortex are the last to become opaque. Accordingly as the lens becomes opaque, it often swells somewhat, and the anterior chamber consequently becomes a little shallower.

Until the whole cortex is opaque a clear interval will be present between the iris and the cataractous part, and on examination with the oblique light a shadow of the iris will be thrown on the cataractous part at the side from which the light comes; and the cataract, in this way, is proved to be immature in the strict sense. If the whole cortical substance be opaque, the thickness of the capsule alone will intervene between the pupillary margin and the opacity. In addition to this examination with the focal light the pupil should be dilated, and the lens examined by transmitted light from the ophthalmoscope mirror, when a completely opaque cataract should permit of no red reflection being obtained in any direction from the fundus oculi.

As soon as the whole of the cortical substance has become opaque, the swelling of the lens begins to subside, and the anterior chamber finally regains its normal depth. If there be no glittering sectors in the cortex, the cataract is now "mature," or "ripe" for operation—*i. e.*, if an extraction operation be now undertaken, it is possible to deliver the lens in its entirety; whereas, prior to this stage, some cortical substance would have been liable to adhere to the capsule, and be left behind.

But a cataract is immature, despite the absence of shadow from the iris, of the illuminable pupil, and even though the anterior chamber be of normal depth, if the cortex present well-marked, glittering sectors. The glitter of the different sectors varies

with the angle of illumination, so that the surface appears faceted. In such a lens there are thin transparent flakes, as well as opaque flakes, close beneath the capsule; and, if extraction be undertaken, the former are very apt to remain within the eye in spite of every effort to remove them. A few months later the sectors lose their sharp contour, break down, and finally disappear. We can then depend upon the exit of the whole cataract.

Yet in persons over sixty years of age, in whom the nucleus is usually large, many a cataract can be completely removed which does not come up to the strict standard of maturity just laid down; and, at that time of life, I would not hesitate to operate, without waiting for absolute maturity, if the patient were materially incommoded for want of sight.

The foregoing is the most common course of events in the progress of a senile cataract; but there is a rather rare form of it, in which total opacity of the cortical layers never does come about. In this form the lens is occupied by radiating linear opacities up to the very capsule; but between these opaque lines there are clear intervals, which may even admit of the fundus oculi being examined, although dimly, and which allow of a certain amount of sight. These cataracts can be successfully removed.

After the stage of maturity a cataract gradually goes on to be hypermature. Here one of two changes takes place: either the cortical substance breaks down, and becomes fluid, the nucleus retaining its consistency, and gravitating to the lowest part of the capsule (Morgagnian cataract); or, more commonly, the cortical substance dries up, as it were, and finally comes to form, with the nucleus, a hard flat disc. Accompanying these changes in the lens substance are changes in the epithelium lining the inner surface of the anterior capsule, which result in a thickening of the capsule. In a Morgagnian cataract the fluid cortex finally undergoes absorption, and the anterior and posterior capsules come in contact (cataracta membranacea). In some cases the capsule remains more or less transparent, and the sight may greatly improve. Some cases are on record of spontaneous cure of cataract, due to intracapsular absorption.

The investigations of Priestley Smith[1] have shown that a diminished rate of growth of the lens precedes the formation of cataract; and it is held that the cataractous process in the

senile lens is the result, in the first instance, of a rapid sclerosis and shrinking of the nucleus. If the process of sclerosis and shrinking be very gradual, cataract does not appear, because the cortical layers of the lens have time to accommodate themselves to the altered state of things; but if the shrinkage be rapid the cortical layers cannot so rapidly accommodate themselves, and then the fibrillæ of these layers become separated somewhat from each other, and fluid collects in the interspaces. This fluid it is which causes the disintegration of the lens substance, gradually leading to opacity of the whole lens. As the opacity increases, more fluid is present in the lens, and it is this which causes the swelling of the lens already referred to. When the whole cortex has become opaque the fluid contents begin to diminish, and the lens returns to its normal size. Senile cataract, then, is entirely a local process, and is not dependent on any disordered state of the general health.

The dimensions of the nucleus vary a good deal. In some cataracts it is very small, and these are called soft cataracts, as they consist chiefly of the soft cortical substance. In others—and as a rule in patients over sixty years of age—the nucleus is large, and these are called hard cataracts, although they are not hard throughout. The size of the nucleus can be estimated pretty accurately by the extent and intensity of the yellowish or brownish reflection, which is obtainable by focal illumination from the center of the cataract.

In some senile cataracts the sclerosis is not confined to the nucleus, but extends to the cortical layers as well. This causes much disturbance of sight, and the term cataracta nigra is given to these lenses, from their very dark hue, although they are not cataracts in the true sense of the term. They require operation, and, as they are always of large size, wide openings have to be made to deliver them.

In the lenses of young people there is no nucleus; consequently, in the complete cataracts of children and of young adults, there is no nucleus; the whole lens becomes opaque, and the cataract is always soft. Although the starting-point of cataract in children and young adults cannot be a shrinking of the nucleus, as there is none, yet the opacity is no doubt due to the taking up of fluid by the lens.

The natives of India are very prone to cataract, and at an early time of life, as are also persons in cold countries whose

business exposes their eyes for hours at a time to the heat and glare of furnaces—*e. g.*, glass-blowers, smiths, cooks, etc.

The Symptoms to which senile cataract gives rise consist, in the earliest stages, in the appearance of motes before the eyes and of monocular polyopia. Motes are complained of also in disease of the vitreous humor; but in those cases they float over a large portion of the field of vision, while in commencing cataract they occupy always the same relative position in the field. The polyopia is the result of irregular refraction in the media, which causes many images of the objects looked at to be formed on the retina. This symptom seems to annoy the patients more especially in the evening, when they look at gas or candle flames, the moon, etc. It is often complained of before there is any actual opacity in the lens, at a time when only the clefts filled with fluid between the fibrillæ can first be detected with weak transmitted light from the ophthalmoscope, as dark lines vanishing and reappearing according as the incidence of the light is altered.

In some cases of incipient cataract there is an increase in the refracting power of the lens, with the result that the patient becomes slightly myopic, if, previously, he have been emmetropic.

Gradually, as the opacity of the lens extends to other parts of it, the acuteness of vision becomes affected; and this is the more marked the more the cortex at the anterior and posterior poles of the lens is involved. In those cases where the equatorial parts of the lens are but little affected, while the polar regions are a good deal affected, the patients see better in the dusk, or with their backs to the light, than when their eyes are exposed to a strong light. The reason for this is that in the dusk the pupil is dilated, and light can pass through the clearer periphery of the lens, while in a strong light the pupil is contracted. On the other hand, when the opacity is confined rather to the equator of the lens, a strong light is not disturbing to sight; or, if the center of the lens be quite clear, a strong light may even be pleasant to the patient.

But, according as the lens becomes more and more opaque, the acuteness of vision is reduced, until, finally, even large objects cannot be discerned, and only quantitative perception of light is left. Some cataracts, however, when quite ripe, still admit of finger-counting at a few feet.

In advanced stages of the disease, as the opacities occupy a

great portion of, or the entire cortex, they are easily recognized even by ordinary daylight, often giving a grayish appearance to the pupil. Inflammatory exudation in the area of the pupil would afford a somewhat similar appearance, but would be attended by other signs of the previous inflammatory process, such as synechiæ, disorganization of the iris, etc., and it would be seen to lie more in the plane of the iris than does any lental opacity.

The length of time occupied by the ripening of a cataract varies in different cases from a few months to many years. In the very old the progress is, in general, more rapid than at an earlier time of life. That form which commences at the equator as fine lines is slower than that with flocculent opacities, or than that in which the cortex around the nucleus is likewise implicated at an early period.

All examinations as to the condition of the lens are rendered easier and more conclusive if the pupil be previously dilated with atropin; but the tension of the eye should be ascertained before atropin is instilled, lest glaucoma, or a tendency to it, be present.

Treatment.—No external local applications, nor internal medicines, are of any avail in the treatment of cataract at any stage. Removal of the cataract from the eye by operation is the only cure for blindness caused by it.

In cases of incipient cataract, or in those, rather, which have advanced somewhat beyond this stage, we often find that vision is improved, or made more pleasant, by the wearing of tinted glasses to moderate the light. With commencing cataract, where slight myopia has come on, low concave glasses for distant vision will be found of service; while, for reading, stenopeic glasses sometimes give good results. Yet, as a rule, patients are unwilling to use any of these aids.

Dilatation of the pupil with atropin is in many cases of the greatest benefit, especially where the nucleus is much more opaque than the cortical portion; but sometimes the diffusion of light resulting is most distressing to the patient, and greater impairment and confusion of vision are produced, and for this reason care in the prescription of atropin is demanded.

Patients with incipient or advancing cataract may, with immunity, be allowed to make every use they can of the sight they possess; and the surgeon should give them hints as to the arrangement of light in their rooms, and for their work, etc., so as to enable them to employ their sight to the best advantage.

The truly distressing period in the progress of cataract, when both eyes are affected, lies between the advent of that degree of blindness which incapacitates the patient for reading or writing, or for making his way about alone, and the occurrence of maturity, or of that degree of maturity which is deemed requisite for successful removal. This is often a lengthened time; it may be months or years. Fortunately, in many instances one cataract is much more advanced than that in the other eye; and then no such trial need be gone through.

Artificial Ripening.—In order to hasten the maturity of a cataract, puncture of its anterior capsule has been proposed and practiced with success, but has not been generally adopted, from the fear that it might set up iritis, or produce increased tension from excessive swelling of the cataract. Förster [2] effected artificial ripening by performing an iridectomy, which can afterwards be utilized for the extraction. This in itself often expedites the ripening, probably by disturbing the arrangement of the lens-fibers when the aqueous humor flows off, and the disturbance can be promoted by gently rubbing or stroking the lens through the cornea, immediately after the iridectomy, with the angle of a strabismus hook. This same massage of the crystalline lens may be employed with good result after simple tapping of the aqueous humor without iridectomy. Soon after this, a rapid increase in the opacity is often noticed, so that in from four to eight weeks extraction can be undertaken. The difficulty of this rubbing or massage of the lens lies in the estimation of the pressure to be applied; for if it be excessive the zonula may easily be ruptured, with the result of loss of vitreous when the extraction comes to be performed. The best results are obtained in cases of cataract where there is a firm and somewhat opaque nucleus, and where a certain amount of opacity already exists in the anterior cortical substance. I have occasionally employed the method, with satisfactory results; but some operators have seen iritis follow the proceeding.

The question whether the cataract in one eye should be extracted until both eyes are blind is often asked by patients. The answer is: A patient with one mature cataract, and the other progressing towards maturity, should have the ripe cataract removed. Hypermaturity is thus avoided, and also the stage of blindness above referred to. Again, if there be a ripe cataract in one eye, and not even incipient cataract in the other, it is often

advisable to operate for the purpose of increasing the binocular field of vision.

Complete Cataract of Young People.—The spontaneous occurrence of total cataract in the youthful lens is of rare occurrence, and its pathogenesis is still unknown.

Treatment.—Discission.

Diabetic Cataract.—This is a complete opacity of the crystalline lens occurring in diabetes, and due to disturbed nutrition. It has been proved by experiment that cataract can be produced by injecting solutions of sugar into the blood; but analysis of the aqueous humor in diabetic patients shows that the amount of sugar contained in it is not sufficient to account for the cataract. The cataract does not differ in appearance or consistency from other cataracts, according to the time of life of the patient.

Treatment and Prognosis.—Contrary to a very general opinion, these cases are not very unfavorable for extraction operations. I have operated on several cases of this kind, and always with success, save once, when the eye was lost by intra-ocular hemorrhage; and I have also seen such cases operated on successfully by others. There is no other method of restoring sight to these patients, who often live a long time. Ophthalmic surgeons of distinction inform me that occasionally patients operated on for diabetic cataract die of coma within about a fortnight or so after the operation; and they seemed to think that this was not diabetic coma of the ordinary kind, but coma caused in some way by the nervous system being upset by the operation. I have not had this untoward experience.

The operation of discission in these cases is apt to be followed by severe iritis.

Complete Congenital Cataract.—Children are sometimes born with crystalline lenses opaque in all their layers, while the other tissues of the eye are healthy. With congenital cataract defects of the chorioid or retina, or congenital amblyopia without ophthalmoscopic appearances, are also sometimes present, and these are usually indicated by nystagmus.

Treatment.—Discission.

Black Cataract.—This name, as above stated, is sometimes given to cases of extreme sclerosis of the lens, in which it assumes a dark brown color; but in other cases the lens is really black, the pigment being derived from the blood (hemin, or hematin). An instance has recently been observed in which the lens was

jet-black from this cause. The prognosis in these cases is not good, as they are often complicated with disease of the chorioid, or with hemorrhages in the vitreous humor.

Partial Cataracts.

These are nearly all congenital.

Central Lental Cataract.—This is a congenital and usually non-progressive form. It is an opacity of the central, or oldest, lens-fibers, while the peripheral layers remain clear.

Treatment.—Discission or iridectomy.

Zonular, or **Lamellar, Cataract.**—This is congenital, or forms in early infancy, and is the most common kind of cataract in children. It usually is present in both eyes, but it has been seen in one eye only. In it the very center of the lens is clear (Fig. 108), while around this is a cataractous layer or zone, and outside that again the peripheral layers are transparent. Most of these cases



Fig. 108.

are non-progressive, but occasionally the whole lens does become opaque, and usually then there have been previously some slight opacities in the otherwise clear cortical layers.

With oblique illumination the cortical layers of the lens are seen to be clear, while towards the center of the lens a uniform gray circular opacity will be observed. The diameter of this opacity may be small, perhaps not more than 3 mm. or 4 mm., or it may extend very nearly to the equator of the lens. If the pupil be dilated, and the lens examined with transmitted light, the cataractous portion will be seen as a more or less dark disc in the center of the lens, while all round it is seen the red light reflected from the fundus oculi. The center of this disc is either of the same degree of darkness as its margin or but very little darker; and this point serves to distinguish this form of cataract from one in which the whole center of the lens is opaque. In the latter case it is evident that the center of the opacity must be darker than its margin. In many cases small radial opacities

are seen round the equator of the lens, passing from the anterior to the posterior surface, their concavity embracing the circumference of the central opacity.

It is probable that lamellar cataract is due to some transient disturbance of nutrition in utero, occurring at the time the affected layers of the lens are being laid down. But against this view is the fact that one-half of the lens only may present the appearance of zonular cataract. The subjects of it are usually rickety, as shown by the irregular and imperfect development of the teeth, and by rachitic alterations in the bones of the skull. Convulsions during infancy in these patients are common.

The Treatment of central lental cataract and of zonular cataract is similar, and consists in either discission or iridectomy. The latter is very decidedly to be preferred in those cases in which the central opacity is so small that, on dilatation of the pupil, the acuteness of vision, with the aid of a stenopeic slit, is increased in a satisfactory degree. When the improvement is but slight, the breaking up of the lens with a needle is indicated. The advantage of iridectomy over discission, when the former can be adopted, is that no spectacles are afterwards required, and that the power of accommodation is retained.

Congenital cataracts may be needled any time after dentition is completed.

Anterior Polar, or **Pyramidal, Cataract** may be either congenital or acquired. In the former case it must be referred to some inflammatory disturbance occurring about the third period of development of the lens. In both cases the mode of origin of the opacity is the same, whether it be punctiform, flake-like, or pyramidal—namely, by contact of the lens with an inflamed cornea. In fetal life this may occur without any perforation of the cornea, as there is then no anterior chamber. After birth a perforating ulcer of the cornea is a necessary precursor of it, but the ulcer need not be central (p. 104). This contact with an inflamed and ulcerating cornea may lead to subcapsular cell-proliferation at that portion of the capsule which is exposed in the pupillary area, and consequent subcapsular opacity in this small area.

No *Treatment* is required, as vision is not affected.

Fusiform, or **Spindle-Shaped, Cataract** is also congenital, and is rare. It consists in an axial opacity extending from pole to pole, and may be combined with central or lamellar opacity.

The foregoing forms of cataract, with the exception, perhaps,

of the anterior polar and genuine black cataract, are primary; that is to say, they are not dependent on, or the result of, disease in other parts of the eye.

Secondary Cataract.

Some diseased states of the eye give rise to secondary cataract.

Of this a partial kind is

Posterior Polar Cataract.—This form is seen, with transmitted light, as a star-shaped or rose-shaped opacity in the most posterior layers of the posterior cortical substance, its center corresponding with the posterior pole of the eye.

Posterior polar cataract is usually found in eyes which are the subjects of disseminated chorioiditis, retinitis pigmentosa, or diseased vitreous humor. It sometimes progresses, and becomes a complete cataract; and then the prognosis for sight after extraction is not very good, owing to the disease which is present in the deep parts of the eye.

The additional disturbance of sight caused by the presence of posterior polar cataract depends a good deal upon its density.

Total Secondary Cataract often ensues upon contact of the lens with inflammatory products in the eye—*e. g.,* where false membranes have been produced by inflammation in the uveal tract. It is sometimes then called *Cataracta Accreta,* when the iris or ciliary processes are adherent to it. Cataract is also caused by detachment of the retina, intra-ocular tumor, absolute glaucoma, dislocation of the lens, etc. The reason of this is that the lens, in these cases, imbibes abnormal nutrient fluid from the diseased tissues with which it is in contact.

Such cataracts often undergo a further degeneration, and become calcareous. Calcareous cataracts are easily recognized by their densely white or yellowish-white appearance; and almost always indicate deep-seated disease in the eye, even when the functions, so far as they can be tested, are fairly good.

These secondary cataracts rarely come within the range of *Treatment,* as the diseases which give rise to them are usually destructive of sight. When, occasionally, they can be dealt with they should be extracted.

The term secondary cataract is also used in cases in which, after a cataract extraction, the capsule of the crystalline lens,

which is left behind, presents an obstacle to good sight. This will be referred to again further on, and is not to be classed with the conditions dealt with in this paragraph.

Capsular Cataract.

By this term is meant an opacity of the anterior capsule or of the capsular epithelium. It is usually confined to the center or anterior pole, and is most frequently seen in overripe senile cataracts and in secondary cataracts.

Traumatic Cataract.

Every injury which opens the capsule of the lens is liable to cause cataract, by reason of the admission of some of the surrounding fluids to the lenticular substance.

Perforating injuries with sharp instruments, or the entrance of small foreign bodies—in both cases, as a rule, through the cornea—are the most common injuries that produce traumatic cataract. But blows upon the eye, without any perforating wound, also, although rarely, produce cataract. In these latter cases there is a rupture of the capsule, either at the equator of the lens, or on its posterior or anterior surface.

Within a few hours after a perforating injury of the anterior capsule, the lens substance in the immediate neighborhood of the opening becomes opaque, swells, and protrudes as a gray fluffy-looking mass, through the opening in the capsule into the anterior chamber, where it gradually breaks up, dissolves, and becomes absorbed. It is immediately followed by other portions of the lens which have become cataractous, until, after a time, the whole lens will have disappeared, and the pupil again become black. Marcus Gunn suggests [3] that the explanation of the solution of the cataract in the anterior chamber consists in the fact that globulin is normally soluble in a weak solution of chlorid of sodium, such as we have in the fluid of the anterior chamber. The absorption of a traumatic cataract takes many weeks; and ultimately the eye sees well if a suitable convex lens be put before it.

But the course of events just sketched is the most favorable one, and is hardly likely to take place in a case which is wholly untreated. In the first place the swelling of the lens—especially if it be rapid, in consequence of a wide opening in the capsule—

is liable to irritate the iris, and to cause iritis; or to push the periphery of the iris forwards against the periphery of the cornea, block the angle of the anterior chamber, and cause secondary glaucoma.

Moreover, violent plastic or purulent uveitis may come on, as the consequence of the introduction of infective matter on the perforating object, or foreign body, which causes the cataract. Where this occurs the case enters into the category of diseases of the uveal tract; and the cataract, as such, becomes a minor consideration.

Again, we sometimes meet with traumatic cataracts which do not undergo any absorption process, but simply remain stationary; or, in the course of years, undergo secondary changes similar to those which occur in senile cataract. In these instances the trauma is usually a blow on the eye, not a perforating injury; and it is believed that the rupture of the capsule closes soon after the blow, and hence no lens matter can escape into the anterior chamber; also, the rupture in many of these cases is probably at the equator of the lens, where the aqueous would not readily get access to the lenticular substance.

Where the cataract is produced by a small foreign body flying through the cornea and into the lens, it is a matter of importance, for the prognosis, to decide whether the foreign body be in the lens or have passed through it into the deeper parts of the eye. In the former case we may hope to extract it with the cataractous lens; while in the latter case we must fear that it will set up dangerous inflammatory reaction. In such cases the lens should be well searched with focal illumination, and the transmitted light may also be of use; but it must be remembered that in these traumatic cataracts there are often glittering sectors in their deep parts, which may readily be mistaken for a metallic foreign body. If the foreign body be of steel or iron, the sideroscope may be employed for its detection, or the Röntgen rays may be utilized.

Treatment.—The pupil should be kept dilated with atropin, in order to draw the iris out of the way of the swelling lens matter; and nothing more is necessary if complications do not arise. But should iritis or high tension come on—and the surgeon must constantly test the tension—it is important, without further delay, to extract as much as possible of the cataract. This may be done either without an iridectomy, through a linear incision some 10 mm. long in the upper third of the cornea,

or with an iridectomy, through an incision in the upper margin of the cornea.

If a foreign body be present in the lens, extraction of the latter with the foreign body should invariably be undertaken.

Where violent purulent or plastic uveitis is set up by the trauma, the treatment resolves itself into that for these inflammations.

Operations for Cataract.

With regard to the *State of Health of the Patient* about to be operated on it is desirable, as in every operation, that it should be good. Still, we have so often in these cases to deal with very old people, that we cannot in every instance require sound organs and a robust constitution; and, as a matter of experience, I have not found serious disease of the heart, lungs, and liver, even when they all existed in the same individual, any impediment to a successful operation. Diabetes is no absolute contra-indication, and even in the presence of Bright's disease I have operated successfully. Very advanced years form no obstacle. I have frequently operated for cataract on persons over eighty years of age, and always with success.

The State of the Eye itself should be carefully investigated prior to proposing or undertaking an operation for cataract, and is a much more important matter than the general health. Above all things, it is to be determined whether there be intra-ocular complications, which would neutralize the result of a successful operation, such as detachment of the retina, disseminated chorioiditis, atrophy of the optic nerve, etc. The examination of the eye before the lens has become opaque, if the surgeon have had that opportunity, will be the most reliable basis upon which to go; and for this reason a careful note should be taken of the condition of the fundus in each case of incipient cataract. The examination of the fundus of the other eye, if its lens be clear, may help in determining the point in so far as those intra-ocular diseases are concerned which are apt to be binocular. Again, the condition of the anterior capsule of the lens should be observed, for a defined glistening white square patch, about 2 mm. broad, situated in the center of the capsule, tells the tale of intra-ocular mischief. It cannot be confounded with the more diffused striated and punctated capsular alterations due to overripeness.

Finally, the functions of the eye should be examined. With

an uncomplicated cataract of the most opaque kind good perception of light should be present, so that the light, say, of a candle some two meters distant may be distinguished. In less dense cataracts fingers may be counted at 1 m. or 1.5 m. when full maturity has been attained. The field of vision must be examined by means of the "projection of light"—*i. e.*, a lighted candle held in different parts of the field should be recognized by the patient, who is required to point his finger in the direction of the light, as it is moved rapidly from one part of the field to another. This examination can also be made by means of the light reflected from the ophthalmoscope mirror. If the patient fail to project the light in any direction, a diseased condition in the corresponding part of the retina may be suspected. Yet in cases of very old uncomplicated cataract the patients often project the light in some one direction, no matter where it may come from. A certain degree of intelligence on the part of the patient is required for this test.

By the foregoing means most intra-ocular complications of a serious nature can be detected; but there is at least one against which I know of no safeguard, namely, a small circumscribed spot of chorioido-retinal degeneration at the macula lutea (central senile chorioiditis). After removal of a cataract from an eye affected in this way, the patient's vision is so much improved as to enable him to go about alone, but reading will still remain an impossibility for him.

The Cornea should be Examined.—Such corneal opacities as would seriously compromise vision may contra-indicate the operation; but slighter opacities, discernible only with oblique illumination, would merely diminish the future acuteness of vision, and would require a corresponding prognosis to be given before operation.

The Condition of the Appendages of the Eye, too, must be examined. Should there be any conjunctivitis, blepharitis, or dacryocystitis, it ought to be cured or alleviated before the operation is undertaken. Very successful operations, it is true, may be performed in the presence of chronic dacryocystitis, or of granular ophthalmia; but the risk is great, and it is in all respects wiser to reduce their activity to a minimum. Some surgeons, in cases of dacryocystitis, temporarily obliterate the lacrimal puncta by introducing a red-hot needle. A better means is extirpation of the lacrimal sac.

Extraction of Cataract.

Linear Extraction.—The extraction through a linear incision in the cornea is applicable only to soft, or fluid, cataracts, in persons under the age of twenty-five. The instruments required are: A spring lid elevator (Fig. 109), a fixation forceps, a wide




Fig. 109.

keratome (Fig. 110) or a Graefe's cataract knife, a cystotome (Fig. 111), and a Critchett's vectis (Fig. 112).

The speculum having been applied, a fold of conjunctiva close to the margin of the cornea, and at the inner end of the horizontal



Fig. 110.



Fig. 111.



Fig. 112.

meridian of the latter, is seized (Fig. 113) with the fixation forceps, and the eye fixed by it throughout the operation. The point of the keratome is now entered into the cornea in its horizontal meridian, about 4 mm. from its outer margin, and is passed into

the anterior chamber; or, the incision may be made in the upper part of the cornea. The blade of the knife is then laid in a plane parallel to that of the iris, and pushed on until the corneal incision has attained a length of 6 or 7 mm. The point of the knife being now laid close to the posterior surface of the cornea—in order that no injury may be done to the iris or lens




Fig. 113.

when the aqueous humor commences to flow off—the instrument is very slowly withdrawn, so that the aqueous humor may come away gradually, without causing prolapse of the iris. In withdrawing the knife it is well to enlarge the inner aspect of one or other end of the wound, by a suitable motion of the instrument in that direction.

The keratome being now laid aside, the cystotome is passed




Fig. 114.

into the anterior chamber (Fig. 114) as far as the opposite pupillary margin, care being taken, by keeping the sharp point of the instrument directed either up or down, not to entangle it in the wound or in the iris. The point is now turned directly on the anterior capsule, and, by withdrawing the cystotome towards the corneal incision, an opening in the capsule of the width of the

pupil is produced. The cystotome is then removed from the anterior chamber, with the same precautions as on its entrance.



Fig. 115.



Fig. 116.



Fig. 117.



Fig. 118

The edge of the vectis is then placed on the outer lip of the corneal incision, and the latter is made to gape somewhat,

gentle pressure being at the same time applied to the inner aspect of the eye by the fixation forceps, and in this way the lens is evacuated. When the pupil has become quite black the operation is concluded. If pressure does not at first clear the pupil completely, the speculum should be removed, the eyelids closed, a compress applied, and a few minutes allowed to elapse, in order that some aqueous humor may be secreted. A renewal of the efforts to clear the pupil will probably now be successful, or, if not, another pause may be made, and then fresh attempts employed until the pupil is quite clear. It is unwise to insert the vectis into the eye to withdraw the fragments; and if some of these should be left behind, no ill results need necessarily follow, although iritis is more apt to supervene than if the lens be thoroughly evacuated. Fragments left behind become absorbed. If there be a prolapse of the iris which cannot be reposed it must be abscised.

Von Graefe, Waldau, and Critchett endeavored, by increasing the size of the incision, placing it in the corneo-sclerotic margin, performing an iridectomy, and introducing a vectis for delivery of the cataract, to make the linear extraction applicable to senile cataracts. The successes derived from these modifications were not, however, more satisfactory than those obtained from the old Flap Operation. But these experiments led von Graefe to the operation, a modification of which is now very generally employed. He called his operation

The Modified Peripheral Linear Extraction.—The instruments required are: A wire lid-speculum, a fixation forceps with spring catch, a von Graefe's cataract knife (Fig. 115), a curved iris forceps, an iris scissors or a de Wecker's forceps-scissors (Fig. 116), a bent cystotome, a hard rubber spoon (Fig. 117), and a hard rubber, tortoiseshell, or silver spatula (Fig. 118).

Before proceeding to operate, the eye is thoroughly cocainized by the instillation of about three drops of a 2 per cent. solution of hydrochlorate of cocain, at intervals of two or three minutes.

Antiseptic Measures, similar to those used for the Three Millimeter Flap Operation (*vide infra*), are to be carefully attended to.

The Operation.—The speculum having been applied, the eye is steadied by seizing a fold of conjunctiva with its subconjunctival tissue, close to the lower margin of the cornea, and in a prolongation of the vertical meridian of the latter. The eye is

now drawn gently downwards, the patient assisting in the motion. The point of the Graefe's knife, its cutting edge being directed upwards, is then entered into the corneo-sclerotic margin at a point (*A* in Fig. 119) about 1.5 m. from the outer and upper corneal margin, and 2 mm. below the level of the tangent which would pass through the highest point of the corneal margin. The blade is held in a plane parallel to that of the iris, and is pushed on into the anterior chamber until its point reaches the point *C*, some 7 or 8 mm. of the blade being now in the anterior chamber. The handle of the knife is then lowered, so that the point of the blade is brought up to *B*, where it is made to pass out through the corneo-sclerotic margin, this counterpuncture corresponding in position, with reference to the corneal margin, to the point of entrance *A*. The edge of the knife is now turned slightly forwards, and by one or two sawing motions the incision *A B* is completed in the corneo-sclerotic margin. The blade still lies under the conjunctiva, which is divided, the edge



FIG. 119.

of the instrument being turned more forwards, or even somewhat downwards, as it is not desirable to have too large a conjunctival flap.

The advantage of this incision lies in its peripheral position, which is almost in the plane of the crystalline lens, and, consequently enables the cataract to be delivered without revolution on its axis. At a later period von Graefe altered the incision, so that, puncture and counterpuncture lying as described, the center of the incision passed through the apex of the clear cornea instead of through the corneo-sclerotic margin. This, by making the incision more nearly a segment of a greater circle of a sphere, made it as linear as possible, and consequently, in his opinion, its margins adapted themselves more readily.

The next step in the operation is an iridectomy, a portion of iris corresponding to the whole length of the wound, or nearly as much, being excised. This iridectomy is necessary, or advisable, chiefly because of the peripheral position of the wound, which

would render prolapse of the iris very liable to occur; but it also facilitates the delivery of the lens and cortical masses. The subsequent stages—capsulotomy and delivery of the lens—are similar in their details to those in the Three Millimeter Flap Operation, to be presently described.

It was found that the advantages of the position and form of the incision in this procedure were largely counterbalanced by the danger of prolapse of the vitreous, the difficulty of proper reposition of the angles of the coloboma, and the liability to cyclitis, all entailed by the peripheral incision, and consequently this incision has been abandoned by nearly all operators.

Out of this method grew that one which is known as the

Three Millimeter Flap Operation, or, more commonly, **The Combined Operation** (*i. e.,* combined with an iridectomy), first proposed by de Wecker. I shall describe the operation as I am in the habit of performing it; and I may here say that for success in the cataract operation it is necessary, not only to select the method which seems the most rational, but also to devote the utmost attention to a series of minute details in its performance.

Preparation of the Patient.—A gentle purgative is given the day before the operation, so that the bowels need not be disturbed for two days after the operation. The face is washed with hot water and soap shortly before the operation.

Preparation of the Eye.—Half an hour before the operation a drop of a sterilized 2 per cent. solution of sulphate of eserin is dropped into the eye, and this is repeated a quarter of an hour later. Just before the operation, at intervals of two minutes, three drops of a sterilized 2 per cent. solution of hydrochlorate of cocain are dropped into the eye. Finally, the lids having been everted, the conjunctival sac is washed out with sterilized physiological solution of common salt, particular attention being paid to the fornix of each lid, and to the inner and outer canthus. Then the skin of the eyelids and immediate surroundings of the eye are freely washed with the same solution.

Preparation of the Instruments.—The instruments required are the same as those for the Modified Linear Extraction. Immediately before the operation they are sterilized by boiling; they are then plunged for a moment into absolute alcohol, and laid on a sterilized porcelain tray, under a sterilized cloth, until required for use.

During the Progress of the Operation small bits of lint, wet

with the sterilized salt solution, are employed to wipe away coagula, cortical masses, etc., and are not employed a second time. An assistant should place the instruments in the surgeon's hand in their turn, and take out of his hand those he has used, in such a manner as to render it unnecessary for him to look away, even for a moment, from the field of operation.

The Operation.—A spring wire lid-speculum is applied. The eye is fixed with a catch fixation-forceps by a fold of conjunctiva and subconjunctival tissue, below the vertical meridian of the cornea, or a little to one side of this line (Fig. 120).

The point of the knife is entered just in the margin of the clear cornea, at the outer extremity of a horizontal line which would pass 3 mm. below the summit of the cornea. This line is easily found by placing the knife, which is about 2 mm. broad, horizontally across the cornea, so that a margin of clear corneal tissue 1 m. broad may remain exposed between the knife and the summit of the cornea. The knife is then passed cautiously through the anterior chamber, and the counterpuncture is made in the corneal margin at the inner extremity of the horizontal line described, and the incision is then finished in the corneal margin by a few slow to-and-fro motions of the knife.

Owing to the action of the eserin, the iris does not prolapse. The incision, between puncture and counterpuncture, lies in the clear cornea at its very margin, as represented by the dotted line in Fig. 120. This incision is no longer linear, but slightly curved. It is found, however, to adapt itself readily, and, being less peripheral than the true von Graefe incision, the objections to the latter are obviated.

The Second Stage of the Operation consists in an Iridectomy. The fixation of the eye having been given over to the assistant, the iridectomy is performed by passing a curved iris forceps into the anterior chamber, seizing the smallest possible portion of the sphincter of the iris at a point corresponding to the center of the incision, drawing it out, and with the forceps-scissors excising a very small central bit of iris. This is done either by making two snips in the iris, one at either side of and close to the forceps, each of them reaching to the periphery of the iris, and then a third cut which joins these two at the base; or, the forceps-scissors being approached from over the cornea—*i. e.*, at right angles to the wound—the coloboma may be formed with one snip of the instrument, and, if care be taken to keep the blades close to

the forceps, a narrow, neat coloboma may thus be obtained. It is unnecessary to excise a large portion of iris, although in von Graefe's original operation a portion corresponding to the entire length of the wound used to be taken away. A small coloboma, say of 2 mm. to 3 mm. in width, as in Fig. 100, is sufficient to allow of an easy delivery of the lens by doing away with the resistance of the sphincter iridis, and to prevent secondary prolapse of the iris (*vide infra*); and its advantages over a wide iridectomy, from an esthetic point of view, are obvious. It is always, therefore, my object to obtain the smallest possible coloboma. The procuring of a neat coloboma is much facilitated if, prior to the operation, the pupil has been contracted (see Fig.




Fig. 120.

120) by the instillation of one or two drops of solution of sulphate of eserin, as above recommended.

The Third Stage of the Operation is the Capsulotomy. The operator takes the fixation forceps from his assistant, who then raises the speculum and eyelids slightly off the globe, in order that no pressure may be exerted on the latter during the remainder of the operation. The surgeon, passing the cystotome into the anterior chamber, divides the anterior capsule of the lens by two incisions, each from the lower pupillary margin upwards, one directed outwards, the other inwards, as far as the anterior surface of the lens can be seen, while finally a third incision is made along the upper periphery of the lens. An extensive opening in the capsule is of great importance, as otherwise difficulty in delivery of the lens may be experienced, and because a small opening renders the occurrence of secondary cataract more likely. In

dividing the capsule it is important not to dig into the lens, as this, in the case of a hard cataract, is apt to dislocate it. A rather oblique application of the cystotome to the capsule is, for this reason, the best.

The cystotome often drags a tag of the capsule into the corneal wound, where it lies until the end of the operation, and where, owing to its transparency, it may easily pass unnoticed. Such a tag acts as a foreign body, and may subsequently form the starting-point of troublesome complications.

Capsule forceps have been invented for the purpose of taking away a large portion of the anterior capsule; but this does not altogether obviate the danger of capsule in the wound, nor does it do away with the likelihood of secondary cataract. I have no objection to the method, but it does not seem to have any advantages over that just described in cases where the capsule is not thickened. But when the anterior capsule is thickened it is always desirable to tear away a central portion of it with forceps.

Gayet of Lyons [4] and Knapp of New York [5] have proposed a method of opening the capsule termed peripheral division—*i. e.*, they make only one opening in the capsule at the upper periphery of the lens with a very sharp needle cystotome, which is passed along the whole length of the corneal section, a wide iridectomy having been made for this purpose. The chief advantages claimed for this method are: Safety from a tag of capsule in the wound, and safety from iritis caused by irritation from particles of lenticular substance left behind after delivery of the lens. On the other hand, it has the disadvantages of the secondary operation on the capsule, which becomes necessary in a much larger proportion of the cases than where a free central opening is made.

The Fourth Stage is the Delivery of the Cataract. The eye is drawn gently downwards—the patient being called on to assist in this motion by looking towards his feet; the convex edge of the hard rubber spoon is placed just below the lower edge of the cornea, and gentle pressure is exercised on this place, the pressure to be gradually increased until the upper margin of the lens presents itself in the wound, when, the same pressure being maintained, the spoon is advanced over the cornea in an upward direction, pushing the lens before it and out through the wound. As soon as the greatest diameter of the lens has passed the wound the pressure of the spoon should immediately be diminished, lest rupture of the zonula be caused. The fixation-forceps and

speculum are now removed from the eye, and a cold sterilized compress is laid on the closed lids.

It may be noted that the fixation-forceps and the speculum are used until this late stage in the operation is reached. Probably most operators do likewise. Some, however, use neither fixation-forceps nor speculum from beginning to end of the operation; while others discard the fixation-forceps when the corneal section is completed, but retain the speculum until after the iridectomy only, delivering the lens with the finger placed on the lower lid. I think that the use of the fixation-forceps and the speculum until after the lens is delivered gives more security and stability to the operator than the other methods, and I see in it no counter-disadvantages.

The Fifth Stage consists in Freeing the Pupil of any Cortical Masses which may have been rubbed off in the passage of the lens through the wound, and in what is called the Toilet of the Wound.

The presence of cortical remains is recognized by the pupil not having become quite black, or by the vision not being such as it ought to be (fingers counted at several feet), or by inspection of the cataract just removed showing that some portions of it are left behind. The use also of focal electric illumination for the detection of cortical fragments is very advantageous. If any fragments be present, the cold sterilized compress having lain on the eye for a few minutes to enable some aqueous humor to collect, the operator, facing the patient, raises the upper lid with the thumb of left hand, while, with the first and second fingers of the right hand laid on the lower lid, light rotatory motions are made with this lid over the cornea so as to collect the masses towards the pupil, and then a few rapid light motions upwards, with the margin of the lid, drive these masses towards, and out of, the wound.

Care and delicacy of touch are required in order to perform this lid-maneuver successfully, without rupturing the hyaloid by undue pressure.

With an iris-forceps the blood-clots which may adhere to the wound are now removed.

I then employ the following means, to prevent the possibility of any portion of capsule being incarcerated in the wound during healing: A bent iris-forceps is passed open between the lips of the wound, closed, and drawn gently out again. Frequently

a tag of capsule will have been captured by the forceps, and is snipped off with the scissors, or it may be that no capsule is caught. The forceps is then similarly inserted at an adjacent part of the wound; and in this manner the wound is searched from end to end for capsule. In about 25 per cent. of the cases a tag of capsule is found present. I regard this measure as an important one, for I believe that it effectually removes the one serious drawback to the valuable operation under consideration.

Finally, the coloboma has to be seen to. The peripheral portions of the iris corresponding to the ends of the wound are apt to have become prolapsed in the course of the operation, and to have displaced the angles of the coloboma upwards. If this be not corrected the prolapsed portions of the iris heal in the wound, and cause bulgings there later on, the pupil in the course of some months becoming drawn up towards the cicatrix. Hence, in every case, even where everything seems to be in order, it is important to pass the narrow spatula into the anterior chamber, and to gently stroke down each pillar of the coloboma as far as it can be brought. The instillation of eserin before the commencement of the operation will cause the sphincter iridis to assist in producing the desired result. All this is aptly termed the toilet of the wound.

The sight of the eye should then be tested by finger-counting, as this affords the patient satisfaction, and lends him courage for the next few days of strict quiet. Finally, the conjunctival sac is flooded with the sterilized saline solution.

The dressing is now applied. A piece of dry sterilized lint, sufficiently large to extend 1-4 inch beyond the orbital margin in every direction, is laid on the closed eyelids. Pieces of sterilized absorbent cotton-wool are laid on this, the hollows at the inner canthus, etc., being carefully filled up; so that, when the bandage is put on, it may exert equal pressure on every part of the eye. I apply three turns of a narrow roller bandage over the dressing and round the head, in the manner which was customary in von Graefe's clinique; but various other, and doubtless equally good, forms of bandage are in use. The pressure of the bandage need only be sufficient to maintain the dressing firmly in its place. It is usual to keep the other eye closed by a light bandage.

I am opposed to the after-treatment of cataract operations without bandage, as advocated by some surgeons. It is by no

means a new method, and I do not doubt that many cases recover under it. I do not believe, however, that in a long series of cases the same percentage of recoveries can be obtained by it as with the bandage.

Accidents Liable to Occur during the Operation.—The Wound may be made Too Small, and the delivery of the lens, consequently, may be so difficult that the margins of the wound become contused, and consequently suppuration may be promoted. The zonula, too, may be ruptured by the excessive pressure, from efforts to force the lens out through the narrow aperture, and prolapse of the vitreous may ensue. If the directions above given be carefully attended to, the vast majority of both hard and soft cataracts may be extracted without difficulty; but should the wound be made too small, it can best be enlarged by the forceps-scissors, or a blunt-pointed knife made for the purpose. Where the presence of an unusually large hard cataract is diagnosed, it is important to make the incision larger *ab initio,* by placing puncture and counterpuncture nearer to the horizontal meridian of the cornea than above directed.

Hemorrhage into the Anterior Chamber may take place. It may be from the iris, from the corneo-sclerotic margin, or from the conjunctiva. Pressure with the spatula on the cornea, which causes the wound to gape, is often successful in clearing the chamber of blood, which might interfere with accurate division of the capsule. Yet, when this cannot be completely got rid of, the capsulotomy can be performed by the exercise of greater care. Cocain, by its power to contract the blood-vessels, has rendered this hemorrhage a less common complication than it used to be.

Prolapse of the Vitreous Humor.—This may be due to a too peripheral position of the wound, support being thus taken away from the zonula, and the danger of its occurrence was a disadvantage of the completely corneo-sclerotic wound practiced at one time by von Graefe. The Three Millimeter Flap Operation is less liable to be attended with loss of vitreous. This accident may also be caused by undue pressure made on the eyeball by the speculum, fixation forceps, or spoon, or by the lower lid during the lid-maneuver. It may be due to defective zonula with fluid vitreous humor. When the vitreous prolapses prior to delivery of the lens, the latter falls back into the eye, and can only be delivered by at once drawing it out with a Critchett's, Taylor's, or

other suitable vectis; and this may be regarded as one of the most serious accidents which can occur in the course of the operation. Loss of vitreous after delivery of the lens is less serious; indeed, a considerable portion of the vitreous may then be lost without ill result to the eye; yet it increases the traumatism, and renders inflammatory reaction more liable to occur. Opacities in the posterior chamber of the eye are frequently an ultimate result of loss of vitreous; but a much more serious consequence is sometimes seen in detachment of the retina.

Normal After-Progress.—Soon after the completion of a normal operation, the effect of the cocain having passed off, some smarting commences, and continues for four or five hours. After that time the patient has no unpleasant sensation in the eye, unless it be some itching, or a slight momentary pain or sensation of a foreign body, especially when the eye is moved under the bandage. The first dressing is made in forty-eight hours, in a manner similar to that immediately after the operation, a drop of atropin being instilled, as also at each successive dressing; and the sterilized salt solution is used for freely washing the margins of the eyelids, some of it being allowed to trickle into the conjunctival sac. At this first dressing it is well to abstain from a very minute or lengthened examination of the eye; but, if the lid be gently raised, the wound will be found closed, the cornea clear, the anterior chamber completely restored, and the pupil semi-dilated and black. The subsequent dressings are made night and morning, for the purpose of instilling atropin. On the third day after the operation the patient may be allowed to sit up, the room being kept moderately dark; and on the fifth or sixth day the bandage may be left aside permanently, and dark glasses worn in its stead. In the course of a few days more the patient, having been gradually used to more light, may be allowed out of doors. It is desirable to continue the use of atropin for about a fortnight longer, or until all abnormal vascular injection of the white of the eye has disappeared, as until then there is danger of iritis. (For selection of glasses in aphakia see end of this chapter.)

Irregularities in the Process of Healing.—The pain may continue longer than four or five hours, and it is then well to quiet it by a hypodermic injection of morphia in the corresponding temple. Should severe pain come on some hours later it is apt to be due to an accumulation of tears under the eyelids, and it

immediately subsides on the bandage being removed and exit given to the tears by slightly opening the eye.

Late Appearance of the Anterior Chamber.—At the first dressing it will sometimes be found that there is no anterior chamber, although the appearance of the wound is quite satisfactory; but this need occasion no alarm, as the anterior chamber is sometimes not restored for a week. Should a more lengthened absence of the anterior chamber be noticed, it may be due to the presence of a small tag of capsule in the wound, and it is then desirable to search the latter with a forceps, and to cut off any capsule which may be found there.

Striped Keratitis.—At this dressing also it may sometimes be observed that there is a more or less well-marked striated cloudiness of the cornea, extending over nearly the whole of it, or occupying only a part in the immediate neighborhood of the wound.

This opacity is, according to some, the result of injury to the endothelium of the posterior surface of the cornea during the operation by instruments, or by the chemical action of an antiseptic lotion, when such a lotion has been used. Leber has shown that the entrance of even the aqueous humor, through a loss of substance in the endothelium, is sufficient to cause the fibers of the true cornea to swell and become opaque, just as the crystalline lens is acted on if its capsule be opened. The endothelium of the posterior surface of the cornea in fact it is which protects the latter from being infiltrated by the aqueous humor.

The explanation given by Hess,[6] however, seems a very reasonable one, namely, that it is due to folding of the posterior layers of the cornea, on account of the difference in tension in the vertical and horizontal direction. His conclusions are based on microscopic examination and experiment.

This so-called striped keratitis is, for the most part, of no serious import, as it usually passes away in a few days, and leaves the cornea perfectly clear; and folding of the posterior layers would account for these cases. But now and then cases do occur in which the process is very intense, and where a permanent white opacity remains in the cornea over the pupillary area, with consequent serious deterioration of vision. These severe cases are most apt to be caused by the introduction of an antiseptic solution into the anterior chamber; for the chemical action of the antiseptic on the corneal tissues is more damaging, and therefore the opacity it produces more permanent, than is the action of the

aqueous humor. Sublimate lotion is the antiseptic which has been most often to blame, probably because it is the antiseptic in most general use. With the 1 in 5000 solution which I at one time employed I never had the severe form, and rarely the mild form; but then I never deliberately introduced the solution into the anterior chamber. I have had only one case of the severe form, and in it, by mistake, a sublimate lotion of 1 in 2500 was used for irrigation of the surface of the eye. At a later period I employed a solution of only 1 in 10,000, but this I have abandoned in favor of the sterilized physiological solution of salt, as I have stated.

Suppuration of the Wound.—This is a danger which is very much rarer than it was prior to the introduction of aseptic surgery; indeed, it is almost banished from the cataract operation. When it occurs it usually does so between the twelfth and thirty-sixth hour after the operation, rarely earlier or later, and is a very serious event; for in the vast majority of cases, do the surgeon what he may, it leads to loss of the eye. Its onset is made



Fig. 121.

known by severe pain of a continuous aching kind in and about the eye; and is thus easily distinguished from the slight, short, stabbing pain, with long intermissions, which some patients complain of, and which has no evil import. On removing the bandage the eye will be found full of tears, and the wound covered with a layer of muco-pus, which can be removed with the forceps in one mass, while the aqueous humor and cornea may already present some opacity. In some hours more the corneal opacity increases considerably, the iris becomes distinctly inflamed, and the pupil filled with a mass of inflammatory exudation. The inflammatory process may remain confined to the wound and iris, and when, in the course of some weeks, it entirely subsides, it leaves the pupil drawn up towards the wound, so that an appearance as in Fig. 121 is presented; or, the inflammation may strike into the ciliary body and chorioid, and produce purulent panophthalmitis, with total destruction of the eye.

To combat Suppuration the best method is the immediate cauterization of the corneal wound in its whole extent with the

galvano-cautery. Also, the wound may be opened up from end to end with a spatula, the aqueous humor evacuated, and the anterior chamber washed out with injections of sublimate solution 1 in 10,000, while the conjunctival sac is irrigated with the same solution. If necessary, these measures are to be repeated at intervals of eight or ten hours. Subconjunctival or intra-ocular injections of sublimate may also be tried.

Iritis.—Apart from the iritis which occurs in connection with suppuration of the wound, this complication is most usually due to irritation from masses of cortical lens-substance left behind, or to infection during the operation, which can show itself in this way instead of by suppuration. Iritis does not usually come on for some days after the operation. It is ushered in with the usual symptoms of pain, and is generally of the plastic variety. General plastic uveitis may be set up, and sympathetic ophthalmitis may result. *Treatment* consists in strict confinement to a dark room, atropin, warm fomentations, leeching, and, internally, salicylate of soda is most useful. In these cases vision is liable to be damaged by pupillary exudation, which remains as a permanent obstruction to vision.

Detachment of the Chorioid.—Fuchs [7] has pointed out that detachment of the chorioid occurs some days after cataract extraction, in some of the cases in which the anterior chamber does not form, or in which, having formed, it becomes empty again. The detachment occurs, too, occasionally after iridectomy unconnected with cataract extraction. It can be seen with the ophthalmoscope, and sometimes even with the focal illumination. Vision while the lesion is at its height is seriously affected, but the prognosis is good, for the detached portion always becomes reposed.

Cystoid Cicatrix.—After convalescence, all foregoing dangers having been escaped, the cicatrix in the corneal margin sometimes bulges and becomes semi-transparent, presenting the appearance of a vesicle, and may attain a large size. The extremities of the late incision are the most common positions for this condition, but it may occupy the entire length of the cicatrix. It does not generally come on for some weeks, or more, after the operation. In some cases it is caused by a tag of iris which is incarcerated in the wound; but in other cases by a small piece of capsule, which has similarly healed in the wound. Irregularity in curvature of the cornea, and consequent irregular astigmatism,

are the least of its evil consequences. If the condition be caused by incarceration of iris, the pupil will be gradually drawn close to the upper corneo-sclerotic margin; while, if it be caused by a portion of capsule, iridocyclitis may be produced. Whether the iris or the capsule be the cause, these eyes are always exposed to the danger of a sudden onset of purulent iridochorioiditis (see p. 263). All this demonstrates the immense importance of attention to those details of the operation which are calculated to obviate incarceration of iris, or of capsule, in the cicatrix.

Cataract Extraction without Iridectomy, or more commonly, **The Simple Operation.**—This method is older than the Linear, von Graefe's, or the Three Millimeter Flap Operation, and used to be known as the Flap Operation. It has been revived within recent years by many distinguished operators. It differs from the Three Millimeter Flap Operation in that the incision occupies a greater extent (about one-third) of the circumference of the cornea, and that no iridectomy is made. Formerly the knife used was triangular in shape (Beer's knife), but von Graefe's cataract knife is the instrument now employed. The round pupil, and consequent somewhat prettier appearance of the eye, is the one advantage which can be claimed for this procedure over the Three Millimeter Flap Operation as above described; for vision with a circular pupil is not better than where a small iridectomy has been made. As a set-off against the circular pupil, the extraction without iridectomy exposes the eye to the serious danger of prolapse of the iris into the wound. Those who operate after this method make it a rule to perform an iridectomy in all cases where they cannot satisfactorily repose the iris after delivery of the lens; but, even where they can repose it well, they are not secure against the occurrence of a prolapse within the first two or three days after the operation; nor do they find that eserin, or any other means, provides the desired safeguard. It is admitted that prolapse of the iris takes place after a number of these operations, and that there is no means of foretelling in what eyes it will occur. The prolapsed portion of iris heals in the wound, which then, in a few weeks, becomes more or less cystoid and bulging, causing displacement of the pupil and irregular curvature of the cornea, with resulting deterioration of vision. Nor is this all; for such eyes are liable—weeks, months, or even years after the operation—to take on severe iridocyclitis, ending in total loss of sight. Another disadvantage of this

operation is, that removal of cortical remains cannot be so effectually performed as where a coloboma has been made.

Therefore, while admitting the charm of a circular pupil, I am of opinion that the question is not whether the appearance of some of the eyes operated on is pleasing to us and to others who inspect them, but rather what advantage the greatest number of persons operated on derive from the operation. With sentimental talk about "mutilation" of the iris I do not sympathize.

The explanation why, in the simple extraction, prolapse of the iris with subsequent incarceration is more liable to occur, even some days after the operation, than in the combined operation, and why it is so difficult to devise a sure means for preventing the accident, as, also, how it is that even a very narrow coloboma is, almost always, sufficient to protect the eye from this disaster, is the following: Within a few hours after the operation the wound in the corneal margin commonly closes, the aqueous humor collects, and the anterior chamber is restored. But it takes many hours more for the delicate union of the lips of the wound to become quite consolidated, and during this time it requires but little—a cough, a sneeze, a motion of the head, the necessary efforts in the use of a urinal or bed-pan, no matter how careful the nursing—to rupture the newly formed union; and, as a matter of fact, this often does take place. The aqueous humor then flows away through the wound with a sudden gush, and, where the simple extraction has been employed, carries with it the iris. In this event, it is that portion of the aqueous humor which is situated behind the iris which is chiefly concerned in the iris-prolapse; the aqueous humor in the anterior part of the anterior chamber probably flows off without influencing the position of the iris.

Many who perform the simple operation endeavor to prevent secondary iris-prolapse by a spastic contraction of the pupil, produced by eserin, which is instilled at the conclusion of the operation, and, again, by some a few hours afterwards. In most instances the desired end is by this means effected. But there is still a considerable percentage of the cases in which the contraction of the sphincter iridis is overcome by the pressure of the aqueous humor from behind, and iris-prolapse takes place.

The formation of even a narrow coloboma prevents prolapse of the iris when the wound bursts, but this is not because the portion of iris which is liable to prolapse has been taken away, for

that would be nothing less than the whole of that part of the iris which corresponds to the length of the opening in the corneal margin. The coloboma averts secondary iris-prolapse, because it provides a way, a sluice, for the aqueous humor contained in the posterior part of the anterior chamber to escape directly through the wound, without carrying with it the iris in its rush; and it is evident that the narrowest coloboma which can be formed will be amply sufficient for the purpose. To my mind a narrow iridectomy here is no "mutilation of the iris," but rather a measure which rests upon a sound basis, and which is calculated to insure the safety of the eye in an important particular.

As to disfigurement of the eye, there is practically none, when the coloboma is so narrow, and is situated in the upper part of the iris. The pupil, too, is movable, almost, if not quite, as much so, as in most cases of simple extraction. For a narrow coloboma does not render the pupil immovable. Where there are no adhesions between the pupillary margin and the capsule, as frequently happens, the reaction to light is active, a drop of atropin will dilate the pupil widely, and a drop of eserin will contract it.

Mental Derangements after Cataract Extractions.—After cataract extractions, during the period of confinement to bed, passing mental disturbances are sometimes seen in old people. This usually takes the form of confusion of ideas, hallucinations, and terror. It is hard to assign a cause for it, but probably it is mainly due to the quiet, and to the exclusion of light following on a period of some anxiety and excitement. A few doses of sulphonal, and permission to sit up—at least in bed—with the admission of more daylight, will be the best measures to adopt in such a case; and speedy restoration of mental equilibrium may be looked for with confidence. Care should be taken not to mistake the symptoms of atropin poisoning for this form of mental disturbance.

Secondary Glaucoma after Cataract Extraction occurs now and then, by whatever method the extraction may have been performed. This is contrary to what one would have expected, in view of the diminished contents of the globe, by reason of absence of the lens, and especially where an iridectomy has been made. High tension in these instances may come on soon after recovery from the cataract operation, or after a good result has existed for many years. Treacher Collins'[8] microscopic investigations show that in these cases either the iris, the capsule, or the hyaloid has

become entangled in the wound and incarcerated in the subsequent cicatrix, and it seems that this leads in some cases to closure of the filtration angle in its entire circumference. Mr. Collins says: In all eyes in which glaucoma comes on after extraction of cataract there is adhesion to or entanglement of the lens capsule in the extraction scar. This adhesion or entanglement keeps the root of the iris, or the anterior of the ciliary process if the iris has been removed up to the periphery, in close contact with the back of the cornea in the region of the coloboma, and so keeps the angle of the chamber blocked in that situation. The advanced position which the capsule takes, by reason of its attachment to the cornea, draws forwards the iris lying in front of it, and in this way approximates the root of the iris, elsewhere than in the region of the coloboma, to the periphery of the cornea. In some cases, especially in those in which the extraction scar is very corneal, the advance in the position of the capsule is so great that the apposition of the cornea and iris is actually occasioned. In such cases the increased tension follows as soon as the wound has closed after the operation. The adhesion of the lens capsule to the extraction scar is sometimes composed of inflammatory exudation; this, on organizing and contracting, tends gradually to advance more and more the position of the capsule, and consequently that of the iris. Such a gradual advance would explain those cases in which the increased tension does not make its appearance until some time after the cataract has been removed.



FIG. 122.

A wide iridectomy, or a sclerotomy, should be made as soon as possible after the high tension shows itself, and by this means many of these eyes may be saved. Simple division of the capsule has produced a good effect in some cases.

Discission or **Dilaceration** means the tearing of the anterior capsule of the lens with a needle, so as to give the aqueous humor access to the lenticular fibers, which causes them to swell, and gradually to become soft, and then to be absorbed. The larger the capsular opening the more freely is the aqueous brought in contact with the lens, and the more rapid is its swelling. The rapidity of the swelling and absorption depend, also, on the consistence of the lens. The softer it is the more rapid is

the process, the completion of which may require from a few weeks to many months. It is wise to make the first discission of moderate dimensions, in order to test the irritability of the eye, especially in adults.

The instruments required are a spring speculum, a fixation-forceps, and a Bowman's stop-needle (Fig. 122). The pupil is to be dilated with atropin.

The eye having been cocainized, the speculum applied, and the eye fixed close to the inner margin of the cornea, the needle is passed perpendicularly through the cornea in its lower and outer quadrant, at a point corresponding to the margin of the dilated pupil. It is then advanced upwards to the upper margin of the pupil (Fig. 123), where it is passed into the capsule, but not deeply into the lens, and a vertical incision is effected by




FIG. 123.

withdrawing the instrument slightly. If an extensive opening in the capsule be wished for, a horizontal incision can be added to the vertical one by a corresponding motion of the needle. During these manuevers the cornea, at the point of puncture, must form the fulcrum for the motions of the instrument. The instrument is then withdrawn from the eye, and some aqueous humor escapes through the opening. Atropin is instilled, and the bandage applied. The patient is kept in bed for a day, and then the bandage may be dispensed with, and dark spectacles worn. The iris is to be kept well under the influence of atropin, until the absorption of the lens is completed. Repetition of the operation is called for, if the opening be so small as to admit of but a very slow absorption of the lens, or if, as sometimes happens, the opening should become closed up.

This method is applicable to all complete cataracts up to the twenty-fifth year of age, and to those lamellar cataracts up to the same age in which the opacity approaches so close to the periphery of the lens that nothing can be gained by an iridectomy. After the above age, the increasing hardness of the nucleus, and the increasing irritability of the iris, render the method unsuitable.

Discission is a safe procedure, when used with the above indications and precautions. The danger chiefly to be feared is iritis, from pressure on the iris of the swelling lens masses. When this occurs, or is threatened, removal of the cataract by a linear incision in the cornea should be at once performed. A safeguard against iritis may be had in a preliminary iridectomy (von Graefe), and it is perhaps well to do this in all cases over fifteen years of age, the discission following some weeks afterwards.

Another danger consists in glaucomatous increase of tension (secondary glaucoma), which may come on without any subjective symptoms, although severe pain usually attends it, while the absorption of the lens runs its proper course. It may happen, consequently, that, when absorption of the cataract is completed, the eye will be found blind from glaucoma. Frequent testings of the tension of the eye during the cure are therefore a most important precaution. Should the tension rise, removal of the lens through a linear incision in the cornea is at once indicated, or the suction operation may be employed.

Suction Operation of Cataract.—This method can only be used for semi-fluid or soft cataracts.

The pupil having been well dilated with atropin, and the eye cocainized, a free opening is made in the capsule of the lens with a discission needle. A linear incision is then made in the cornea about halfway between its center and its margin, and the point of a Bowman's or a Teale's syringe introduced through it, and through the opening in the capsule, into the substance of the lens. Gentle suction is then applied, and the lens substance drawn into the syringe. The syringe should not be passed behind the iris. If it be thought that the cataract is not sufficiently soft, it is desirable to allow some time (a fortnight or so) to elapse between the discission and the suction, in order that the lens substance may undergo disintegration by the action of the aqueous humor. The suction operation is now much less frequently used than formerly.

Secondary Cataract and its Operation: Capsulotomy.—The term secondary cataract, as here used (compare p. 352), usually means a closure of the opening which is present in the anterior capsule after the removal of a cataractous lens, along with a thickening of the capsule in some of the cases, by which an impediment is offered to the rays of light in passing through the pupil. The thickening may have pre-existed in the capsule, or it may be due to subsequent proliferation of the epithelial cells on the inner surface of the capsule. The term is also used with reference to those cases in which no central opening has been made in the capsule (peripheral capsulotomy), and where the latter causes imperfect vision. It




Fig. 124.

is also used in those cases where, after cataract extraction, an exudation in the pupil, consequent upon iritis, has occurred. Finally, and very incorrectly, it is applied to the cases which Fig. 121 represents, in which, after suppuration of the wound with iridocylitis, the iris is dragged upwards, and the pupil is consequently obliterated.

The most simple form of secondary cataract occurs as a very fine cobweb-like membrane—the capsule of the lens—extending over the whole area of the pupil, which can often only be discovered by careful examination with oblique illumination. It may not cause any trouble of vision until some months after the extraction, when some thickening of it may have taken place.

It is a simple matter to make a rent in this delicate membrane with a discission needle.

Where there are thick opacities in the capsule, or inflammatory exudation into the pupil, with, probably, adhesions of the iris to the pupillary membrane, extraction of the latter has been proposed and practiced, but is associated with so much danger, from the unavoidable dragging on the ciliary body and iris, that the proceeding is not often employed.

Sir W. Bowman's Method, with two needles, is here much preferable. The point of a discission needle is passed through the inner quadrant of the cornea, and into the center of the opacity (Fig. 124), and then, with the other hand, a second needle is passed through the outer quadrant of the cornea, and into the membrane, close beside the first needle. The points of the needles are now separated from each other by approximation of their handles, and in this way a hole is made in the membrane.

A very small opening in the capsule, if quite clear, is sufficient to establish good vision.

Noyes' Method.—A Graefe's cataract-knife is entered in the horizontal meridian of the cornea at its temporal margin, and a counterpuncture made in the same meridian at the inner corneal margin. The point of the knife is now withdrawn into the anterior chamber, and made to puncture the secondary cataract, and is then removed from the eye. Two blunt-pointed hooks are then entered into the anterior chamber, one through each corneal puncture, and the point of each passed through the opening in the membrane made with the knife. By traction on the hooks this opening is enlarged, without any dragging on the iris or ciliary body.

Knapp's Method.—Knapp has designed a needle-knife for dividing the capsule. This instrument has a blade 4 1-2 mm. in length. It cuts on one side only, and the blade and the evenly rounded shaft are so proportioned that the shaft fills exactly the opening made by the blade, and consequently the needle can be moved within the anterior chamber in every direction, without escape of aqueous or bruising of the cornea. The instrument must be of the utmost sharpness in point and edge, so that it may cut, and not tear. The point of the needle-knife is entered through a thin part of the capsule, and an opening is cut in it, hard and inelastic bands being avoided.

Iridotomy.—For the cases as in Fig. 121, where the iris forms

a complete and tightly stretched curtain across the pupil, iridectomy is the operation which readily suggests itself. In very few cases, however, does it give a satisfactory result, owing to the inflammatory products which lie behind the iris, and which close up any artificial pupil by their proliferation, which is set going afresh by the dragging of the iris with the forceps. Repeated iridectomies may finally produce a clear pupil, but iridotomy, in which there is no dragging of the iris, is a better operation in these cases.

There are several modes of performing iridotomy, that of de Wecker being the best. A vertical incision having been made in the cornea, about 3 mm. long, and the same distance removed from its inner margin, the closed blades—one of which has a sharp point—of de Wecker's forceps-scissors are passed into the anterior chamber. The blades are then opened, and the sharp point of one of them is forced through the stretched iris, and some 3 or 4 mm. behind it. By now closing the blades the tightened iris fibers are cut across, and on their retraction a central clear pupil is formed in the iris and retro-iridic tissue.

Dislocation of the Crystalline Lens.—This may be congenital, and due to arrested development of the zonula of Zinn; or it may be the result of disease, such, for example, as anterior sclerochorioiditis; or it may be caused by a blow or other trauma.

The dislocation may be partial or complete. In the former case it is often so slight as to be discoverable only when the pupil is widely dilated, the margin of the lens becoming then visible as a curved black line, in some one direction, by aid of the ophthalmoscope mirror; or the displacement may be so great as to bring the margin of the lens across the center of the undilated pupil, in which case one part of the eye will be highly hypermetropic, while in another part it will be myopic. Complete dislocation may take place into the anterior chamber, into the vitreous humor, or even under the conjunctiva, if the sclerotic have been ruptured.

The symptoms in partial dislocation are those of loss of power of accommodation, and monocular double vision. Iridodonesis (*i. e.*, trembling of the iris when the eye moves) is present, as a rule, in consequence of the loss of support provided by the lens. In complete dislocation the symptoms are those of aphakia—*i. e.*, extreme hypermetropia, and want of power of accommodation.

Treatment.—In partial dislocation it is rarely that any treatment can be of service. The prescribing of spectacles suited, so far as it is preferable, to the faulty refraction is indicated. In complete dislocation of the lens into the anterior chamber, its extraction is usually required, especially if it cause symptoms of irritation. Dislocation into the vitreous humor is generally unattended by irritation; but when the latter does arise, removal of the lens by aid of a spoon, through a peripheral corneal incision, has to be attempted.

Lenticonus is a very rare congenital anomaly of the lens, in which its anterior surface, or, still more rarely, its posterior surface, is cone-shaped.

Aphakia(*ἀ,priv.; φακός, a lentil, lens*), or **Absence of the Crystalline Lens.**—The condition of the emmetropic eye after the removal of a cataract is one of high hypermetropia, and the power of accommodation is wanting. Consequently, in order that the eye may have the best possible sight for distant objects, a high convex glass has to be experimentally found to suit it, and stronger lenses must be prescribed for shorter distances.

The degree of vision obtained varies considerably in different cases; frequently V $= \frac{6}{6}$ is obtained,but V $= \frac{6}{18}$ may be regarded as a satisfactory result; and even lower degrees, which enable the patients to find their way about with comfort, are classed as successful operations. The vision often improves for some months after the operation, patients who at first had only $\frac{6}{18}$ or so advancing up to $\frac{6}{9}$ or $\frac{6}{6}$ For writing, reading, etc., at about 25 cm., a still higher convex glass must be provided. If the correcting lens for distant vision be + 10 D, its power, for vision at 25 cm., must be increased by the lens which would represent the amplitude of accommodation from infinite distance up to 25 cm. This lens is 4 D (because $\frac{100}{25} = 4$); therefore + 14 D is the lens required. With these two lenses most patients are satisfied. For distinct vision at middle distances they learn to vary the power of the lenses by moving them a little closer to, or farther from, the eye; but, if necessary, a lens can be prescribed for distinct vision at any desired distance.

In the case of hospital patients one is often obliged to select the + glasses in a fortnight or three weeks after the operation, but the result is more satisfactory when the selection can be postponed for six weeks or two months. Permanent wearing of the + glasses should not be permitted until all redness of the eye

has passed off, and this varies in different cases. Until then, also, dark protection spectacles should be worn.

In the majority of cases, after cataract operations, the best vision is not obtained until a certain degree of astigmatism is corrected. This astigmatism is caused by a flattening of the vertical meridian of the cornea, due to the incision at its upper margin, and hence it is against the rule (see p. 50). An obliquity in the incision often produces an obliquity in the principal meridians of the astigmatism. The degree of astigmatism varies, and may be very high. It rapidly reaches its maximum after the operation, and then gradually diminishes for weeks or months, and in some cases completely disappears; hence it is that glasses for permanent use can be better prescribed a month or two subsequently to the operation.

For an account of Erythropsia after cataract extraction see chap. xvii.

References.

[1] "Trans. Ophthal. Soc., U. K.," 1883, p. 79.
[2] "Archives of Opthalmology," xi. 3. p. 349.
[3] "Ophthalmic Review," 1889, p. 235.
[4] "Gazette Hebdomadaire," 1875, No. 35.
[5] "Archives of Ophthalmology and Otology," vi. p. 545.
[6] "Von Graefe's Archiv," xxxviii. 4. p. 1.
[7] *Ibidem*, li. p. 199.
[8] "Trans. Ophth. Soc., U. K.," x. p. 108; and "Anatomy and Pathology of the Eye," p. 107. London, 1896.

CHAPTER XIV.

DISEASES OF THE VITREOUS HUMOR.

Purulent Inflammation of the Vitreous Humor (to which, unfortunately, the name pseudo-glioma is sometimes applied) occurs only as the result of perforating injuries, or of the lodgment of a foreign body, or as an extension of a purulent process from the chorioid (p. 263).

Ophthalmoscopically, a purulent deposit in the vitreous humor gives a yellowish reflection. It is to be distinguished from a somewhat similar appearance in glioma of the retina by the history, by its early complication with more or less severe iritis, by the very frequent retraction of the periphery of the iris, with bulging forwards of its pupillary part, and by the diminished tension of the eye, while a lobulated appearance is not so usual in it as in glioma. Again, in glioma the vitreous humor remains clear, while in this disease it is hazy.

The condition, if at first confined to the vitreous humor, soon extends to the surrounding tissues, and usually leads to panophthalmitis and complete destruction of the eye.

Inflammatory Affections of the Vitreous Humor, other than the purulent form, are for the most part the consequence of diseases of the chorioid, ciliary body, or retina, and display themselves as opacities of various kinds. These are either cells derived from the primarily diseased tissue, or they are secondary changes (connective tissue development), the result of this cellular invasion.

The chief *Varieties of Vitreous Humor Opacities* are: (1) The Dust-like Opacity so characteristic of syphilitic disease of the retina and chorioid. It may occupy the entire vitreous humor, but is frequently confined to the region of the ciliary body, or to that of the posterior layers of the vitreous humor. (2) Flakes and Threads. These occur with chronic affections of the chorioid or ciliary body, and may be the result also of hemorrhages into the vitreous humor. They invade every portion of the humor. (3) Membranous Opacities, which are rare, and are probably

the result either of extensive hemorrhagic extravasations or of chorioidal exudations.

Hemorrhages into the vitreous humor are not uncommon, and are the result of certain diseases of the retina and chorioid, which are accompanied by hemorrhages in those membranes. They are common in old people, but very large hemorrhages also occur in young adults (see Apoplexy of the Retina). They are also caused by blows on the eye, which rupture the chorioidal or retinal vessels. Most of the alterations occurring in the vitreous humor are attended with, or give rise to, fluidity of it, and may lead to detachment of the retina.

The Diagnosis of opacities in the vitreous humor is made with the ophthalmoscope mirror and a not very bright light, or with the plane mirror. If a very bright light and a concave mirror be employed, the finer opacities will not be readily seen. The pupil being illuminated, the patient is directed to look rapidly in different directions, when the opacities will be seen to float across the area of the pupil, as they are thrown from one side of the eye to the other.

Opacities in the vitreous can be distinguished from those in the lens by the fact that the latter are fixed, and are arranged mostly in a radiating manner. Opacities which lie behind the center of curvature of the cornea, as examined with the ophthalmoscope mirror, seem to move in the opposite direction when the patient moves his eye; while those which lie in front of that point move in the same direction as the eye. Therefore opacities in the lens and anterior part of the vitreous humor, about 0.6 mm. behind the lens,* will move in the same direction.

Another and very fine method for the detection of delicate opacities in the vitreous is by placing a high + lens, say + 10 D, behind the ophthalmoscope mirror, and then going close to the eye, as in the examination of the upright image. Minute opacities will then be seen as black dots floating in the vitreous humor.

When the vitreous humor is full of blood, no red reflex can be obtained with the ophthalmoscope, and the pupil looks quite black. By focal illumination we can observe, in this case, that the lens is perfectly clear, and sometimes the red color of the extravasated blood can be seen behind it.

* Radius of curvature of cornea, 7.829 mm. Distance from anterior surface of cornea to posterior surface of lens, 7.2 mm.

The ophthalmoscope does not always detect changes in the chorioid or retina, when they are opacities in the vitreous; and in many such cases we are led to the belief that the diseased changes in the chorioid or retina are too fine to be seen with the ophthalmoscope, or that they are situated in the region of the ciliary body which is out of view.

Vision is affected by opacities in the vitreous humor in proportion to their density, and to the extent to which the vitreous humor is occupied by them. The patients often observe them as floating positive scotomata in their field of vision. These entopic appearances are caused by the shadows of the opacities thrown on the retina.

The Prognosis depends on the cause of the opacities. Small hemorrhagic extravasations in young people are readily absorbed, but are liable to recur. The dust-like opacities accompanying specific retinitis are also favorable for absorption, while extensive hemorrhages in older people, and the flake and thread opacities, frequently remain as permanent obstructions. Moreover, by shrinking, many of the more organized opacities give rise to detachment of the retina from the chorioid, and consequent blindness.

Treatment consists, above all, in that for the exciting cause. Besides this, Heurteloup's artificial leech, or dry cupping on the temple, is most useful; and in many cases, soon after the application, a marked clearing up of the vitreous is apparent. Pilocarpin hypodermically is worthy of trial. In one case von Graefe operated on membranous opacities by tearing them with a needle, and with a successful result.

Mouches Volantes, Muscæ Volitantes, and **Myodesopsia** (*μυῖα, a fly; ὄψις, seeing*) are terms applied to the motes which people frequently see floating before their eyes, but which do not interfere with the acuteness of vision, nor can the ophthalmoscope detect opacities in the vitreous humor, or any other intra-ocular disease. These motes are most apparent when a bright surface, such as a white wall or the field of a microscope, is looked at. Mouches volantes have no clinical importance. Those annoyed with them should be strongly recommended not to look for them, as in that case others are very apt to become visible. They depend, probably, upon minute remains of the embryonic tissue in the vitreous humor.

Fluidity of the Vitreous Humor, or **Synchysis** (*σύν, to-*

gether; χέω, to pour) is not rare. It can only be diagnosed with certainty when the humor contains floating opacities. Low tension of the eyeball does not always indicate fluidity of the vitreous, although soft eyeballs nearly always contain fluid vitreous humor. Trembling of the iris is also no sign of fluid vitreous, but merely indicates that the iris is not supported in the normal way by the crystalline lens. Defective zonula of Zinn,




Fig. 125.

however, is often caused by, or is a concomitant of, fluid vitreous, and, by causing displacement of the lens, would allow trembling of the iris.

The Causes of synchysis are chorioiditis and staphyloma of the chorioid and sclerotic, and it also occurs as a senile change.

Synchysis Scintillans is a fluid condition of the vitreous humor, with cholesterin and tyrosin crystals held in suspension in it. The ophthalmoscopic appearances are very beautiful, re-

sembling a shower of golden rain. A satisfactory explanation for the occurrence of these crystals in this position has not yet been given. They usually occur in old people, and seldom cause any marked deterioration of vision.

Fluidity of the vitreous humor is not, *per se*, a condition of serious import, unless the eye come to be the subject of an opera-




Fig. 126.

tion involving an incision in the corneo-sclerotic coat, when it renders prolapse of the vitreous more liable to take place.

Foreign Bodies in the Vitreous Humor and Interior of the Eye in General.—One of the most common and most serious accidents to the eye is perforation of the sclerotic, or of the cornea and crystalline lens, by a small foreign body (shot, morsel of iron, copper, stone, or glass), which then lodges somewhere in the interior of the eye, very frequently in the vitreous humor.

In cases where the ophthalmoscope fails us, owing to ex-

travasation of blood, etc., it is sometimes not easy to say whether the foreign body be in the eye, or whether it may merely have punctured the sclerotic without passing through it, and then fallen to the ground. If it be known to have been a small foreign body, which has flown against the eye with force, the probabilities are that it is lodged in the eye.

But, if the case be seen immediately, or soon after the accident,




FIG. 127.

and there be no intra-ocular hemorrhage to obscure our view, the foreign body may frequently be detected with the ophthalmoscope in the vitreous humor as a dark or glittering body, according to its nature, and focal illumination with dilated pupil will often help us to discover a foreign body situated in the anterior part of the vitreous humor. Or, if it cannot be seen, an opaque streak through the vitreous humor, one end of which corresponds

with the sclerotic wound, may indicate the track taken by a foreign body.

In case the foreign body has perforated the cornea, and reached the vitreous humor through the circumlental space, a counter-opening will be found in the iris; while, if it be supposed to have passed through the cornea and lens, the openings both in the anterior and posterior capsule of the lens should be sought for.

Of recent years the *Röntgen Rays* have been very successfully employed in determining not only the presence, but also the exact position of foreign bodies in the eyeball and orbit, and to Mackenzie Davidson is due the credit of having devised a most complete and practical method for the application of the rays, and for the obtaining of skiagraphs. It is as follows:

The patient sits upon a chair in an upright position with his head fixed in a headpiece (clamped to a table) to keep it steady, while at the same time a photographic dry plate can be placed against the temple on the side of the eye which is to be photographed.

Fig 125 shows the patient in position for having his left eye skiagraphed. (If the right eye is to be photographed he sits the other way, so that this headpiece is constructed for taking either the right or left eye.)

Fig. 126 is an enlarged picture of a patient's head in position for taking the right eye. The back of the head rests against a board, and another board, with a thumbscrew sliding in a groove, serves to press and fix his head laterally against two stretched piano-wires, behind which again the photographic plate is placed. The chin is supported on an adjustable projection.

Fig. 127 is a side view of the same patient. The stretched piano-wires are shown. The patient, while the skiagram is being taken, is made to fix his gaze on a distant object, so that his optic axis is parallel to the horizontal wire. Previously, a small piece of lead wire, exactly 1 cm. long, is placed on the lower eyelid, and secured by two strips of adhesive plaster, and the relative position of the point of the wire (nearest the eye) is carefully noted in relation to the cornea (*e. g.*, so many millimeters vertically below the center of the cornea, or so many millimeters vertically below any corneal scar which may happen to be present); also whether the point is on a level with a vertical line from the center of the cornea (as it usually

is), or how far behind or in front of this plane. These are all the adjustments necessary to be made with the patient.

Before the patient is placed in position, the Crookes tube is adjusted, so that the fine point on the anode, from which the linear X-rays originate, shall be exactly opposite the point of intersection of the two stretched piano-wires. When the tube is worked by the coil, this point shows as a bright incandescent spot on the anode, if it be of osmium; and by means of a fixed "sight," placed on this side of the wires, the tube can be so adjusted that this point is exactly opposite the intersection of the wires. The distance is carefully noted; it is usually 28 to 30 cm. The tube-holder is fixed to a bar of wood, which slides horizontally, and by means of marks placed on the bar itself and upon the edge of the groove in which it slides, it can be displaced in a plane exactly parallel to the horizontal wire. It is to be displaced 3 cm. to one side of the vertical or zero point. Then a photographic plate, protected, as usual, in black paper, is placed against the wires (see Fig. 127), and an exposure given of from ninety seconds to two minutes. With exceptionally good osmium tubes ten seconds is enough. The tube is then displaced 3 cm. to the other side of the zero point—the photographic plate having been removed and a fresh one put in its place—and a second similar exposure is given. The result is two negatives taken from two points of view 6 cm. apart.

A transparent sheet of thin celluloid has two cross-lines marked upon it at right angles to each other. One side is varnished, so that it will readily take pencil marks. Immediately after development and fixing, this sheet of celluloid is placed over the film side of the negative, so that its two lines are exactly superimposed upon the white lines left by the wires in the headpiece; while firmly held in position, the shadow of the leaden wire or landmark, placed on the lower eyelid, is carefully traced. Then the foreign body, or foreign bodies, is, or are, traced in the same way. This process of tracing is repeated with the other negative. The result is that upon the sheet of celluloid two tracings of the leaden landmark wire, and two tracings of the foreign body, side by side, are obtained.

This celluloid tracing is now placed upon the horizontal glass stage of the Cross-Thread Localizer. The latter has two fine silk threads coming from two points, which are so adjusted as to occupy relatively the two positions occupied by the anode of the

Crookes tube, and to be at the same distance from the celluloid tracing, and also in the same relative position to the cross-lines that the anode of the Crookes tube had to the photographic plate and to the cross-wires of the headpiece, when the photographs were being actually taken.

The silk threads are now used to trace the linear paths of the X-rays. The intersection of the two threads fixes the position of the object in space.[1] Its geometrical relations to the known data can then be measured. First, the three co-ordinates of the known point are ascertained, then the three co-ordinates of the unknown foreign body, and then, by simple subtraction, the minor co-ordinates are obtained, and thus the position of the foreign body is accurately determined. The surgeon is enabled to say how far horizontally inwards or outwards the foreign body lies from the point of the landmark lead wire; from that point how far vertically upwards or downwards it lies; and, finally, how far directly backwards, parallel to the visual axis, it is situated. If care be taken, the position of a foreign body, however small, can be ascertained with great accuracy by this method. Its size also can be discovered. Moreover, the two negatives are stereoscopic, so that, when viewed either in a Wheatstone's reflecting stereoscope, or by converging the optic axes, and so fusing the pictures, a single picture in relief is seen, showing the relative position of the parts in a very beautiful manner.

The Sideroscope[2] is an instrument devised by Asmus for the detection of the presence of atoms of steel or iron in the eye; but the first suggestion for it was given by T. R. Pooley, of New York, many years earlier. It consists in a magnetic needle hung by a fine thread, and so mounted that, when it is brought close to the eye containing the foreign body, its deflections can be read by means of an astronomical telescope which is attached. The sideroscope is also used for ascertaining the position of the foreign body, which is nearest to the part where the deflection of the needle is greatest. This, of course, is only an approximate localization.

The danger of leaving a foreign body in the eye is great. It is rarely that, whether it remains free or, as sometimes happens, becomes encapsuled, it is tolerated permanently in any part of the interior of the eye, and that event should never be reckoned on in the treatment of such a case.

As a rule foreign bodies in the vitreous, or elsewhere within the eye, soon produce violent inflammatory reaction. This occurs, either by reason of infective micro-organisms being introduced into the eye with the foreign body, or it may be caused by the oxidization of the foreign body, when, as is most common, it is of iron or copper. The form of inflammation may be either a plastic or purulent uveitis, in the latter case with purulent infiltration of the vitreous humor and hypopyon.

An eye which contains a foreign body that is not, or cannot be, at once removed, may be regarded as lost; and such an eye becomes, moreover, one of the surest sources of sympathetic ophthalmitis.

It is, consequently, of the utmost importance to remove every foreign body from the interior of the eye if possible, where there is a reasonable prospect of saving the eye, even partially, by so doing—and this with the least possible delay; or, carefully to watch the eye, and at any sign of inflammatory reaction to excise the eyeball. Indeed, in view of the fact that this inflammatory reaction almost invariably comes on sooner or later, I should be inclined to excise most of these eyes, as soon as it has been decided that the foreign body cannot be extracted.

Removal of the Foreign Body is very often an extremely difficult and disappointing undertaking, but it should always be attempted when, being neither steel nor iron, it is visible within the eye, so that its position can be determined with the ophthalmoscope or by focal illumination, and when it seems that the eye may be saved, at least to some extent.

The introduction of the magnet for the removal of fragments of iron or steel has made it unnecessary that they should in every case be visible, and here the chances of success have been much enhanced since the foreign body can be accurately localized by the Röntgen rays. In all these operations it is necessary that the patient should be deeply under the influence of an anesthetic, in order that as little vitreous humor as possible may be lost. And, again, strict aseptic measures must be observed, lest by our operation the very form of mischief be produced which it is our desire to avert.

There are several methods of proceeding. Atoms of glass, copper, stone, etc., may sometimes be removed through an incision in the sclerotic which is either an enlargement of the opening made by the foreign body, or is a special one, at a point

more nearly corresponding to the actual position of the latter in the eye. This incision should lie between two recti muscles, should have an antero-posterior direction, and, in order that it may gape but little, should be a puncture with a broad keratome. Prolapse of the vitreous is then produced by pressure on the eyeball, and the foreign body is evacuated.

This method should only be tried when the foreign body is situated in the periphery of the vitreous, and towards the equator of the eye, where the opening for its exit can be made in its immediate neighborhood; but the proceeding is often attended with disappointment, much vitreous being lost, while the foreign body remains in the eye.

Or, a forceps is passed in through the opening, and while the foreign body is kept in view with the ophthalmoscope it is seized and drawn out. This plan is also unsatisfactory, as, loss




Fig. 128.

of vitreous occurring, the cornea becomes flaccid, and the view of the foreign body is soon obscured.

Again, some surgeons prefer to make their opening not close to the foreign body, but exactly at the opposite side of the eyeball, by which means they can often reach the foreign body with greater ease, and with less injury to the tissues.

The magnet, thanks to M'Keown, of Belfast,[3] has of late years come into use for the removal of fragments of steel and iron from the interior of the eye, and especially from the vitreous humor. M'Keown used a permanent magnet; but electro-magnets are those now employed for this purpose, the instruments of Hirschberg [4] and of Simeon Snell [5] being the most suitable. Fig. 128 represents Snell's instrument in two-thirds its actual size. It is a core of soft iron, around which is placed a coil of insulated copper wire, the whole inclosed in an ebonite case. To one extremity of the instrument are attached the screws to receive the

connections of a small accumulator. At the other extremity the core projects just beyond the ebonite jacket, and is tapped, and into it screws the point. Points of various kinds or shapes can be adjusted to the magnet, according to the case to be dealt




Fig. 129.

with. A point having been passed through the sclerotic opening, it is advanced towards the foreign body, when the latter adheres to it, and is drawn towards the wound. To prevent too much stirring up of the vitreous, which might lead to its shrinkage, with consequent detachment of the retina, it is desirable,

previously to introduction of the magnet point, to divide the vitreous with a Graefe's knife until close to the foreign body. Much care is required in drawing the foreign body through the opening, lest it be rubbed off the point in its passage. A forceps is generally used at this part of the proceeding, either to dilate the wound, or to seize the foreign body and extract it.

Haab's Giant Electro-Magnet [6] (Fig. 129) is also used for extracting morsels of iron or steel from the eye. It is an immense and very powerful magnet, which is placed close to the eye. Care is required in its use, lest even more injury be done to the delicate tissues of the eyeball by the foreign body in its passage towards the magnet than by its entrance into the eye.

As a rule, Haab recommends that the center of the cornea, in the first instance, be brought opposite and close to the point of the magnet, for by so doing entanglement of the foreign body in the ciliary processes, from which it is not easy again to disengage it, may best be avoided. Foreign bodies which are in the vitreous humor, or which are not too firmly fixed in the retina, slide round the lens and bulge the iris forwards. As soon as this occurs the current is turned off, and the patient's head withdrawn from the magnet. By a suitable turning of the patient's head and eye, the latter being again approached to the magnet and the current turned on, the foreign body is drawn from behind the iris, through the pupil, which has been well dilated with atropin and cocain, and into the anterior chamber. Should it not be possible to get the foreign body away from behind the periphery of the iris with the magnet, an iridodialysis may be formed with a keratome, and the foreign body drawn away with a Snell's magnet or with a forceps. It must not be attempted to draw the foreign body by the magnet through the iris, or the latter may be partially or entirely pulled away. Having got the foreign body into the anterior chamber through the pupil, an incision with a Graefe knife is made in the cornea, if possible without allowing the aqueous humor to flow away, and through the incision the sharp point of the giant magnet is passed, and the foreign body extracted. Even foreign bodies which enter through the sclerotic are best removed through the anterior chamber. When the foreign body is firmly fixed, it may often be loosened by rapid turning on and off of the current, or it may first be drawn towards the equator, and then towards the anterior chamber.

Cysticercus in the Vitreous Humor is not of rare occurrence

in some parts of Germany, but there have not been many such cases observed in the British Isles. Hill Griffith,[7] however, has published so many as seven cases from Manchester.

The original seat of the entozoön is usually beneath the retina (see chap. xv.), through which it breaks to reach the vitreous humor; but it also sometimes makes its first appearance in the vitreous. It is recognized by its peculiar, somewhat dumb-bell shape, its iridescence, and its peristaltic motions. The vitreous humor often becomes full of peculiar membranous opacities, as a consequence of the presence of the cysticercus.

Treatment.—Removal by operation. The prospects for the eye are very much worse than in the case of a subretinal cysticercus.

Blood-Vessels are sometimes formed in the vitreous humor. They spring from the retinal vessels, often in connection with connective tissue formations which accompany hemorrhages; but sometimes small loops arise in the neighborhood of the disc, without any hemorrhagic disease.

Persistent Hyaloid Artery.—In intra-uterine life the hyaloid artery is a prolongation of the central artery of the retina, and runs from the papilla to the posterior surface of the crystalline lens. It completely disappears prior to birth, except in those rare cases where it remains as an opaque string, which may stretch the whole way from papilla to lens, or may extend only part of the way. It is then thrown into wave-like movements by the movements of the eyeball, and is easily recognized with the ophthalmoscope. It does not usually cause any disturbance of vision.

Detachment of the Vitreous Humor from the Retina, although probably a common diseased condition, cannot as yet be recognized with certainty during life, and rarely becomes the immediate cause of blindness. Its danger lies in its liability to bring about detachment of the retina.

Detachment of the vitreous may be either idiopathic or due to trauma. In the idiopathic cases chronic chorioiditis is the primary disease, which gives rise to a change in the fine connective tissue elements of the vitreous, with consequent shrinking of this body. Yet, with the ophthalmoscope, the chorioid may seem normal; and, moreover, although floating opacities may be present in the vitreous chamber, yet it is quite possible for a perfectly clear vitreous to be detached.

The condition occurs in connection with high degrees of myopia, where chorioiditis is also common, and is probably the most important factor in the production of the detachment of the retina, so frequent in these eyes. Anterior staphyloma, hemorrhages into the vitreous humor, and neoplastic growths between the vitreous and retina also gives rise to detachment of the vitreous.

Detachment of the anterior portion of the vitreous occurs in many cases of iridocyclitis.

With regard to traumatic cases, all perforating injuries attended with loss of vitreous, including cataract operations—and sometimes, when the wound is in the sclerotic, without loss of vitreous—are liable to be followed by detachment of the vitreous.

I observed a case [8] in which detachment of the vitreous was the chief lesion in the eye, and was the cause of blindness, the



Fig. 130.

vision being reduced to perception of light. The detachment had probably been brought about by an idiopathic hemorrhage from the ciliary body into the anterior part of the vitreous. It lay (Fig. 130) immediately behind the lens and in contact with it, and presented the appearance of a grayish opacity, very like a detached retina but for the absence of retinal vessels. Suspicion of an intra-ocular tumor existing, the eye was removed. The vitreous lay against the ciliary body and lens, while the vitreous chamber was filled with serous fluid, and the retina was in its normal position. In the retina, towards the ora serrata, there were a few minute hemorrhages.

References.

[1] "British Medical Journal," January 1, 1898.
[2] "Das Sideroskop und seine Anwendung." Wiesbaden, 1898.
[3] "Brit. Med. Journal," 1874, vol. i. p. 800.

[4] "Centralblatt für prak. Augenheilkunde," 1879, p. 380.
[5] "The Electro-magnet," etc. London, 1883.
[6] "Bericht. d. Ophthalmologischen Gesellschaft." Heidelberg, 1892. *Ibidem,* 1902. "Klin. Monatsblätter f. Augenheilkunde," 1902, p. 193.
[7] "Trans Ophthal. Soc., U. K.," xxvii. p. 220.
[8] *Ibidem,* ii. p. 41.

CHAPTER XV.

DISEASES OF THE RETINA.

Diseases of the retina may, for the purpose of description, be conveniently grouped as follows: Alterations in Vascularity, Inflammation, Atrophy, Diseases of the Blood-Vessels, Injury by Strong Light, Tumors, Parasitic Disease, Detachment, and Traumatic Affections.

Alterations in the Vascularity of the Retina.

Hyperemia and **Anemia** of the retina, due to changes in the capillary vessels, cannot be seen with the ophthalmoscope, hence these terms are used to denote apparent enlargement or diminution of the principal branches of the central vessels. *Venous Engorgement* may occur as a local condition, as in papillitis, retinitis, thrombosis of the central vein, or as part of general venous obstruction in cardiac and pulmonary diseases. *Contraction of the Arteries* may also be due to local disease of the vessels (embolism, albuminuric retinitis, etc.) and spasm (malaria, quinin), or, more rarely, to diminished blood supply from general causes (cholera). The opposite conditions, namely, diminution in the size of the veins, and enlargement of the arteries, are rarely noticeable.

Inflammations of the Retina: Retinitis.

Retinitis, in general, is characterized by the following ophthalmoscopic appearances: *diffuse cloudiness,* especially of the central portion of the fundus, due to loss of transparency in the retina, and consequent veiling of the chorioid; the *optic papilla* becomes more or less congested, with indistinctness of its outline which in the erect image resolves itself into a delicate striation; *vascular engorgement,* the retinal veins especially becoming enlarged and tortuous. The inflammation in some cases may subside at this stage, but as a rule *hemorrhages* and whitish *exudations* soon make their appearance.

The various forms of retinitis are distinguished by the predominance of some of the above signs, and also by the peculiar appearance and grouping of the exudations.

If the optic papilla be not merely congested, but also swollen, the condition is called Neuro-Retinitis.

In some cases of retinitis the chorioid is also involved, and to these the name chorio-retinitis is given.

Inflammation of the retina is rarely a local affection, being in most cases due to general diseases, and hence it commonly occurs in both eyes.

Syphilitic Retinitis (or Syphilitic Chorioido-Retinitis).—Inherited or acquired constitutional syphilis is liable to induce a form of chronic diffuse retinitis. In the acquired disease it is a later secondary symptom, coming on between the sixth and eighteenth month, often only in one eye.

With the Ophthalmoscope a slight opacity of the retina is seen extending from the papilla some distance into the retina, and very gradually disappearing towards the equator of the eye. The papilla is but slightly hyperemic, while its margins are indistinct, like those of the moon seen through a light cloud. The artery is not generally altered, and the vein but slightly distended. Opacities in the vitreous humor are not uncommon. They may be membranous or thread-like, but a diffuse dust-like opacity, filling the whole vitreous humor, is almost pathognomonic of a syphilitic taint, and may create much difficulty in the ophthalmoscopic diagnosis of the retinal affection.

Disseminated chorioidal changes, in the form of small yellowish spots with pigmentary deposits, are very frequent, especially towards the equator of the eye. Many observers, indeed, hold that the whole process is primarily in the chorioid, and that the retina is only secondarily affected. Fine whitish dots and pigmentary changes often occur about the macula lutea.

Vision may be but slightly affected, but in the advanced stages it is usually much lowered. Central, or peripheral, or ring scotomata, or concentric defects of the field, are found. The scotomata are often positive—*i. e.*, they can be seen by the patient as dark spots in the field. Night-blindness is a constant symptom, and the light-sense is enormously diminished. The patients sometimes complain of sparks or lights, which seem to dance before their eyes, and occasionally also of a diminution in the size (micropsia) of objects, or of a distortion (metamorphopsia) of

their outlines. The micropsia is believed to be due to a separation from each other of the elements of the layer of rods and cones by subretinal exudation. The image of an object then comes into relation with fewer of these elements, and hence the mental impression is that of a smaller object than is conveyed by the image formed in the sound eye, or on a sound part of the same retina.

The Progress of the Disease is very slow, and is liable to relapses. In the late stages extensive pigmentary degeneration of the retina may come on, or disseminated chorioiditis. But if the cases come under suitable treatment in an early stage, a cure may often be effected.

Treatment.—The only remedy which is of real value is mercury, and that in an early stage. It should be used in a protracted course of some weeks by inunction, combined at discretion with small doses of calomel internally. Perchlorid of mercury hypodermically, in 1-20 grain doses once a day, is also a suitable measure. If mercurialization be effected, it should not go further than a very slight stomatitis. Turkish baths, and the artificial leech at the temple, may be employed as adjuncts to the treatment. When the mercurial course has been completed, iodid of potassium should be prescribed as an after-treatment. Complete rest of the eyes, and protection from strong light by dark glasses, are also necessary in this, as in many forms of retinitis.

Hemorrhagic Retinitis.—In this affection the retina contains a number of small hemorrhages. They occur chiefly between the fibers of the inner layer, and consequently present a flame-like appearance as seen with the *Ophthalmoscope.* Any which lie in the outer layers are more apt to be round or irregular in shape. In addition to the hemorrhages there is diffuse opacity of the retina, and sometimes white spots of degeneration. The papilla is often much swollen, and the retinal veins distended and tortuous, while the arteries are small; but these appearances, as well as the number of the hemorrhages, vary much in different cases. When there are but few hemorrhages, they are situated in the neighborhood of the papilla and macula lutea. The appearances occasionally resemble those of albuminuric retinitis, but in the latter, as a rule, the proportion of white spots to hemorrhages is greater than in this affection. Probably many cases described as hemorrhagic retinitis are due to thrombosis of the

central vein. (See p. 412.) In a small proportion of cases glaucoma (p. 340) comes on.

Causes.—The affection is found most commonly in connection with cardiac disease—*e. g.,* valvular insufficiency, and hypertrophy of the left ventricle; or with diseases of the vascular system—*e. g.,* atheroma, and aneurysms of the large vessels. Where it is due to disease of the coats of the arteries, the ophthalmoscope will occasionally reveal an arterial branch altered to the appearance of a white thread; but usually the degenerative change does not interfere with the transparency of the vascular coats. In the majority of cases dependent on cardiac or vascular disease the retinal affection is monocular. This, and the frequently sudden onset of the retinitis, suggests some second factor for its occurrence, probably multiple embolisms of the small branches of the central artery. Suppression of menstruation, or other wonted discharge—such as that from piles—has been observed as an immediate cause of hemorrhagic retinitis.

A peculiar form of hemorrhagic retinitis is sometimes associated with secondary syphilis. In addition to the usual opacity of the retina in syphilitic retinitis, a portion of the retina is covered with numbers of small round hemorrhages lying in the different layers of the retina, while a connective tissue development is occasionally found in the nerve-fiber layer, in the form of white striæ along the course of the blood-vessels.

The disturbance of vision is considerable, especially if the neighborhood of the macula lutea be much involved.

The Prognosis is unfavorable in severe cases of hemorrhagic retinitis. Relapses are common, while the ultimate tendency is towards atrophy of the retina and papilla. In very mild cases recovery may come about.

The Treatment must be chiefly expectant, or directed, at most, towards procuring rest for the general system, or for the organ primarily at fault. Dry cupping on the temple, hot footbaths, and iodid of potassium internally may be employed.

Retinitis Albuminurica occurs as a complication in many cases both of acute and chronic nephritis, and in the albuminuria of pregnancy. It is most common with the small granular kidney, but may attend any chronic form of Bright's disease, and occurs in 6 or 7 per cent. of these cases.

The Defect of Vision in the chronic form, although often an early or even the first symptom, is never associated with an early

stage of the kidney disease, but rather with a late stage of it, and with dilated left ventricle. Both eyes as a rule are affected, although often not equally so. Vision is much lowered, and even perception of light may be wanting; but the blindness is not always all due to organic changes in the retina, being often largely the result of uremia.

Ophthalmoscopic Appearances.—These are venous hyperemia and swelling of the papilla, and of the retina in its neighborhood; hemorrhages on the papilla, and in the nerve-fiber layer of the retina; and round or irregularly shaped white spots in the retina, arranged in a zone around the papilla, some three papilla-diameters removed from it. These changes take place in the order in which I have enumerated them. The hyperemia and engorgement of the veins, often very great, become less according as the white spots become more developed. Near the macula lutea no very coarse changes usually occur; but fine white dots are found, with a star-like arrangement converging towards the macula. In some cases the spots spread out only on the inner side of the macula towards the papilla. The degree in which all these different changes are present varies in different cases, no one of them being pathognomonic of the kidney affection, but rather the grouping of the whole picture being suggestive. Sometimes the papillitis is so intense as to simulate that formerly known as congestion papilla in cases of intracranial tumor; while the white spots are sometimes developed to such a degree as to become confluent, and to form one large white plaque. Again, the papillitis, or white spots, or both, may be but slightly marked. The number and size of the hemorrhages are also liable to great variation. Detachment of the retina has been observed in a few cases, and in some the hemorrhages burst into the vitreous humor.

Some of the white spots are caused by fatty degeneration of the outer layers of the retina (the retinal vessels passing over them), others by hypertrophy of the nerve-fiber layer (the retinal vessels hidden by them). The fine dots about the macula lutea are the result of fatty degeneration of the inner ends of Müller's fibers. Small aneurysmal dilatations of the arteries occur very occasionally.

The connection between the renal and retinal affections is not known with certainty, but the theory that the latter is due to chronic uremia is probably correct.

Prognosis.—In these chronic cases the prognosis as regards the patient's life is bad. The majority die within eighteen months or two years; but if the general disease remains stationary, or improves, or recovers, the retinal changes may improve or disappear, and may leave the retina with normal appearances and functions; or the swelling, hyperemia, white spots, and hemorrhages may give place to optic atrophy, with diminution in size of the arteries, pigmentary alterations in the retina, and blindness. In the albuminuria of pregnancy, and in acute nephritis, the retinal complication may disappear with the renal disorder, leaving good vision.

Treatment.—Dry cupping at the temple is about the only remedy which can be employed directly for the retinal affection, and I cannot say that it is of much use. Taking into consideration the serious import of this eye-symptom for the life of the patient, it is a question whether, in many cases of pregnancy with albuminuric retinitis, abortion should not be resorted to, especially if the pregnancy have still some months to run. But on the whole the prognosis is more favorable in the albuminuria of pregnancy than in interstitial nephritis.

Retinal Affections in Diabetes.—There is no one condition of the retina characteristic of diabetes, although undoubtedly retinal affections occasionally do complicate it in an advanced stage. Small retinal hemorrhages, with fine changes in the form of glistening dots, about the macula lutea, somewhat similar in appearance to those which occur in Bright's disease, except that they rarely form the well-marked star, are perhaps the most common and suggestive appearances. In other cases retinal hemorrhages alone are found, and in others hemorrhagic retinitis; while, again, the so-called typical appearances of Bright's disease may be presented. There are often opacities of hemorrhagic origin in the vitreous humor, and iritis may come on.

Leber lays down the important rule that in all cases of retinal hemorrhages and of retinitis hemorrhagica the urine should be examined for sugar.

Retinitis Leukemica.—In not more than one-third or one-fourth of the cases of leukocythemia does a retinal affection occur, and it is not always of the same type. It may consist in a light diffuse retinitis, accompanied by some extravasations of pale blood; while the blood-vessels are also pale, the veins being much enlarged, and rather flattened than overdistended, the

arteries small, and the chorioid of an orange-yellow color. Or, it may resemble a case of ordinary hemorrhagic retinitis.

The Appearances most characteristic of the affection are: a pale papilla with indistinct margins; slight opacity of the retina, especially along the vessels; small hemorrhages; round white, elevated spots up to 2 mm. in diameter, with a hemorrhagic halo, situated by preference towards the periphery of the fundus and at the macula lutea, but not at all, or only in very severe cases, in the zone between the macula and the equator of the eye. These white spots consist of extravasations of leukemic blood, the results, probably, of diapedesis.

Vision may be but little affected if the macula lutea be fairly free. Hemorrhage into the vitreous humor may cause complete blindness.

Retinitis Punctata Albescens.—This disease commences in early childhood, or is perhaps congenital. It often occurs in more than one member of a family, and the parents are frequently blood-relations. The main symptom is night-blindness; in good daylight central vision is usually not defective to any marked degree. The field of vision is contracted. Ophthalmoscopically the fundus is sprinkled over with innumerable small white dots, which, for the most part, are free from any pigmentary disturbance in their neighborhood. In some cases, towards the periphery of the fundus, signs of chorioidal atrophy are present. It is thought by some that this disease is related to retinitis pigmentosa.

Treatment is of no avail.

Development of Connective Tissue in the Retina, or **Retinitis Proliferans.**—Extensive white striæ, formed of connective tissue, are sometimes seen in the retina, and may even conceal the vessels and papilla. They are the result of hemorrhages, traumatic or otherwise, according to Leber, and of an inflammatory process according to Manz, and are formed by proliferation of Müller's fibers and new growth of connective tissue. Hemorrhages in the retina, or in the vitreous humor, or in both, are generally present at some period. Vision is often but slightly affected, but the danger of recurrent intra-ocular hemorrhages renders the ultimate prognosis unfavorable as a rule.

Treatment—Heurteloup's leech. Iodid of potassium, or perchlorid of mercury. Protection spectacles.

Retinitis Circinata is a rare disease, first described by Fuchs.[1]

It occurs in old people, chiefly women, and is characterized by very remarkable appearances. At the macula is a gray or yellowish cloudy patch, which may attain the size of the papilla, and sometimes presents hemorrhages on its surface; surrounding this, but separated from it by a healthy zone, is a ring composed of numerous closely set, small white spots, which are confluent in places. The sight gradually becomes much deteriorated. A large central scotoma develops, and vision is finally reduced to finger-counting centrally, although for a long time the peripheral field may not become contracted. Total blindness rarely results.

Purulent Retinitis is observed as the result of septic embolism of the retinal arteries in septicemia after surgical operations, etc., and very frequently in cases of metria, and it is usually, in the latter condition, a fatal sign.

In an early stage the *Ophthalmoscope* shows a number of small hemorrhages in the retina, with general cloudiness of the retinal tissues, while the actual embolisms, which are usually multiple, may not be visible.

The inflammation makes rapid progress, soon destroying sight, and extending to the chorioid, iris, and vitreous humor, until finally panophthalmitis is reached. The retina is sometimes alone the primary seat of the embolic attack, and sometimes the chorioid is also involved. The embolisms are often little more than masses of micrococci.

The retina becomes secondarily implicated in many purulent processes, which commence in other parts of the eye.

Atrophy of the Retina.

Atrophy, or degeneration, of the retina is characterized by diminution, or even complete obliteration, of the retinal vessels, accompanied by more or less atrophy of the optic papilla. It may be caused by severe forms of retinitis, and also by embolism or thrombosis.

Retinitis Pigmentosa is a degenerative rather than an inflammatory affection of the retina. It is extremely chronic in its progress, coming on most commonly in childhood, and often resulting in complete, or almost complete, blindness in advanced life.

Vision is much affected, but the symptom most complained

of is night-blindness, due rather to defective power of retinal adaptation than to defective light-sense. The field of vision, moreover, becomes gradually contracted, until only a very small central portion remains; so that, although the patient may still be able to read, he cannot find his way alone—a function for which the eccentric parts of the field are the most important. A ring scotoma in the field of vision is present in some cases. Finally, the last remaining central region becomes blind.

The Ophthalmoscopic Appearances consist in a pigmentation of the nerve-fiber layer of the retina, which commences in the periphery, but not at its extreme limits, and in the course of years advances towards the macula lutea. The pigment is arranged in stellate spots, of which the processes intercommunicate, so that the appearance reminds one of a drawing of the Haversian system of bone. Pigment is also deposited along the course of many of the vessels, hiding them from view. The degree of pigmentation varies much, and in some cases is quite absent, and the diagnosis then has to depend upon the other appearances and on the symptoms. The papilla is of a grayish-yellow color, never white, and the vessels are very small.

The chorioid is sometimes slightly affected, irregularity in its pigmentation being observable. At the posterior pole of the crystalline lens there is often a star-shaped opacity.

Pathology.—The pigment in the retina is believed to wander into it from the pigment-epithelium layer. The other pathological changes in the retina consist in hyperplasy of its connective tissue elements, and thickening of the walls of the vessels at the expense of their lumen.

The chorioidal vessels, too, are altered, owing to endarteritis, which causes hypertrophy of their coats, with more or less obliteration of their lumen. In fact, it seems probable that the primary seat of the diseased process is in the chorioid; and that it is the changes in it which cause the pigment from the pigment-epithelium layer to wander into the retina.

Causes.—Retinitis pigmentosa often affects more than one member of a family; and the patients, too, are frequently defective in intelligence or deaf and dumb. Many of them are the offspring of marriages of consanguinity, and in others an inherited syphilitic taint is present, while in others no cause can be assigned. Other congenital defects, supernumerary digits, etc., are sometimes present.

Treatment is of little use. At best one may stimulate the torpid retina temporarily by hypodermic injections of strychnia or with the continuous current.

Gyrate Atrophy of the Retina and Chorioid (Fuchs).[2]—The disease, according to Fuchs, who has seen only a few cases, is apt to occur in more than one member of the same family, and in children whose parents are blood-relations. The first symptom appears in childhood as night-blindness. The optic papilla is atrophied, as in retinitis pigmentosa, and atrophy of the retina is shown by the narrowing of its vessels. The characteristic feature is the peculiar form of chorioidal atrophy. In a zone with the papilla for its center, and extending nearly to the latter, white atrophic dots with sharp margins form, and gradually increase in size, until they become confluent. The atrophy involves both the pigment epithelium and the stroma of the chorioid. The papilla is finally surrounded by a broad white girdle, from which it is separated by a band of normally colored fundus. The edge of the girdle towards the papilla is scalloped, because the separate rounded parts of which it is composed extend backwards in varying distances, while the remains of the normal fundus project forwards between them in sharp processes. There is often, as in retinitis pigmentosa, a star-shaped posterior polar cataract. In addition to the night-blindness, central vision is much contracted. Fuchs believes the disease is closely related to retinitis pigmentosa, being differentiated from the latter by the prominence of the chorioidal atrophy.

Diseases of the Retinal Vessels.

Apoplexy of the Retina.—This differs from hemorrhagic retinitis in that the hemorrhages are found in a retina free from other diseased appearances, retinitis in particular.

With the Ophthalmoscope the extravasations of blood appear as red, or almost black, spots of various sizes and shapes. Their number and position in the fundus are also variable. They may be in any layer of the retina, and sometimes burst into the vitreous humor, and sometimes become extravasated between the retina and chorioid.

Vision is interfered with according to the position and extent of the hemorrhages. Wherever an apoplexy be situated, the function of the retina at that place is suspended. If it be at

the macula lutea, central vision will be seriously impaired; while the scotoma produced by an apoplexy at the periphery of the fundus may pass unnoticed by the patient.

Causes.—Retinal apoplexies are most common in advanced life, with atheroma of the blood-vessels, and are then valuable as a warning of possibly impending cerebral mischief. Other causes are: Hypertrophy of the left ventricle; suppression or irregularity of menstruation or at the climacteric period; the sudden reduction of tension of the eyeball after iridectomy for glaucoma; the gouty diathesis (Hutchinson); progressive pernicious anemia, or anemia from loss of blood (hematemesis, etc.), or from exhausting diseases.

In young people of both sexes, from the fourteenth to the twentieth year of age, large retinal apoplexies, which may extravasate into the vitreous humor, are sometimes seen, and it is difficult to assign a cause for them. Some of the subjects are weak or anemic, while many of them are in perfect health.

Prognosis.—The apoplexies are observed, in the course of weeks or months, to become paler and smaller, often leaving after them chorioidal changes, or grayish spots dependent on degeneration of the retina, and in some extreme cases atrophy of the whole retina may result.

Occasionally absorption of the hemorrhages is accompanied by complete restoration of vision, but usually the scotomata remain. Recurrences of the hemorrhages are very common. Glaucoma comes on as consecutive to retinal apoplexies in some instances, and is then known as hemorrhagic glaucoma, an incurable form of the disease (p. 340).

In other cases the hemorrhage, having invaded the vitreous humor, gives rise to dense permanent opacity in it, followed, perhaps, by detachment of the retina.

Treatment.—Active measures are of little use. Cold compresses at first, with a pressure bandage and dry cupping to the temple, may be employed. The general state of the patient must be attended to, and no violent muscular efforts permitted.

Embolism of the Central Artery of the Retina.—Sudden or very rapid blindness, beginning at the periphery of the field, and advancing towards the center, is the only symptom experienced by the patient.

Immediately after the attack the *Ophthalmoscope* shows a marked pallor of the papilla, while the artery and its branches

are empty of blood, resembling fine white threads, and the veins are diminished in size at the papilla, but increase somewhat towards the periphery. Pressure on the eyeball produces neither pulsation nor change in caliber of the vessels, as it does in a sound eye. Usually, on the following day, the central region of the retina begins to assume a grayish-white opaque appearance, consequent on disturbance of nutrition, in the midst of which the macula lutea is seen as a purple-red spot. De Schweinitz has seen this cherry-red spot at the macula twenty minutes after the embolism took place. The little blood contained in the vessels may soon be observed to divide into short columns with colorless interspaces, and these short columns move along the vessels with a slow jerky motion. Minute hemorrhages often occur, most commonly between the macula and the papilla.

The peculiar appearance of the macula lutea is certainly not due to hemorrhage. According to Liebreich it is merely a contrast effect, the red color of the chorioid shining through, where no nerve-fiber layer is present. Leber suggests that the color may be due to the retinal purple.

The infiltration of the retina passes away in a few weeks, and also the peculiar appearance of the macula lutea, while atrophy of the retina and papilla usually supervenes.

Embolism of a branch only of the central artery has been observed. In these cases the infiltration and the defect of vision are confined to the part of the retina supplied by the embolized branch.

Prognosis.—Vision may improve for a time, but when atrophy commences it falls back again, and finally power of perception of light is lost. Cases of embolism of a branch are more likely to recover.

Causes.—Endocarditis; mitral disease; atheroma of the large arteries of the body; aneurysm of the aorta; pregnancy; Bright's disease. Two cases of chorea with embolism of the central artery are recorded.[3] But it is said also to occur in healthy persons without any discoverable cause.

Treatment.—Repeated paracentesis of the anterior chamber has been tried, and also iridectomy, with the object of reducing the tension, and in this way promoting a collateral flow of blood, by means of the only ascertained (Leber) communications between the retinal and chorioidal vascular systems—namely, at

the entrance of the optic nerve. These attempts have been unsuccessful.

Several cases have been published in which the circulation, which probably was not completely impeded by the embolus, was restored and good vision regained, the recovery being probably due to the manipulations of the eyeball made in each case for the purpose of observing the effect of pressure on the vessels. So long as the pressure was maintained, a column of blood was being stored up behind the embolus, and, on removal of the pressure, it rushed forward against the impediment, carrying the latter into some more remote vessel or into the general vascular system. In fresh cases massage of the eyeball suitably applied would, therefore, always be worth the trial.

Thrombosis of the Retinal Artery.—Blocking of the artery may occur spontaneously, from thrombosis due to failure of the heart's action and slowing of the arterial flow, the result, in its turn, of cardiac disease, spasm of the blood-vessels, disease of the walls of the vessels, or alterations in the quantity and amount of blood.

The *Ophthalmoscopic Signs* are in all respects similar to those of embolism.

The *Diagnosis* between thrombosis and embolism of the central artery can only be made by certain symptoms, which precede or accompany the attack in thrombosis, but are wanting in embolism. These are: previous attacks of transient blindness in the blind eye, a simultaneous attack of blindness in the fellow eye, and faintness, giddiness, and headache at the onset of the blindness.

Treatment.—When transient attacks of blindness are complained of it is important to overhaul the patient's general state, and to correct, so far as possible, any condition which might be the cause of feeble circulation. When the true attack comes on, manipulation of the eyeball applied immediately, or paracentesis of the anterior chamber, might prove of use.

Thrombosis of the Retinal Vein is seen chiefly in old people with atheromatous arteries or cardiac troubles. Orbital cellulitis, from erysipelas or other causes, may also produce it.

The Ophthalmoscopic Appearances are: extreme engorgement of the retinal veins, with great narrowing of the arteries; the whole fundus is thickly studded with dark hemorrhages; the optic papilla after a time becomes pale, and undergoes atrophy,

and the hemorrhages, having become absorbed, leave an atrophied retina with thready arteries.

The Prognosis is very bad, sight becoming permanently damaged or lost, and *Treatment* can only be directed to the general condition.

Aneurysm of the Central Artery of the Retina occurs either as a relatively large dilatation on a main branch of the artery (a very rare condition), or as small miliary aneurysms, which may indicate the presence of others in the small arteries of the brain. Two interesting cases of the latter kind have been recorded, one by Story and Benson,[4] and the other by Perinow [5] in men aged respectively twenty and forty. In one of these cases there were also extensive connective tissue changes in the retina, the veins were dilated in places, and only one eye was affected. The minute aneurysmal dilatations were either globular, and situated laterally on the vessels; or they were fusiform, and involved the whole of its lumen. The number of aneurysms in an eye varied from three to nine. Neither case was followed to its end; but it is to be presumed that such eyes would run great risk of being ultimately lost through intra-ocular hemorrhage.

A rational *Treatment* for the condition can hardly be devised.

Sclerosis of the Retinal Vessels (Perivasculitis, or, more rarely, Endarteritis) reveals its presence by narrowing of the blood column, and by the appearance of white lines along the vessels. It usually begins in the large trunks on the papilla, and may not extend much beyond the latter, as in some cases of optic atrophy; while in other cases (Bright's disease, hereditary syphilis) it involves the small branches as well, and promotes thrombosis and retinal apoplexies, and may even ultimately lead to obliteration of the lumen of the vessels, so that they look like white branching streaks. The arteries are more liable to this condition than the veins.

Quinin Amaurosis.—Quinin in large doses, and very occasionally in small doses, is liable in some individuals to cause amblyopia, which may come on almost suddenly, and which may amount to absolute blindness, accompanied for some hours or days by great deafness. This absolute blindness is rarely more than temporary, although it may last for some weeks—indeed, there is but one case of permanent quinin amaurosis on record; but in severe cases concentric contraction of the field is apt

to remain permanently, with or without some defect of central vision. In the only instance of this more serious result which I have seen, the color and light senses, notwithstanding the contracted field and marked seeming optic atrophy, were normal; but the adaptation of the retina, as shown by considerable night-blindness, was defective.

Major Yarr [6] finds that doses of sulphate of quinin of more than 20 grains are dangerous to the sight, and that more than 40 grains should not be given in twenty-four hours. During the early stages the pupils are widely dilated, and the cornea and conjunctiva are sometimes anesthetic.

In what may be called the acute stage, the *Ophthalmoscopic Appearances* are sometimes normal, but pallor of the optic papilla, with scarcity and smallness of the retinal vessels, is the usual condition. Where the case is chronic—the fields remaining contracted, although central vision has improved—the ophthalmoscope may discover a very pale optic papilla with minimal vessels.

The retinal ischemia is doubtless the immediate *Cause* of the amblyopia, and is the result of diminished heart's action and lowered arterial tension, both of which have been shown to be produced by large doses of quinin. Destruction of the ganglion cells of the retina towards its periphery has been found, and to it may be referred the permanent contraction of the field of vision in some cases.

Treatment.—Cessation of the quinin. Digitalis internally to raise the arterial tension, nitro-glycerin, hypodermic injections of strychnia, and general tonic treatment. Yarr has found that nitrate of amyl causes only temporary improvement of vision.

Injury of the Retina by Strong Light.

Blinding of the Retina by Direct Sunlight.—This is especially likely to occur on the occasion of solar eclipses, from observation with unprotected eye.

Immediately after the exposure the patients complain of a dark or semi-blind spot in the center of the field of vision—a positive scotoma, in short, which may even be absolute, and which interferes with vision in proportion to the length of the exposure. There may also be a central defect for colors, which may extend over a larger area. A peculiar oscillation, or rotary movement, is frequently observed by the patient in the

scotoma, and is very persistent. Objects may also seem twisted or otherwise distorted (metamorphosia).

The Ophthalmoscopic Appearances may be normal, but as a rule some changes exist, such as an alteration or loss of the light reflex at the macula, or a minute pale orange spot near the fovea, with, especially in the later stages, some darkening or pigmentation. When the cases are not severe, improvement in vision takes place, but complete recovery is not common. Hitherto no case in which the vision had been reduced to less than $\frac{1}{3}$ has regained $\frac{6}{6}$.

Treatment.—Hypodermic injections of strychnia, the constant galvanic current, and dry cupping on the temple afford the best chances for promoting the cure. Rest and dark protection glasses are very important.

Snow-Blindness.—Exposure of the unprotected eyes for a length of time to the glare from an extensive surface of snow produces, in some persons, a peculiar form of ophthalmia, which may be followed by temporary or even permanent amblyopia. Although this condition is chiefly an affection of the conjunctiva, it is described here in order to compare it with the effects of sunlight and electric light.

Snow-blindness begins with sensations of a foreign body in the eye, photophobia, blepharospasm, and lacrimation; later on chemosis, with small opacities, or ulcers, of the cornea, comes on. The condition passes off in three or four days without leaving any permanent ill results, except in rare cases, when there may be some secondary hyperemia of the retina.

Treatment.—The preventative treatment consists in the wearing of dark smoked glasses when traveling on the snow; while for the ophthalmia cold applications and cocain are recommended, to relieve the distressing symptoms.

Effects of Electric Light on the Eyes.—The degree of intensity of light required to produce injurious effects on the eye is not known; but this much is certain, that no bad results have been observed from the ordinary use of the incandescent light. Two groups of symptoms are observed from the action of a strong electric light on the eyes.

(*a*) *Electric Ophthalmia.*—This has been chiefly seen in those employed in electric welding operations, and less frequently in electricians who use strong arc-light. The symptoms begin shortly after exposure to the light, always within twenty-four

hours, and are the same as those present in snow-blindness; the lids also are swollen, and even erythematous at times. The pupils are contracted. A slight muco-purulent secretion from the conjunctiva appears after the subsidence of the above symptoms. Recovery takes place in a few days, with complete restoration of vision, except in rare cases.

(*b*) *Blinding of the Retina.*—This is the same affection as in blinding of the retina by sunlight. The central scotoma may persist after an attack of electric ophthalmia, or may occur without it. The injurious action of the electric light on the eye has been attributed to the chemical action of the ultra-violet rays, to the accompanying heat rays, and to dazzling of the retina. Widmark's experiments show that changes can be produced in the retina by the electric light, without any heat coagulation. These changes consist in edema, with more or less destruction of the nervous elements of the retina—namely, the outer layers, including the rods and cones, and the inner layer of nerve fibers.

Treatment.—The preventive treatment consists in the use of colored glasses. Yellow glass has been recommended by Maklakoff. In the electric welding works in Germany a combination of deep blue and red is used, while the Sheffield workers prefer several layers of ruby glass. For the rest of the treatment see the paragraphs on snow-blindness, and blinding by sunlight.

It may be as well to mention here that for domestic illuminating purposes electric light possesses many advantages over gas, in so far as the use of the eye is concerned. It has a greater illuminating power, produces less heat and no products of combustion, and hence it does not vitiate the atmosphere, or tend to cause conjunctival irritation. The electric light is much steadier than gas; and, on account of the smaller quantity of red rays which it emits, it forms a nearer approach to sunlight than does gas.

Tumor of the Retina.

Glioma of the Retina.—This is a malignant growth which is found almost exclusively in young children, and may even be congenital. It is the only growth which occurs in the retina. Owing to the age of the patients, the incipient stages of the disease are seldom observed, for they are unattended by pain or inflammation.

The growth commences as small, white, disseminated swell-

ings in the retina, usually in one or other of the granular layers, more rarely in the nerve-fiber layer. The retina is apt to become detached at an early period; but there are exceptions to this, especially when the disease starts from the nerve-fiber layer. In the early stages there is no iritis, cyclitis, or opacity of the vitreous humor, and the iris periphery is not retracted—points which especially enable us to distinguish it from pseudo-glioma (*vide* Purulent Inflammation of the Vitreous Humor, p. 384). Secondary glaucoma finally comes on. The optic nerve may become involved at an early period; but sooner or later it invariably does so, leading then by extension to glioma of the brain. When the tumor has filled the eyeball, it bursts outwards, usually at the corneo-sclerotic margin, and then grows more rapidly, and often to an immense size, as a fungus hematodes. The orbital tissues become involved, and even the bony walls of the orbit; while secondary growths in other organs, more especially in the liver, are not rare.

The diagnosis between glioma of the retina and tubercle of the chorioid (p. 294), when the latter occurs in young children, is sometimes difficult or impossible; but, in view of treatment, it is not of great importance, as in either case the eye must be enucleated.

Treatment.—The only hope of saving the patient's life lies in enucleation at an early stage, or before the optic nerve becomes diseased. It is important in removing the eyeball to divide the nerve as far back as possible; and, if the orbital tissues be already diseased, to remove all suspicious portions of them. Several cases in which there was no return of the growth have been observed.

Parasitic Disease.

Cysticercus under the Retina.—The cysticercus of the tænia solium in the eye is extremely rare in these countries, but is not so very rare in Germany. Its most frequent seat is between the retina and chorioid, where it is recognized with the ophthalmoscope as a sharply defined bluish-white body, with bright orange margin. At one point of the cyst there is a very bright spot, which corresponds with the head of the entozoön. Wave-like motions along the contour of the cyst should be looked for to confirm the diagnosis. The cysticercus may move from its origi-

nal position, and in so doing cause considerable detachment of the retina. Delicate veil-like opacities are apt to form in the vitreous humor, and are almost characteristic of the presence of cysticercus.

The entozoön may become encapsulated behind the retina; or it may burst into the vitreous humor (p. 396); and finally chronic iridocyclitis, with total loss of sight and phthisis, is apt to come on.

Treatment.—We are not acquainted with any anthelmintic which will act upon the entozoön in the eye. Removal of the cyst by operation is the only means by which the eye can be saved; and this measure can only be resorted to when the position of the cysticercus is unfavorable for it—*e. g.,* when it is close to the equator of the eyeball. By a well-placed puncture through the sclerotic and chorioid the entozoön may then be evacuated.

Detachment of the Retina.

This condition consists in a separation of the retina from the chorioid, the intervening space being occupied by a clear serous fluid. It is not usual to employ the term when it is a solid neoplasm only that lies between retina and chorioid.

If the media be clear and the detached portion extensive, the diagnosis is not difficult.

The Ophthalmoscope shows a grayish reflex from a position in front of the fundus oculi, and to the surface from which the reflex is obtained a wave-like motion is imparted when the eyeball is moved. Over this grayish surface the retinal vessels run, and they serve to distinguish a detached retina from any other diseased condition with a somewhat similar appearance. They seem black, not red, in consequence of absorption of the transmitted light, and are hidden from view here and there in the folds of the detached retina. In many cases a rent in the detached retina, through which the chorioid can be discerned, will be discovered.

The detachment may commence in any portion of the fundus, but most commonly above; yet, owing to gravitation of the fluid, it ultimately settles in the lower half of the fundus, and hence this is the most common place to find it, the part first detached having become replaced. The diagnosis is more difficult if there be but little fluid behind the retina, or if there be opacities in the vitreous humor.

Vision is affected according to the position and extent of the detachment. Central vision may be quite normal if the macula lutea and its immediate neighborhood are intact. The patients complain of seeing objects distorted, of a black veil which seems to hang over the sight, and sometimes of black floating spots before the eye, due to opacities in the vitreous humor. These symptoms often come on suddenly in an eye which has hitherto had good sight.

The field of vision, on examination, will show a defect corresponding to the position of the detachment. If, for example, it be below, the defect will be in the upper part of the field. If the detachment be fresh, the retina not having yet undergone secondary changes, and if the quantity of subretinal fluid be not great, the defect in the field may only amount to an indistinctness of vision; while later on, when infiltration and connective tissue degeneration of the detached part come about, fingers may not be counted at the same place. The phosphenes * of the detached portion are wanting.

Should the detachment become complete, little more than mere power of perception of light may be present. Total detachment is followed by cataract, and often by iritis, cyclitis, and phthisis bulbi. The detachment may remain stationary, and not extend to the whole fundus, or the retina may return to its normal position; but such a happy event is most rare.

Causes.—Myopic eyes—which we know are so frequently affected with chorioiditis and disease of the vitreous humor—are those most subject to detachment of the retina; but idiopathic detachment occurs also in eyes which are apparently healthy. Blows upon the eye may produce detachment, the retro-retinal fluid being serous or bloody. Some punctured wounds of the sclerotic, also, in the course of healing, by dragging on the retina, give rise to it. Chorioidal tumors, especially those situated in the posterior segment of the fundus, usually cause detachment in

* Phosphene is the subjective sensation of light experienced when the eyeball is pressed upon. For clinical purposes it is best tested by gentle pressure with a blunt point (head of a bodkin or large-sized probe) applied to the eyeball through the eyelid. The phosphene of any region is tested by applying pressure to that part of the globe. Thus, if in a healthy eye the individual look down, and pressure be applied to the upper part of the globe through the eyelid, the phosphene will be seen appearing below; but if there be a detachment of the retina at the place pressed on, no phosphene is seen.

an early stage of their growth, and the complication renders their diagnosis more difficult (p. 292).

Leber [7] observed that in non-traumatic detachment a perforation or rent in the detached portion is very frequently to be seen with the ophthalmoscope, and holds that it is probably always present, although sometimes, from being hidden behind a fold of the retina, it cannot be found. He was led from this, and from his pathological investigations and experiments upon animals, to think that the detachment was due to shrinking of a diseased vitreous, which first became slightly separated from the retina, and that then—at some place where the retina and hyaloid had become adherent from the inflammatory process—a rent was produced in the retina by the shrinking process in the vitreous. He concluded that through this rent the fluid, which is always present behind the vitreous in cases of detachment of that body, makes its way behind the retina, and separates the latter from the chorioid. Nordenson's pathological researches [8] went to corroborate this. He ascertained, too, that disease of the ciliary body and chorioid is the primary cause, although we may not be always able to detect it with the ophthalmoscope, and that the pathological change in the vitreous humor consists in an alteration in its connective tissue elements, resulting in the deleterious shrinking.

Raehlmann,[9] however, from the results of experiments, and also from clinical observation, concludes that detachment of the retina is due to exudation from the chorioidal vessels of a fluid which is more albuminous than the fluid in the vitreous humor. Hence, he thinks, diffusion takes place through the retina, and a greater quantity of the less albuminous vitreous fluid passes through the retina, thus producing and increasing the detachment. Rupture of the retina is not, in his view, a necessary factor in the causation, but it may occur if the tension behind the retina be higher than that in front of it.

Treatment.—Evacuation of the subretinal fluid by puncture of the sclerotic was proposed by Sichel, and cultivated by de Wecker. The instrument resembled a broad needle, with a sharp point and two blunt edges, which is entered through the sclerotic and chorioid at a place corresponding to the position of the detachment, but not so deeply as to reach the retina, lest thereby it be further displaced. The instrument is then given a quarter of a rotation, to make the wound gape, so as to admit of the flowing off of the fluid. If possible a position near the equator of the

globe, and between two recti muscles, should be selected for the operation. Moreover, the incision should lie parallel to the direction of the orbital muscles, so that the chorioidal vessels may be injured as little as possible. A firm bandage is applied, and the patient kept in bed for eight or ten days.

The dorsal position in bed, with a pressure bandage on the eye, and diaphoretics internally, the treatment being continued for from four to six weeks, brings about reposition of the detachment in some cases. The method, if properly carried out, is trying to the patient, but it is only rarely of use.

The cures which have been accomplished by these means probably depend upon the retina again coming in contact with the chorioid, and, owing to some slight inflammatory process, adhering to it. For the most part the cure is but temporary, and in such cases we may suppose that no adhesion sprang up, but that the temporary cure was due to a return of the subretinal fluid, through the hole in the retina, to its original position between the retina and vitreous. Soon, however, it makes its way back again through the opening, and the detachment recurs.

To promote adhesion between retina and chorioid, Dor [10] touches the sclerotic corresponding to the detachment lightly with a small cautery, injects rather strong (10 per cent.) solutions of common salt under the conjunctiva, and keeps the patient in bed. He reports some cures by this method.

Schoeler [11] injects tincture of iodin into the vitreous humor in front of the detached retina, in order to press it back to the chorioid, and to produce a plastic chorioido-retinitis, which may unite the two coats. He has reported several good results by this method, but some who have tried it have experienced violent inflammatory reaction in the eyes operated on, with disastrous consequences, and the treatment is discredited.

Deutschmann [12] has proposed the following method of treatment. Adopting the theory of retraction, he passed a double-edged narrow knife through the sclerotic, chorioid, and retina, into the vitreous humor, dividing cords and liberating fluid from before and behind the retina. The vitreous of a rabbit's eye diluted with a few drops of a 7 to 100 solution of chlorid of sodium was triturated. Of this a few drops were injected with a hypodermic syringe into the vitreous cavity of the eye. By this procedure it was intended to press the retina permanently against the chorioid, until the ensuing uveitis had established agglutination

of the retina to the chorioid. In a good many cases where this operation has been tried it has done harm rather than good.

Grossmann [13] tried aspiration of the subretinal fluid, with simultaneous increase of the pressure in the vitreous humor, by injections into the latter of four or five drops of an indifferent fluid, namely, a 0.75 per cent. tepid solution of common salt. The results obtained were encouraging in the three cases reported.

Galezowski [14] simply aspirates the subretinal fluid.

Electrolysis has been tried.

Pilocarpin used hypodermically has been praised by some as a mode of treatment, as, also, salicylate of sodium internally.

Formerly an active mercurial treatment used to be employed, with the object of obtaining absorption of the fluid.

The Prognosis of every case of detached retina is bad, spontaneous cure being extremely rare, and the cures by any one or by any combination of the above methods of treatment being few and far between; and when the retina does return to its place, there is the danger of a recurrence of the detachment. Moreover, both eyes are often affected, one after the other. The most favorable cases are those due to chorioiditis, the most unfavorable those due to posterior staphyloma.

Traumatic Affections of the Retina.

In addition to detachment and rupture of the retina, the undermentioned conditions occur as the results of injuries.

Traumatic Anesthesia of the Retina.—A blow on the eye from a fist, cork from a bottle, etc., is liable to produce considerable amblyopia, with concentric contraction of the field, which may continue for a long time, while the *Ophthalmoscopic Appearances* are normal. Ultimately these cases usually recover, an event which may be decidedly promoted by the use of strychnin hypodermically; but very defective sight sometimes remains permanently.

Commotio Retinæ, or **Traumatic Edema of the Retina,** is the result of a blow upon the eye. Immediately after the blow there is marked episcleral injection, and the pupil can be dilated but slowly with atropin. Within a few hours after the accident the *Ophthalmoscope* reveals a white cloudiness of a portion of the retina, usually in the neighborhood of the optic papilla and

macula, but sometimes more eccentrically; and sometimes there are two opaque patches. The opacity increases in intensity, and spreads somewhat. The retinal vessels remain normal; there may be some small hemorrhages, and sometimes the papilla is redder than normal. These appearances completely disappear in the course of a few days. Vision is only slightly affected, and recovers according as the retinal changes pass off.

References.

[1] "Von Graefe's Archiv," xxxix. 3. p. 227.
[2] "Arch. f. Ophthal.," xxvii. p. 484.
[3] "Roy. Lond. Hosp. Rep.," 1875. "Ophthalmic Review," v. p. 1.
[4] "Trans. Ophthal. Soc., U. K.," vol. iii. p. 108.
[5] "Centralbl. f. Augenheilkunde," 1883, p. 392.
[6] "Journ. Tropical Medicine," November, 1898.
[7] "Bericht d. Ophthal. Gesellsch." Heidelberg, 1882, p. 18.
[8] "Die Netzhautablösung." Wiesbaden, 1887.
[9] "Archiv für Ophthal.," xxvii. part 1. p. 1.
[10] "Bulletins de la Société francaise d'Ophtalmologie," 1895, 1896.
[11] "Zur operativen Behandlung und Heilung der Netzhautablösung." Berlin, 1889.
[12] "Beiträge zur Augenheilkunde," xx. p. 1.
[13] "Ophthalmic Review," 1883, p. 89.
[14] "Recueil d'Ophtalmologie," Mars, 1888.

CHAPTER XVI.

DISEASES OF THE OPTIC NERVE.

Optic Neuritis.— *The Ophthalmoscopic Appearances* of inflammation of the optic nerve vary a good deal with the intensity of the process. Common to every case is hyperemia and swelling of the papilla, with haziness (so-called "woolliness") of its margins, and increase in the size of the central vein, while the central artery remains of normal dimensions, or is contracted. The swelling and haziness extend but a short distance into the surrounding retina, and the distention of the vein is also not continued to the periphery of the fundus. In slight cases these appearances may barely exceed the normal.

In extreme instances the papilla is swollen to a great size, and may even assume quite a dome shape, while the veins are enormously distended and tortuous, and the arteries are contracted so as to be barely visible. Grayish striæ extend from the papilla into the surrounding retina, some flame-shaped hemorrhages are present on or near the papilla, and, occasionally, white spots in the retina, and a stellate arrangement of small white dots about the macula lutea produce an appearance which cannot be distinguished from albuminuric retinitis. This extreme form is still sometimes termed Congestion Papilla, or Choked Disc (*Steuungspapille*), although the theory which originally suggested the term has been abandoned. Papillitis (Inflammation of the Optic Papilla) is a better term, expressing, as it does, more truly the pathological condition.

The Vision, even in cases where the ophthalmoscopic signs are highly developed, is frequently but little below the normal; while, again, in other, and possibly less well-marked cases, in so far as the appearances are concerned, it may be reduced to perception of light, or even that may be wanting. When due to cerebral tumor, the neuritis appears as a rule before the vision becomes affected. These remarkable differences in the degree of blindness depend, probably, on the extent to which the nervous ele-

ments of the inflamed part are pressed on or altered, and this cannot be gauged by the ophthalmoscopic appearances.

Sometimes the field of vision is normal, while again it is concentrically or irregularly contracted, or it may be hemianopic.

An attack of temporary loss of sight is a very common symptom in cerebral tumors; it may occur several times a day, and may last from a few minutes to half an hour.

Pathologically, the changes in the papilla consist in venous hyperemia, edema, hypertrophy of the nerve fibers, infiltration of lymph cells, and development of connective tissue. Inflammatory changes, although less pronounced, are also present in the trunk of the nerve and its sheaths.

Causes.—Inflammation of the optic nerve is most commonly found in connection with coarse encephalic disease. A Cerebral Tumor (including syphiloma, tubercle, and abscess) in particular is the most common cause, and is, moreover, usually present when the papillitis is of an intense kind (choked disc). The neuritis, except in very rare instances, is bilateral, and it is one of the general symptoms of cerebral tumor. Hemianopsia may coexist as a localizing symptom, if the visual center or fibers on one side be involved. Even a small tumor situated anywhere in the brain is capable of producing optic neuritis, although unattended by meningitis. Cerebral cysts do not often cause it.

Tubercular Meningitis is the next most common cause. Non-tubercular meningitis occasionally gives rise to optic neuritis, and sometimes, also, cerebrospinal meningitis does so.

The Connection between Optic Neuritis and Intracranial Diseases has given rise to much discussion. In cases of tumor, as well as of tubercular meningitis, a considerable exudation of fluid usually takes place into the cavity of the third ventricle. This, along with the new growth, or alone in cases of meningitis, increases the pressure within the cranial cavity. By reason of this increased intracranial pressure, the subarachnoid fluid is believed to be driven into the subvaginal lymph space of the optic nerve, and to produce there that dropsy of the sheath which is found in nearly all these cases on careful postmortem examination.

Leber holds [1] that this fluid probably contains a phlogogenic substance. It seems probable that the reason why some small cerebral tumors may cause optic neuritis, while some large ones do not, is to be sought in the fact that the former may happen to be rapidly growing tumors, and accompanied by much ventricular

dropsy, while the larger tumors may be slow in growth, and attended by but little dropsy of the ventricles. Moreover, the fluid driven into the subvaginal sheath of the optic nerve may not be equally rich in phlogogenic substance in the case of every tumor.

The inflammation, although most intense at the papilla, near which the fluid is collected in greatest quantity in the *cul-de-sac* formed by the termination of the intervaginal spaces, is not confined to that place, as was believed, but extends up the trunk of the nerve, as microscopic examination reveals.

Many observers state that in a large number of cases cerebritis, recognizable only with the microscope, is present, and that an extension of this process down the optic nerve takes place. They have ascertained that the whole trunk of the nerve is involved in the inflammation, and they seem to regard the dropsy of the sheath as of little or no importance in the causation of the optic neuritis.

Again, others maintain that edema, but not inflammation of the optic trunk is conducted from the brain.

Other causes for Optic Neuritis are:

Hydrocephalus.—Here the pathogenesis is probably the same as in the foregoing; but the occurrence of optic neuritis is, on the whole, not very common in this connection.

Tumors of the Orbit.—The path by which these growths bring about papillitis is still unknown.

Inflammatory Processes in the Orbit, such as caries, inflammation of the retro-orbital areolar tissue, erysipelas of the head and face extending to the orbital tissues, and periostitis. The presence of the latter may often be recognized by pain on motion of the eyeball, pain in the eye and forehead, and especially by pain on pressure of the globe backwards, and is frequently of rheumatic origin. Often in these cases one eye only is affected; and, although the *Ophthalmoscopic Appearances* are sometimes very slight, yet vision may be quite lost in a few hours or days, atrophy of the nerve then rapidly setting in.

Very many of the cases, however, do not go on to atrophy, but end in recovery of useful vision.

Exposure to Cold, especially if the skin be heated and perspiring.

Suppression of the Menstruation.—If during the menstrual period the flow be arrested by exposure to cold, wet feet, etc., acute optic neuritis with rapid blindness may come on. Spontaneous

amenorrhea, or even irregularity of menstruation, and the climacteric period are liable to have a similar, but more chronic, result. Nothing is known with regard to the connection between the uterine and ocular disorder. In these cases the Ophthalmoscopic Appearances, as well as the blindness, are apt to be extreme. Treatment here should be directed chiefly to restoring, when possible, the normal uterine functions. Hot foot-baths and Heurteloup's leech to the temples are of use.

Chlorosis.—Here optic neuritis often is present, due to the disordered state of the blood, and usually yields under the influence of iron.

Syphilis.—The trunk of one or both optic nerves may be the seat of specific inflammation in connection either with congenital or with acquired syphilis, but this primary specific optic neuritis is a relatively rare disease. In cases of acquired syphilis it makes its appearance in from six months to two years after the inoculation.

The Ophthalmoscopic Appearances may be normal (retrobulbar neuritis), or may present any grade of neuritis, even to the most pronounced papillitis. In the latter case it would not be possible to say whether the papillitis is a primary one, or is due to a syphilitic gumma within the cranium. The inflammation often extends as far up as the chiasma.

The Treatment in these cases of specific papillitis must be active mercurialization. By this treatment, even if perception of light be lost for a period of not more than eight to fourteen days, hopes may be entertained of its complete or partial recovery.

Cases of double optic neuritis of syphilitic origin have been observed, in which complete recovery took place, the papilla returning to its normal condition. But, as a rule, some optic atrophy, at the least, with slight concentric contraction of the field, results. The prognosis is all the better the sooner the optic neuritis follows upon the primary syphilitic affection.

Rheumatism.—There is no doubt whatever but that the rheumatic diathesis is occasionally the cause of optic neuritis, although the fact is not unreservedly accepted by every author. Other manifestations of rheumatism are sometimes well marked, but may be slight—*e. g.*, in a case which I saw, neuralgia of the face and head in damp weather, and even with a shower of rain, was the only other sign of rheumatism. One or both optic nerves may be attacked.

The Ophthalmoscopic Appearances often amount to extreme papillitis, but in many cases fall short of this.

If the case come early under suitable treatment the Prognosis is fairly favorable; but when the inflammation is of some standing, consecutive optic atrophy must be feared.

The Treatment consists of full doses of salicin, salicylate of sodium, iodid of potassium or of sodium, Turkish baths, and other recognized antirheumatic measures.

Lead-Poisoning.—In some cases of lead-poisoning optic neuritis, not to be distinguished from that of primary cerebral affections, is found. Sometimes the Ophthalmoscopic Appearances are very slight, and, again, quite pronounced, the changes extending into the retina. They sometimes simulate the retinitis of Bright's disease; and in such cases renal disease is likely to have much to do with the causation of the retinitis. Indeed, there are those who, with good opportunities for forming a correct opinion, deny the existence of a specific lead neuritis, and hold that the neuritic affection in all such cases is to be referred to albuminuria, to effusion into the ventricles of the brain and subrachnoid space, or to accompanying suppression of menstruation. Occasionally optic atrophy is the first ophthalmoscopic appearance seen; but it is probably consecutive to retrobulbar neuritis, as shown by white striæ (perivascularitis) along the vessels.

The Vision is often much affected, and it is stated that sudden complete blindness in connection with an intercurrent attack of lead colic may appear and pass off again. Consecutive atrophy is liable to come on, and then vision may be seriously and permanently damaged.

The Diagnosis depends entirely on the presence of the other well-known symptoms of lead-poisoning, the Ophthalmoscopic Appearances presenting nothing pathognomonic.

The Treatment is that for general lead-poisoning, or for the immediate cause of the neuritis.

In Peripheral Neuritis optic neuritis is occasionally found.

Multiple Sclerosis.—In these cases the inflammation is very ephemeral, and rapidly gives place to atrophy. Uhthoff states that it occurs in about 13 per cent. of the cases of this disease.

Tabes Dorsalis.—A few cases of this disease are published in which optic neuritis was present. It is probable that the latter depended on coexistent syphilitic cerebral disease, rather than on

the spinal disorder as such. In Acute Myelitis inflammation of the optic nerve is sometimes seen, so that optic neuritis with paralytic phenomena does not exclusively indicate cerebral disease.

Hereditary and Congenital Predisposition.—It has been observed that optic neuritis, without immediate cause, may attack several members of a family, and that the tendency to it may extend over several generations. It makes its appearance about puberty or a little later in males, and about the climacteric in females, but is much rarer in the latter. The patients may be perfectly healthy in all other respects, but many of them suffer from other affections of the nervous system. Both eyes are affected, the defect of vision being a central amblyopia (central scotoma), from which recovery is rare; but yet, although the Ophthalmoscopic Appearances gradually become those of atrophy, the peripheral portions of the field retain their functions. Optic neuritis in this form was first thoroughly described by Leber,[2] and is generally known as Leber's disease.

As to the treatment of these cases, due to hereditary and congenital predisposition, Mooren has employed a seton in the back of the neck in the early periods, and, later on, nitrate of silver internally. Leber has found benefit from a mild course of mercurial inunction.

Optic Neuritis also occurs occasionally in fevers; it has been observed in Measles, Scarlatina, Typhoid, and Malaria. It may follow Influenza, causing contraction of the field of vision or central scotoma which usually disappear, but, on the other hand, it may lead to optic atrophy.

The two following diseases—Toxic Central Amblyopia, or Central Scotoma and Optic Neuritis with Persistent Dropping from the Nostril—must be treated of separately, owing to the well-defined etiology of the one and the peculiar symptoms of the other.

Toxic Central Amblyopia, or Central Scotoma.—

Symptoms.—The affection of vision often comes on rather rapidly. The patient may complain of a glimmering mist which covers all objects, especially in a bright light, and the acuteness of vision is reduced. The patient generally states he can see better in the dusk than in bright light. At the commencement there is no defect in the field of vision, but simply a general dimness of vision. At a somewhat later stage examination of the field dis-

covers no defect for a white object; yet, if a small pale green object be employed, it will generally be ascertained that, at a region close to the point of fixation, the color is not recognized, but seems gray or white; pink may seem blue, and red may appear brown or black; while in other parts of the field the colors are recognized up to their normal boundaries. This is a central color-scotoma. As the disease advances a white object will be but indistinctly seen in the scotoma; and in some rare cases all power of perception within its area may be lost, even the flame of a candle not being recognized. Hence the name Central Amblyopia. The scotoma is usually oval in shape, its long axis horizontal, and extends from the fixation point towards the blind-spot of Mariotte (paracentric scotoma); but occasionally it is of much larger dimensions, and sometimes surrounds the fixation point (pericentric scotoma).

Even when the scotoma is very pronounced it remains negative—*i. e.*, it is not observed by the patient as a dark spot in the field, as is a scotoma due to disease in the outer retinal layers. The affection is almost always binocular, and as a rule there is but little difference between the vision of the two eyes.

The Prognosis of the disease is slow, occupying weeks or months. Restoration of normal vision usually takes place if the defect of vision, although of extreme degree, be not of old standing. In the latter case, while recovery of central vision cannot be expected, the functions in the periphery of the field are usually maintained; and, consequently, these people, although incapacitated from reading, writing, and other fine work, do not lose their power of guiding themselves.

Causes.—With but few exceptions the subjects of this disease are men, probably because their habits and modes of life expose them, more than women, to the influences which produce it. These are: Exposure to cold and wet; cold blasts on the body, especially the heated face (Samelsohn); but the most common cause is excess in the use of alcohol, or of tobacco (toxic amblyopia), or of both. I have observed that the kind of alcoholic indulgence most likely to develop the disease is the frequent drinking of small doses of the stimulant. The individual who often gets thoroughly intoxicated, and between times drinks but little, is less liable to central amblyopia than he who, although never incapable of transacting his business, takes many half-glasses of whisky or brandy during the day. Dyspepsia and loss of appe-

tite are constantly present in these cases. Other signs of chronic alcoholism need not be present, but one often sees trembling of the hand and head, sleeplessness, and even delirium tremens. The kind of tobacco most likely, when used in excess, to give rise to central amblyopia is shag or twist. Other kinds of pipe-tobacco and cigars may cause it, but I have not known of a case due to cigarette-smoking.

Excess in alcohol is usually combined with excessive smoking, but cases of pure alcohol-amblyopia certainly do occur—although some English authors deny it—as well as pure tobacco-amblyopia. The most common age for tobacco-amblyopia is from thirty-five to fifty—a time of life when men do well to give up, or to very much reduce, their use of tobacco, as well as of alcohol.

Central amblyopia has also been observed in diabetes, in poisoning from bisulphid of carbon,[3] so largely used in the manufacture of india rubber, from dinitro-benzol,[4] used for explosives, and in iodoform poisoning.[5]

The Ophthalmoscopic Appearances in the beginning are either quite normal or there is slight hyperemia of the papilla and retinal vessels; or, in addition, there may be slight indistinctness of the margins of the papilla, and sometimes white striæ along the vessels, especially before they leave the papilla. All the primary appearances, if any be present, soon pass away, and give place to a grayish whiteness of the temporal side of the papilla, while the nasal portion remains of normal appearance, as do also the vessels. At a very advanced stage, in some cases, the whole papilla presents the appearance of white atrophy.

The Pathological Changes observed by Samelsohn, Nettleship, and Walter Edmunds, and Uhthoff, in the optic nerve, consist of an interstitial nephritis at its axis, commencing so high up as the optic foramen, and leading to proliferation of connective tissue and to secondary descending atrophy of a certain bundle of nerve fibers. These are the fibers which supply the region of the macula lutea. The changes are analogous to those which take place in the liver and brain as the result of chronic alcoholism.

Treatment consists, above all, in total abstinence from the poison in question. The patients are generally ready to promise this, but they often do not act up to their intentions. When they do so, improvement rapidly takes place in most cases which are not too

far gone, without any other treatment; but the cure may be promoted by the use of iodid of potassium in large doses, Heurteloup's artificial leech or dry cupping to the temples, hot foot-baths, and Turkish baths. Strychnin hypodermically (1-30 grain daily) in the temple is often of use, and phosphorus and strychnin may be given internally. Whatever remedy be used internally, care should be taken that it does not produce or increase dyspepsia; and it may be necessary to restrict the internal medicine for a time, or altogether, to a stomach tonic. Sleeplessness should be combated with sulphonal or bromid of potassium. Treatment may have to be continued for some weeks, before a cure may be noted.

Retrobulbar Neuritis.—This is by no means a very common disease. It is ushered in by rapid loss of sight in one eye, sometimes in both, or they may be attacked with a considerable interval between. Examination of the field of vision discovers a central scotoma, which is often absolute. At the commencement, pain in the orbit is complained of, the motions of the eye are somewhat painful, and there is pain on moderate pressure of the globe backwards into the orbit. Often at first there are no ophthalmoscopic changes, but after a time marked optic neuritis shows itself, and this may pass into atrophy, or atrophy may appear without any previous neuritis which can be discerned. It is rare for complete and absolute amaurosis to result, although the optic disc remains white. In most of the cases the central scotoma disappears, and almost normal vision is restored; but in some a more or less well-marked central scotoma, with defective sight, remains. The prognosis is all the better if the case comes under treatment very early.

Exposure to severe blasts of cold wind on the head, rheumatism, and influenza are the most common causes.

Treatment.—Iodid of potassium in large doses, mercury, and salicylate of soda.

Optic Neuritis associated with Persistent Dropping of Watery Fluid from the Nostril.

Twenty-one cases of this remarkable disorder, including a case of his own, have been collected by St. Clair Thomson [6] in a valuable monograph, and others have since then been reported. In eight of these cases the eyesight was affected, there being optic neuritis or secondary atrophy. The patients suffered from a persistent watery discharge from the nose (usually the left nostril),

with more or less severe cerebral symptoms: violent headache, epileptic attacks, vomiting, stupidity, sleepiness, unconsciousness, delirium, weakness of the lower extremities, and a high degree of amblyopia, or even blindness, of both eyes, due to papillitis followed by atrophy. The severity of the head symptoms varies very much in different cases. Headache is the most constant of these symptoms, but even it was absent in one case. In Leber's case, moreover, there was loss of smell, and in Nettleship's case palpitation of the heart and prominence of the eyes. The fluid which runs from the nostril is identical in its analysis with that of the cerebrospinal fluid. If it occasionally ceases to flow, the cerebral symptoms are brought on, or increased in violence. Leber's case was one of internal hydrocephalus, and the other cases were probably of similar nature. He regards the fluid as coming from the third ventricle, through a small opening in the ethmoid bone; or the fluid possibly passed from the subdural space along the lymph spaces, which surround the olfactory nerves.

The affection commences usually in early adult life, and no rational treatment for it has been suggested. The flow may cease spontaneously for periods varying from a few hours to several months. In some cases it ceased altogether, or at least had not recurred after an intermission of five or even fourteen years. Most of the cases have been lost sight of, but some are recorded as having died of meningitis.

Atrophy of the Optic Nerve.—This disease may be secondary to some other optic nerve or retinal affection, or it may be a primary disease. *The Vision* is seriously affected, and complete blindness is the usual result. With the *Ophthalmoscope* the optic papilla is seen to have lost its delicate pink color, and to have become white or grayish, while it is often cupped, and the vessels are apt to be diminished in caliber.

Secondary Atrophy of the Optic Nerve may result:

1. *From Optic Neuritis.*—The ophthalmoscopic appearances consist in a white or grayish-white color of the papilla, with very diminished retinal vessels; and along both sides of the vessels, far into the retina, are seen white lines, which sometimes even obscure the vessels, and which are due to hypertrophy of their coats. The diminution in caliber of the vessel is a sign of neuritic atrophy, but is not always present, and is moreover found with other forms of atrophy. Other signs of this form, also not con-

stant, are: a certain opacity of the papilla, and that the lamina cribrosa is not generally visible, owing to development of connective tissue at the papilla. It is evidently not always possible to recognize any given case as of neuritic origin.

Symptoms.—The acuteness of vision is lowered, and as a rule the field of vision becomes contracted, usually more at the nasal than at the temporal side. Subsequently the temporal side of the field becomes contracted, and finally a small eccentric portion of the field to the temporal side may be all that remains, or even this may disappear, and absolute amaurosis result. The color-vision is always much affected. The light-sense is affected, so that there is diminished sensibility for differences of illumination; while, in chorioido-retinal diseases, there is defect in the quantitative perception of light, the minimum quantity being larger than normal.

2. *From Pressure.*—This may be brought about by a tumor anywhere in the course of the nerve, by inflammatory exudations, by a splinter of bone in cases of fracture of the skull, and, also, by pressure upon the chiasma by the floor of the distended third ventricle in cases of internal hydrocephalus.

3. *From Embolism of the Central Artery of the Retina.*—In these cases the contraction of the vessels is usually extreme.

4. *From Syphilitic Retinitis, Retinitis Pigmentosa, and Chorioido-retinitis.*—The vessels here are much attenuated, and the altered color of the optic disc is a dull yellow rather than white or gray.

Primary Optic Atrophy is often found associated with:

Disease of the Spinal Cord (Spinal Amaurosis), especially locomotor ataxy. Optic atrophy is often an early symptom in the latter disease; but, again, it may not come on until the affection of the gait is well pronounced, while in other cases it is never present at all. It is a remarkable and important fact, first pointed out by Benedikt of Vienna, that there is an antagonism between atrophy of the optic disc and the other symptoms of tabes dorsalis. It is rare for a tabetic patient, in whom optic atrophy comes on in an early stage of his disease, to become ataxic; and, frequently, in these cases, when the blindness has advanced, the pains, too, become less severe. But if amaurosis does not come on until the ataxy is well developed, no improvement in the latter is likely to be noted.

Atrophy is found more rarely with insular sclerosis and lateral sclerosis of the spinal cord; and in general paralysis of the

insane, although spinal disease is not always present, atrophy of the papilla frequently occurs.

It is probable that the disease commences at the papilla in spinal cases. The ophthalmoscope displays a papery-white or bluish-white papilla, which in advanced stages often becomes cupped. The retinal arteries are usually extremely reduced in caliber, and the veins, too, may be small; but, again, the retinal vessels may differ but little, or not at all, from the normal.

Symptoms.—Central vision is affected at an early stage in the disease, and eccentric contraction of the field usually appears at the same time. The contraction may be concentric, or it may be more marked in one direction than another, and opinion is divided as to the direction commonly first involved. This concentric contraction advances gradually towards the center of the field from every side, until it finally engulfs the fixation point.

Occasionally the affection begins as a central scotoma, accompanied by eccentric defects of the field. Color-blindness is an almost constant symptom. As a rule, absolute blindness is brought about in the course of a year or two.

Primary Optic Atrophy of the progressive form just described may occur *As a Purely Local Disease,* without any other defect in the system. The prognosis for the sight in such cases is as bad as in spinal cases.

Treatment.—In neuritic atrophy, so long as there are still signs of active inflammation, antiphlogistic measures—Heurteloup's leech to the temple, hot footbaths, rest of body and mind, dark room, iodid of potassium, and, especially, mercury internally, when otherwise admissible—are to be adopted. At a later period hypodermic injections of strychnia (1-30 gr., increased gradually to 1-20 or 1-18 gr. once a day), and galvanism may be tried. Hypodermic injections of antipyrin (about 7 1-2 grains every second day) have been given with some benefit in these cases.

In spinal amaurosis, and in optic atrophy occurring as a local disease, strychnia hypodermically and the galvanic current sometimes improve vision for a time. Phosphorus internally may be given.

The treatment for optic atrophy, due to causes 2, 3, and 4, is to be directed to the primary disease.

The Prognosis is very serious; for, although every therapeutic measure may have been employed, amaurosis is the ultimate result as a rule.

Tumors of the Optic Nerve.— These are rare affections. They occur at all times of life, but most of the patients are under twenty years of age. The tumor usually commences about the middle of the course of the nerve, and often does not extend to the bulbar end. The symptoms are: Slowly increasing protrusion of the eyeball, in a direction most usually directly forwards, or forwards and outwards. The motions of the eyeball are not greatly restricted, and the center of its rotation is not displaced, owing to the tumor being within the cone of the orbital muscles. The proptosis is unaccompanied by pain. The sight becomes very defective, or is quite lost at a very early stage, from interference with the functions of the nerve by the tumor, or by the optic neuritis, or optic atrophy, to which it gives rise. The tumor is sometimes very soft, so that the eyeball can, as it were, be pushed back into it, and the pressure does not cause pain. The pupil reacts consensually. The tumor may often be felt by palpation in the orbit. These tumors being benign, the patient's health does not suffer.

The diagnosis between a new growth of the optic nerve and one of its sheaths can hardly be made with certainty; but the existence of fairly good vision, while other symptoms are as above, at a rather late period of the proptosis, would point to the sheath as the seat. Such a diagnosis would be important, for, as a brilliant case of Dr. Antill Pockley's [7] shows, it may be possible to remove, by Krönlein's operation, a tumor of the sheath of the optic nerve, while preserving not merely the eyeball, but good vision as well.

These tumors are either myxo-sarcomata, or, less frequently, endotheliomata, and are usually encapsuled by the sheath of the nerve. They are benign, in the sense that they do not lead to glandular enlargements or to metastases; but in rare cases they extend into the cranial cavity.

Treatment.—To remove these tumors there are three methods, namely: Removal of the eyeball with the tumor; Krönlein's operation, by means of which the optic nerve tumor may, probably in most instances, be removed without the eyeball; and Knapp's operation, also for removal of the tumor without the eyeball. It is unnecessary to describe the first of these procedures, which follows very much the lines of an ordinary excision of the eyeball, except that the optic nerve is divided as far back in the orbit as possible. Krönlein's operation will be described in the next chapter.

Knapp's operation is as follows: The tendon of the internal rectus is divided so as to leave a portion adherent to the sclerotic of about 5 mm., the cut end being secured by a suture passed through it, to prevent it from retracting into the orbit. The eyeball is then forcibly everted outwards, a strong scissors is passed into the orbit, and the optic nerve is divided as close to the optic foramen as possible. The globe is now further everted outwards, to expose its posterior surface with the tumor attached, and the latter is removed by dividing the optic nerve close to the eyeball. Finally, the eyeball is reposed, the cut ends of the tendon of the muscle united, and the opening in the conjunctiva closed. A drawback to this operation is that it is not always possible to be certain that the deep portion of the tumor is reached with the scissors.

Hyaline, or Colloid, Outgrowths from the optic papilla are occasionally met with. Seen with the ophthalmoscope, they present the appearance of bluish-gray, mulberrry-like nodules. According to Iwanoff's [8] investigations they originate in the lamina vitrea of the chorioid at the margin of the papilla, or within the area of the papilla; for the lamina vitrea is often prolonged into the papilla, and takes part in the formation of the lamina cribrosa. These outgrowths do not always cause a defect of sight, and rarely cause serious blindness. It is often found that a blow upon the eye has been received some time previously, and it is probable that such a trauma may have to do with the growth by rupturing the very brittle lamina vitrea.

Treatment is of no avail.

Injuries of the Optic Nerve.— In addition to those injuries which result from direct violence with sharp instruments, etc., entering the orbit, the optic nerve may be injured by falls on the head. Fractures of the base of the skull frequently involve injury to the optic nerve. But even where no fracture occurs, blindness with atrophy of the optic nerve may come on, usually only in one eye; and in these cases concussion of the nerve at its passage through the optic foramen, or an extravasation of blood in the sheath of the nerve, is probably the direct cause of the atrophy.

Hemorrhages from the Stomach, Bowels, or Uterus are capable of giving rise to serious and incurable blindness.

Blindness during or immediately after a severe hemorrhage is probably due to insufficient blood-supply to the nerve-centers

and retina, accompanying general exhaustion of the system. For such cases the prognosis is favorable.

But there is another class of cases of very much more serious import. In these the defect of vision does not come on for from two to fourteen days after the hemorrhage, when the general system is recovering. Even comparatively slight hemorrhages, which caused no marked anemia, are said to have been followed by blindness. The connection between the loss of blood and of sight in these cases is not yet clearly made out. Leber inclines to the belief that the blindness here is due to an extravasation of blood at the base of the skull and into the sheath of the optic nerve; but even then the relationship between this and the stomachic or uterine hemorrhage is not made clearer. Papillitis has been several times noted with the ophthalmoscope in these cases; and this circumstance makes it probable that neuritis is the immediate cause of blindness—even in those cases which show no ophthalmoscopic sign of it—and hydremia may be presumed to be the influence which calls forth the neuritis.

The Defect of Vision may be but slight, or it may amount to absolute amaurosis. Both eyes are usually affected in equal degree. But cases have been observed in which one eye was completely amaurotic, while the vision of the other eye was quite normal; and one such case is sufficient to prove that the lesion is peripheral—in fact, that it lies in each instance on the distal side of the optic chiasma. The field of vision is frequently contracted, either concentrically or segmentally; and even when central vision recovers the field may remain contracted.

The Ophthalmoscopic Appearances which are present immediately on the occurrence of the blindness have not as yet been observed. A few weeks later they have been found to be different in different cases. They have been found at this period normal; or presenting slight paleness of the papilla and contraction of the arteries; or there was marked paleness of the papilla, and the arteries were extremely contracted, with slight distention of the veins; or paleness of the papilla was present, but its margins were indistinct, and the surrounding retina somewhat swollen, while the retinal vessels were normal. Small hemorrhages have repeatedly been seen in the neighborhood of the papilla. At later periods well-marked optic atrophy is frequently observed.

Prognosis.—If in the beginning the defect of vision be merely amblyopia, and not complete blindness, hopes may be entertained

of marked improvement or of complete recovery. But Mooren has seen slight amblyopia pass into permanent amaurosis.

Hemorrhages from the stomach are those which are followed by the most complete and permanent blindness, while uterine hemorrhages are more commonly followed by less serious degrees of blindness.

The Treatment must consist of internal remedies calculated to correct the general anemia, such as iron, beef-tea, and meat extracts, wine, etc. Strychnin hypodermically, to stimulate the nerve, may be employed.

Glycosuric Amblyopia.—In addition to the retinal affections dependent upon diabetes, we recognize the occasional occurrence in that disease of defects of vision which are referred to disorder of the optic nerve, and which are not always accompanied by ophthalmoscopic changes. These defects of vision are found in the form of (1) General Amblyopia (see p. 430), or, in slighter cases, as amblyopia without central scotoma. Occasionally, higher degrees of amblyopia with concentric contraction of the field of vision, and yet negative ophthalmoscopic appearances, are present. (2) Atrophy of the optic nerve. This may appear in the usual form as progressive blindness, with concentric contraction of the field of vision; or it may come on after the slighter form of amblyopia has existed for some time. (3) Hemianopsia and color-blindness.

It is probable that these apparently different kinds of blindness depend upon similar pathological processes, and merely indicate degrees of the latter. In what these processes consist is still unknown; but the tendency to hemorrhages in the retina in diabetes makes it likely that hemorrhages in the optic nerve are sometimes the source of the amblyopia in question; while in the cases with central scotoma it is no doubt due to retrobulbar neuritis similar to that produced by tobacco, etc.

Amblyopia is sometimes the only symptom of diabetes; and, consequently, it is of the utmost importance to examine the urine for sugar in every case of amblyopia where the ophthalmoscopic appearances are negative, or where the only abnormality is atrophy of the optic papilla.

The Treatment indicated is solely that for the general disease, and the prognosis for vision depends upon the amenability of the latter to treatment, and upon the extent to which organic changes in the optic nerve have proceeded.

References.

[1] "Trans. Internat. Med. Congress," 1881, vol. iii. p. 52.
[2] "Von Graefe's Archiv," xvii. 2. p. 249.
[3] "Trans. Ophthal. Soc., U. K.", vol. v. p. 149.
[4] S. Snell, "Brit. Med. Journ.," March 3, 1894.
[5] P. Smith, "Ophthal. Rev.," 1893, p. 101; and Valude, "Revue d' Ophtal.," 1893, p. 231.
[6] "The Cerebro-Spinal Fluid, its Spontaneous Escape from the Nose," London, 1899.
[7] "Australian Medical Gazette," October 21, 1901, and "Archives of Ophthalmology," xxxi. 2. p. 114.
[8] "Klin. Monatsbl. f. Augenhlk.," vi. p. 425.

CHAPTER XVII.

THE ORBITAL MUSCLES AND THEIR DERANGEMENTS.

Normal Action of the Orbital Muscles.

The eyeball moves round a point on its antero-posterior axis, situated (in the emmetropic eye) 14 mm. behind the cornea, and 10 mm. in front of the posterior surface of the sclerotic. Its motions are effected by means of the six orbital muscles, arranged in three pairs, each pair consisting of two antagonistic muscles; thus the rectus internus and rectus externus are antagonistic, the former rotating the eye inwards, and the latter rotating it outwards. The remaining pairs are the recti superior and inferior, and the obliqui superior and inferior.

The Primary Position of the Eyeball is that one in which, the head being held erect, the gaze is directed straight forwards in the horizontal plane. This is the starting-point from which the actions of the muscles are considered.

The Rectus Externus and Rectus Internus, lying from their origin to their insertion in a plane which corresponds with that of the horizontal plane of the eyeball, move the latter on its perpendicular axis directly inwards and outwards, and have no other action.

The plane of the *Rectus Superior and Rectus Inferior* does not quite correspond with the vertical plane of the eyeball, and consequently the axis on which they rotate the globe is not its horizontal axis, but one which, passing from within and before, backwards and outwards, forms with the antero-posterior axis an angle of 70° (Fig. 131). While, then, their action is mainly to rotate the eyeball upwards and downwards, these muscles rotate it also somewhat inwards. Moreover, the superior rectus giving to the vertical meridian of the cornea an inward inclination,* or inward wheel-motion of the eye (*vide infra*), while the inferior rectus

* In speaking of the inclination of the vertical meridian of the cornea it is the upper extremity of this meridian which is meant.

gives this meridian an outward inclination or outward wheel-motion of the eye, the power of these muscles over the upward and downward motions is greatest when the eye is turned out, for then their axis of rotation coincides most closely with the horizontal axis of the globe; and their influence over the wheel-motion is greatest when the eye is turned in, for then their axis of rotation coincides most closely with the antero-posterior axis of the globe.

The plane of the *Oblique Muscles* of the eyeball also approaches the vertical plane of the eyeball, the axis upon which they rotate the latter passing from within and behind, forwards and outwards,




Fig. 131.




Fig. 132.

and making with the antero-posterior axis an angle of 35° (Fig. 132). The principal action, accordingly, of the oblique muscles is to incline the vertical meridian of the cornea; the sup. oblique inclines it inwards (wheel-motion inwards), the inf. oblique inclines it outwards (wheel-motion outwards). In addition to this action the oblique muscles, respectively, rotate the eyeball downwards and outwards (sup. oblique), and upwards and outwards (inf. oblique). It is evident that the power of the oblique muscles over the upward and downward motions of the eyeball is greatest if the eye be turned in, and that their power over the wheel-motion is greatest when the eye is turned out.

To sum up, then, the superior oblique and superior rectus produce wheel-motion inwards, while the inferior oblique and inferior rectus produce wheel-motion outwards. The action of the obliques on the wheel-motion is greatest when the eye is rotated outwards, and of the recti when the eye is rotated inwards.

In considering the motions of the eyeballs we have to think of the motions of one eyeball as associated with those of its fellow—




Fig. 133.

e. g., the action of the internal rectus of the left eye is associated with the action of the external rectus of the right eye, in rotation of both eyeballs to the right.

The vertical meridian of the eyes becomes inclined to the right or left in different positions of the globes, as was experimentally proved by Donders.

1. In the primary position, as also when the eyes are turned directly to the right, to the left, upwards, or downwards, the ver-




Fig. 134.

tical meridians (*a, b,* Figs. 133-137) maintain their vertical direction (Fig. 133).

2. When the eyes are turned to the *left, and upwards,* the vertical meridian of each eye is inclined at the same angle to the left (Fig. 134). Wheel-motion to the left.

3. When the eyes are turned to the *left, and downwards,* the vertical meridian of each eye is inclined to the right at the same angle (Fig. 135). Wheel-motion to the right.

4. When the eyes are turned to the *right, and upwards,* the vertical meridian of each eye is inclined at the same angle to the right (Fig. 136). Wheel-motion to the right.

5. When the eyes are turned to the *right, and downwards,* the vertical meridian of each eye is inclined at the same angle to the left (Fig. 137). Wheel-motion to the left.

We shall now consider what muscles of one eye are called into




Fig. 135.

action, when an individual requires to place it in the several principal positions.

1. *In the Primary Position* all the muscles are at rest.

2. Motion of the eyeball *directly outwards* is effected by the external rectus alone, and motion *directly inwards* by the internal rectus alone.

3. Motion of the eyeball *directly upwards* and *directly downwards* is effected chiefly by aid of the sup. and inf. recti. But




Fig. 136.

these muscles, acting alone, rotate the eyeball slightly inwards, and give a certain inclination to the vertical meridian, which in this position should be upright. Consequently, in rotation of the globe directly upwards, the inf. oblique, which rotates the eye slightly outwards (as well as upwards) and inclines the vertical meridian outwards, must be associated with the sup. rectus, in order to counteract in these particulars the tendency of its action. In rotation of the eyeball directly downwards, the inf. rectus must

be associated with the sup. oblique, which acts antagonistically to this rectus in respect of rotation inwards and of outward wheel-motion.

4. Rotation *upwards and outwards* is chiefly effected by aid of the rectus superior and rectus externus. But the latter muscle has no influence over the wheel-motion, while the former produces wheel-motion inwards. Yet the inclination of the vertical meridian is outwards in this position; and therefore a third muscle, which will supply this inclination in a high degree, is required—namely, the inferior oblique, whose power over the wheel-motion of the eyeball is greatest when the latter is turned upwards and outwards.

5. Rotation *downwards and outwards* is chiefly effected by the rectus inf. and rectus ext. Inasmuch, however, as the former inclines the vertical meridian outwards, while the latter has no influence over it at all, a third force is required which will bring




FIG. 137.

about the required inward wheel-motion—namely, the sup. oblique, whose influence in this respect is most powerful when the eye is turned downwards and outwards.

6. Rotation *upwards and inwards* is chiefly brought about by the rectus superior and rectus internus. But the effect of the former upon the inward wheel-motion of the eye would be so great as to interfere with parallelism of the vertical meridians of the two eyes, that of the other eye not being inclined outwards in a corresponding degree. A third force, therefore, is required, which will to a certain extent counteract the influence of the sup. rectus in this respect, and this is the inf. oblique, which in this position of the eyeball has but slight power over its wheel-motion.

7. Rotation *downwards and inwards* is chiefly the result of contraction of the rectus inf. and rectus int. The power of the former over the outward inclination of the vertical meridian would, in a similar way, be too great, and must be similarly corrected by the action of the superior oblique.

The Field of Fixation.

The field of fixation contains all the points which the eye can successively see or "fix" without movement of the head. It can be measured with the perimeter, as in testing the field of vision, except that here the patient is made to move the eye as far as possible in each meridian, and the limit of each movement is measured by observing the corneal reflex of a candle flame, or ophthalmoscope mirror, which is moved along the arc of the perimeter. The *binocular* field of fixation contains all those points which can be seen as single with the two eyes and without movement of the head. According to Landolt [1] the averages give, for movement of one eye, inwards 44°, outwards 46°, upwards 44°, and downwards 50°.

Strabismus.

When looking at any object with the two eyes it is necessary, in order to avoid seeing double, that the visual axis of the eyes should meet at the point fixed. When this does not take place, one of the eyes must be in a faulty position, or, as it is commonly termed, it squints. This condition is called *strabismus,* and may arise either from overaction or from paralysis of one of the muscles. Strabismus may occur in any direction, but vertical and oblique deviations are less common than the convergent or divergent forms.

In order to ascertain in slight cases which of the two is the deviating eye, the patient is made to fix a certain object, and one of the eyes, say the left, is rapidly covered with the surgeon's hand; then, if the right eye make no movement, it must have been looking at the object before the left one was covered; but if, on covering the right eye, the left make a movement in order to fix the object, then this eye must be the squinting one. The movement is always in the opposite direction to the deviation. For instance, if the eye be turned inwards too much, it must of course turn outwards to fix the object, when its fellow is covered. Another good method consists in observing the position of the corneal reflex when the patient looks at the ophthalmoscope (see Measurement of Strabismus). But the most delicate test is the character of the diplopia, when diplopia is present.

Apparent Strabismus is due to a large angle γ (p. 24). In this

case, as the visual axes are both directed to the point fixed, there will be no movement of either eye on covering the other, as in true strabismus.

Paralyses of the Orbital Muscles.

Loss of power of one or more of the muscles of the eyeball is, of course, always to be regarded as a symptom, not as in itself a disease.

It may be due to lesions in several different situations, namely: (1) Lesions situated in the orbit. (2) Basic lesions—lesions situated at the sphenoidal fissure, and those at the base of the skull, between the sphenoidal fissure and the pons. (3) Pontine lesions, which may be Fascicular—*i. e.*, involving the ocular nerve fibers in the substance of the pons, or Nuclear—*i. e.*, only attacking the nuclei of the nerves in the aqueduct of Sylvius and floor of the fourth ventricle. (4) Cerebral lesions—lesions above the nuclei, in the internal capsule, corona radiata, or cortex. These four classes differ considerably in their clinical aspect, in their pathological causes, and in their significance for the well-being of the patient.

The first class—loss of power due to orbital lesions—will be considered in the chapters on Diseases of the Orbit.

The second class—those due to basic lesions—provides by far the largest number of cases of paralyses of the orbital muscles. Let us now consider the

General Symptoms of this second class. They include symptoms to be found in each of the other classes. (1) Diplopia. The affected eye being deviated from its correct position, and being more or less incapable of associated motions with the other eye, the image of the object looked at is not formed on identical spots of the retina in each eye, and hence the object seems doubled. (2) Indistinct vision. If the paralysis be but slight, actual diplopia may not be present, but the double images overlapping each other will cause dimness or confusion of sight. (3) Giddiness, due partly to the diplopia, and partly to faulty projection of the object. By faulty projection is meant the false idea of the position of the image in the field of fixation. (4) Some patients turn the head towards the side of the paralyzed muscle, in order to diminish or eliminate the diplopia—*e. g.*, if the left ext. rectus were paralyzed, the head would be turned towards the left; if it

were the left int. rectus, the head would be turned towards the right. By this maneuver the loss of the action of the affected muscle is less felt for those objects which lie straight in the patient's path, while he walks about; because it involves a rotation of the eye towards the side of the healthy antagonist, in which region of the binocular fixation field the diplopia is reduced to a minimum. Some patients close one eye to procure single vision. (5) In peripheral paralysis it is most common to find only the muscle, or muscles, supplied by some one nerve—the third, fourth,




FIG. 138.

or sixth—affected; although, of course, exceptions to this are not rare, especially where a neoplasm forms at the base of the skull.

In studying a case of paralysis of an orbital muscle the following *General Principles* should be borne in mind: (1) The defective mobility and the diplopia increase towards the side of the affected muscle—towards the left, if the left external rectus be paralyzed; towards the right, if the left internal rectus be paralyzed. (2) The secondary deviation (*i. e.*, the deviation of the sound eye while the affected eye fixes) is greater than the primary deviation (*i. e.*, the deviation of the affected eye while the sound eye fixes); because the muscle in the sound eye, which is associated in its action with the paralyzed muscle in the affected eye (*e. g.*, the rect. int. with the rect. ext.), must receive a nervous im-

pulse of equal intensity to that sent to the weak muscle, and, as the latter requires a considerable impulse to excite its action, its associate will be overexcited. Let us suppose the left external rectus to be paralyzed, and that, shading the right eye with a hand, we direct the patient to fix with his left eye an object held somewhat to his left-hand side; we may notice, on removing the shading hand, that the right eye has been rotated inwards to an extent far exceeding that of the primary deviation of the left eye, and has now to make an outward motion in order again to fix the object. (3) The image formed on the retina of the affected eye is projected (*i. e.*, seems to the patient to lie) in the direction of the paralyzed muscle—*e. g.*, if the left ext. rect. be paralyzed, the image corresponding to that eye will be projected to the left of the image belonging to the right eye.

Where the image of the affected eye lies to the corresponding side the diplopia is termed homonymous, and homonymous double vision always indicates convergence of the visual lines. Fig. 138 explains the occurrence of homonymous diplopia in convergent paralytic strabismus.* The right eye fixes the object *o*, and its image on the macula lutea *m;* but the left eye, by reason of the paralysis of the external rectus, is turned in, and its visual axis lies in the direction *m v*, and the image of *o* falls to the inner side of the macula lutea at *a*. Now why should this image not be referred to its correct position along the line *a o?* The reason is that the patient is not conscious of the deviation of this eye; and, having always been in the habit of superposing his fields of vision, so that the visual axes of the eyes meet at the object fixed, he imagines that this is still the case, and that *v m* lies in the position of *o a*, and that the macula lutea *m* is at *m'*. But if this were the case, *a* would be at *a'*, and in this position of the eye, indicated by the dotted line, images formed at *a'* to the inner side of the macula lutea are projected to the outer side of the field, along the line *a' o'*, and the patient imagines that *o* occupies the position *o'*, as seen with the left eye.

If we suppose the internal rectus of the left eye to be paralyzed, the image on the retina of that eye falls then to the outside of its macula lutea, and must therefore be projected to the right of the true position of the object; this is crossed diplopia, and attends divergence of the visual lines.

* For the sake of simplicity in the diagram the effect which rotation of the eye has on the nodal point is omitted.

Paralysis of the External Rectus of the Left Eye.—If this be complete or considerable, it is easy of diagnosis, as marked loss of power and motion of the left eyeball outwards is present, and the patient complains of double vision. He keeps his head turned to the left, in order to diminish the influence of the paralyzed muscle as much as possible.

If, however, the paralysis be but slight, the patient may not complain decidedly of diplopia, but only of indistinctness or confusion of sight, especially when he looks towards the left. To decide the diagnosis in such a case, the double images must be examined. A long lighted candle is used as the object to be looked at; and one eye—let us say here the left eye—is covered with a bit of red-stained glass in order to differentiate the images.* The candle is now held on a level with the patient's eyes, and straight opposite him, at about three meters' distance (eyes in primary




FIG. 139.

position). (*a*) In this position the images are seen very close together or overlapping each other, both of them upright and on the same level, the red candle to the left, the white to the right—*i. e.*, homonymous diplopia = convergence. This convergence must be due to paralysis of one or other external rectus muscle, but we cannot say at this stage of the experiment which of them is affected. (*b*) In order to determine this point the candle must be carried from side to side, and the increasing or decreasing distance of the images from each other noted. If the candle be carried slowly to the right, the patient following it with his eyes without turning his head, the images come still closer together, or only one candle is seen. But if the candle be carried to the patient's left-hand side, the images go farther apart, their relative positions being maintained. We now know that it is the left

* Maddox' Rod Test, described further on, is very suitable here, and in the investigation of other forms of ocular palsy.

external rectus which is affected: because towards the left—the direction in which the action of this muscle is most wanted, and consequently its loss most felt—the distance between the double images increases. The images are erect, as no wheel-motion is caused by action of the external rectus. (*c*) If, however, the candle be held to the left and raised aloft, the image belonging to the left eye will seem to lean away from that of the right eye (Fig. 139). The reason of this is that, owing to the paralysis of the external rectus, the left eye cannot look upwards and outwards as it should, but merely looks upwards. The vertical meridian therefore remains vertical. But the right eye, which is free to follow the candle, looks up and to the left. Its vertical meridian is therefore inclined to the left. That is, the vertical meridians of the two eyes converge at the top, which necessitates a divergence of the upper extremities of the images. The rotation of the right eye in this position is physiological, and its image is therefore judged to be vertical; while the image of the left eye diverging from that of the right, though really vertical, is judged to be oblique. An analogous derangement of the vertical meridian takes place in the position below and to the outside. (*d*) If the patient be told to direct his gaze specially towards the red candle—*i. e.*, the image which belongs to the left, the affected, eye—the distance between the two candles will be much greater than if he direct his gaze towards the white candle. This is explained by General Principle No. 2 (p. 448).

If the patient's good eye be closed, and an object (surgeon's finger) be held up within his reach, but towards his left-hand side, and he be requested to aim rapidly at it with his forefinger, he will aim to the left of it. The nervous impulse sent to his left external rectus, to enable him to turn the eye towards the object, is of such intensity as to lead him to fancy that the object lies much farther to the left than is the case (incorrect projection of the field of view); for we, to a great extent, estimate the distance of objects from each other by the amount of nervous impulse supplied to our orbital muscles in motions of the eyeball.

A prism held horizontally before the affected eye with its base outwards brings the double images closer together; or, if the correct prism be selected, the images will be blended into one.

Paralysis of the Superior Oblique of the Left Eye.—This paralysis will be most apparent when a demand is made for motion of the eyeball downwards and inwards, motion in this direction

being that over which the superior oblique has most influence. Yet absolute defect of motion is sometimes difficult to detect even in complete paralysis of this muscle, owing to vicarious action of the inferior rectus and of the internal rectus. Careful examination of the secondary deviation will often be successful as to this point, but it is on the examination of the double images that we must chiefly rely for the diagnosis.

(*a*) In the whole of the field of vision above the horizontal plane there is single vision. Below the horizontal plane in the median line diplopia appears, the image belonging to the left eye standing lower than that belonging to the right, because the superior oblique being a muscle which assists in rotating the eye downwards, the latter, for want of the action of this muscle, now stands higher than its fellow (right eye), and consequently the image will not fall on its macula lutea (as it does in the right eye),




Fig. 140.

but above it, and will therefore be projected below the image of the right eye. The position downwards and inwards of the eyeballs is that in which the greatest demand is made upon the superior oblique for rotation of the eye downwards: therefore it is in this position its want for this purpose is most felt; and when the candle is held in this position the vertical distance between the double images is greatest. (*b*) The superior oblique assists also in rotation of the eye outwards: therefore loss of its power must commit the eyeball to a certain extent to the power of the muscles which move it inwards, and a rotation in this latter direction (convergence) takes place, with the result of making the image belonging to the left eye stand to the left of the image belonging to the right eye (homonymous diplopia). (*c*) The superior oblique inclines the vertical meridian inwards: therefore, in rotation directly downwards, loss of its power commits the eye to the outward wheel-motion imparted to it by the inferior rectus. This

gives to the image belonging to the left eye an inclination to the patient's right hand. (*d*) The power of the superior oblique to incline the vertical meridian inwards is greatest when the eye is turned downwards and outwards: consequently, in this respect its paralysis will be most felt in this position, and therefore here the inclination of its image to that of the sound eye will be most marked. (*e*) A remarkable phenomenon usually noticed in this paralysis (and sometimes in paralysis of the inferior rectus), and for which a good explanation does not exist, is that the image belonging to the affected eye seems to stand nearer the patient than that of the sound eye.

To sum up, then (*vide* Fig. 140): below the horizontal plane there is homonymous diplopia, while the image (*A*) of the affected eye stands on a lower level, is inclined towards the other image, and seems to be nearer the patient. Furthermore:

(*f*) In an extreme lower and outer position the image of the affected eye may sometimes seem to stand higher than that of the



Fig. 141.



Fig. 142.

Fig. 141.—Paralysis of left sup. oblique. Homonymous diplopia. R, image of right eye. L, image of left eye.

Fig. 142.—Paralysis of right inferior rectus. Crossed diplopia. R′, image of right eye. L′, image of left eye.

sound eye, owing to an excessive outward inclination of the vertical meridian, which throws the image on the lower and outer quadrant of the retina.

In order to do away with or to diminish the diplopia, the patient inclines his head forwards, and turns it to the side of the good eye.

For the prismatic correction of the diplopia, two prisms will be required; one with its base downwards in front of the left eye, to correct the vertical difference, and a second with its base outwards in front of the right eye, to correct the lateral difference.

To make the diagnosis between the foregoing paralysis and paralysis of the right inf. rectus (in which the diplopia is also be-

low the horizontal plane only, and the false image lower than the true one and inclined towards it) it has merely to be remembered that there is here crossed—instead of homonymous—diplopia, because the superior oblique, which now chiefly effects the downward motion of the eyeball, turns it at the same time somewhat outwards. The Figures 141 and 142 will assist in this explanation.

Paralysis of the Internal Rectus, Superior Rectus, Inferior Rectus, Inferior Oblique, and Levator Palpebræ.—Complete paralysis of all the branches of the third nerve produces a remarkable appearance. The upper lid droops (ptosis), the pupil is semi-dilated and immovable, the power of accommodation is de-




Fig. 143.

stroyed, and the eyeball is often slightly protruded, owing to the backward traction of the recti being lost to it. Motion inwards exists but to a slight degree, and motion downwards is effected only by aid of the superior oblique, and is accompanied by marked inward wheel-motion. If the paralysis be of some little standing, the external rectus obtains rule over the eyeball, and rotates it permanently outwards.

The diagnosis, then, in cases of complete paralysis of all branches of the nerve is easily made; but not so, sometimes, if the paralysis be only slight. The examination of the double images, then, is of value. If (see Fig. 143) the left third nerve be partially paralyzed in all or most of its branches, there will be crossed diplopia either in the whole of the field of vision—for want of

power in the internal rectus—or towards the patient's right at the least, and the lateral distance between the images will increase as the visual object is carried farther towards the right. When the visual object is held aloft the left eye will remain behind—for want of the action of both of the muscles which turn the eye upwards—and, consequently, in this position its image will stand, not only to the right of, but also above that of the right eye; while, when the visual object is held below the horizontal plane, the eye will—owing to paralysis of the inferior rectus—remain higher than the right eye, and consequently its image will appear to be lower than that of the right eye. It will, moreover, be inclined




Fig. 144.




Fig. 145.

towards the latter image, in consequence of the inward wheel-motion imparted to the eye by the healthy superior oblique.

When in each eye some branches of the third nerve are paralyzed the diagnosis is often extremely complicated. The ptosis, however, which is nearly always present, and is readily recognized, and the paralysis of the sphincter iridis (mydriasis) and of accommodation, which often exist, and are also easily observed, give valuable aid. Moreover, any loss of motion upwards must be due to paralysis of the third nerve; but if there be loss of motion downwards, the differential diagnosis between paralysis of the inferior rectus and of the superior oblique has to be made. For this see the paragraph on paralysis of the latter muscle.

As may be imagined from the foregoing, it is often difficult in practice to keep clearly before one's mind the different actions

of the orbital muscles, and from the character of the diplopia to deduce the paralysis which may be present. An aid in this respect has been provided by Dr. Louis Werner [2] by means of two diagrams (Figs. 144 and 145).

The form of diplopia which characterizes paralysis of each muscle is expressed by the position of the dotted line bearing the name of the muscle. The dotted lines represent the false images, the continuous lines the true images.*

In the case of the recti (Fig. 144) the false images inclose a lozenge-shaped space situated between the true ones, whereas in the case of the oblique muscles (Fig. 145), the true images, which, for the sake of simplicity, are combined in one line, lie between the four false images, which diverge from one another so as to form an X. It will also be noted that the dotted lines extend upwards and downwards beyond the others, indicating respectively that the false images are higher or lower than the true ones. Another fact which the diagrams indicate is that, in the case of the muscles represented in the upper halves of the figures, the diplopia occurs in the *upper* part of the field of fixation, or, in other words, in upward movements of the eyes. A similar rule holds good with regard to the lower halves.

The method of using the diagrams will be better understood by taking a particular muscle as an example. Suppose, for instance, that we wish to know what kind of diplopia results from paralysis of the *left inferior* rectus, it is simply necessary to look at the *left inferior* portion of Fig. 144 (recti), which gives the diplopia. If we analyze this we find (1) that the diplopia is *crossed,* for the false image corresponding to the *left* eye is on the *right* of the true image—*i. e.,* the right image corresponds to the left eye; (2) that the false image has its *upper end inclined towards the true one;* (3) that the false image is *lower* than the true one, for the dotted line extends *lower* than the other one; (4) that the diplopia occurs in *downward movements* of the eyes, for it is in the *lower* half of the diagram that the false image lies.

The same method applies to the other recti: the diplopia for the *right upper* rectus is found in the *right upper* quadrant, and so on for the rest.

The same rules also apply to the obliques (Fig. 145), with one difference. The recti move the eye in the direction indicated by

* The "false image" corresponds to the affected eye, and the "true image" to the sound eye.

their names, the superior moving it upwards and the inferior downwards; but in the case of the obliques the reverse takes place, the superior oblique moving the eye downwards and the inferior upwards. Therefore, for the *superior* obliques we must look at the *lower* half of Fig. 145, and for the *inferior* obliques at the *upper* part.

This is an extremely simple method. By bearing the figures in mind it is possible to tell immediately what kind of diplopia would result from paralysis of any one of these muscles, and conversely, given the diplopia, to determine to which muscle it is due.

The Causes of Peripheral Paralyses of Orbital Muscles are chiefly of rheumatic or syphilitic nature.

Rheumatic paralysis, to which the external rectus is specially prone, will be noted if there are symptoms of general rheumatism, or if there is a history of exposure to cold or wet immediately preceding the attack.

Syphilis will be suggested as a cause if there be a specific history, and that other causes can be excluded. Peripheral paralyses of the orbital muscles due to syphilis are amongst the later symptoms of the disease, and may depend on exostoses or gummata at the base of the skull, or to syphilitic neoplasms, or meningitis, in the course of the nerve. The third nerve seems to be particularly liable to be attacked by a solitary gumma at the base of the skull, especially at the sphenoidal fissure, ptosis being commonly the first symptom.

Other neoplastic growths can, of course, cause these paralyses in the same way.

Prognosis.—In peripheral paralyses recovery is very frequent; much, however, depending on the nature of the lesion. In cases where a cure is not effected, the antagonist muscle often becomes contracted, and the eye is then rotated permanently and excessively in the corresponding direction. In cases of old standing, a permanent contraction of the muscles of the neck may be brought about, from the inclination of the head which the diplopia has obliged the patient to adopt.

Treatment.—In these cases the medical treatment consists in drugs suitable to the fundamental disease (rheumatism, syphilis, etc.). Local depletion at the temple by the artificial leech in the early stages, and galvanism later on, may be employed with advantage. The most common method of applying galvanism is

through the closed lid; but it is probable that the episcleral method—*i. e.*, with the electrode placed directly over the muscle—is more effectual. Dr. Buzzard's method [3] seems to be a very admirable one. He applies a moistened plate rheophore to the nape of the patient's neck, and connects it with one pole of a Leclanché battery. He then takes the other rheophore, well wetted, in his left hand, and, securing good contact with the skin of his palm, applies the index finger of his right hand to the patient's globe in the situation of the various external muscles of the eye. The finger is covered with a single thickness of well-moistened muslin, and the conjunctiva should be previously rendered insensitive by cocain. The strength of the current advised is from 1.5 to 2 milliampères, and the alternate application and lifting of the finger, by closing and opening the circuit, gives rise to a feeling of a slight electric shock in the terminal point of the finger. The operator should first test the strength of the current upon the patient's cheek. The point of the finger thus employed acts as a sentient rheophore, and can be applied with nicety and delicacy to various parts of the eye, the operator being constantly aware, by the feeling in his finger, of the strength of the current employed.

Passive orthoptic treatment [4] occasionally gives a rapid and brilliant result, while, again, it is useless. It is performed as follows: The conjunctiva at the corneo-scleral margin, near the insertion of the paralyzed muscle, is seized with a forceps, and the eyeball is drawn in the direction of the muscle, and as far as possible beyond its ordinary limit of contraction, and back again. These movements are continued for about a minute once a day, cocain being used.

Prismatic glasses may be used, either to eliminate the diplopia or to excite the weak muscle to exert itself. In the former case the glass selected must completely neutralize the diplopia; but as it can do so only for one position of the eyes, prisms are rarely employed in this way. In the latter case a prism slightly weaker than that sufficient to completely neutralize the diplopia is selected, in order that, with a little effort, the weak muscle may be enabled to bring about single vision, and this effort having been successfully maintained for some days, a still weaker prism is then prescribed, and so on.

It is very important for the patient's comfort while awaiting his cure, unless a cure by prisms as above described is being at-

tempted, that the affected eye should be covered, so that the distressing double vision may be obviated.

Surgical treatment is justifiable only when other means have failed to restore muscular equilibrium. If the deviation amount to 3 or 4 mm., tenotomy of the antagonistic muscle, with subsequent tenotomy of the associate muscle in the other eye, will be sufficient; but if the deviation amount to 5 or 6 mm., advancement of the paralyzed muscle, in addition to the tenotomy, may be required. This surgical treatment applied to the internal and external rectus sometimes gives satisfactory results; but in the cases of the superior and inferior recti it is less useful, while the oblique muscles should not be operated on.

A peculiar and rare form of peripheral or basal paralysis is **Intermittent Paralysis of the Third Nerve of one Eye,** for which Charcot suggested the name of **Ophthalmoplegic Migraine.** The patients are generally children or young adults, who usually suffer from headache on the side corresponding to the paralyzed eye, and sometimes from vomiting. The paralysis may be complete or partial, and the attack varies in its duration from a few days to a few months. Some cases are purely periodical —*i. e.,* in the intervals between the attacks of paralysis all the muscles supplied by the third nerve act in a completely normal manner; while in other cases these muscles, or some of them, do not completely recover their functions in the intervals. We are as yet quite in the dark as to the cause of these periodical paralyses of the third nerve. Some hold that the purely periodical cases are of a functional nature, possibly hysterical or reflex, and that the periodically exacerbating cases alone are due to a lesion of the root of the nerve, of an undefined kind, at the base of the skull; while others are of opinion that both forms depend upon a diseased process at the base. In three cases in which an autopsy was made, there was disease of the trunk of the nerve at the base of the skull.

In intermitting paralysis the *Prognosis* of the purely periodical form is favorable, inasmuch as the attacks in the course of time become fewer and less severe, until, finally, they cease entirely. In the exacerbating form the prognosis for complete recovery is less favorable. Out of twenty-six cases collected by Darquier only one patient died from a cerebral cause.

In view of the obscurity which still surrounds the causation of these intermitting paralyses their *Treatment* must consist, in each

case, in the relief of any general dyscrasia or concomitant symptoms which may be present.

The third class of paralyses of orbital muscles above enumerated—those due to lesions of the nuclei of the orbital muscles in the aqueduct of Sylvius and floor of the fourth ventricle—are known by the term

Ophthalmoplegia Externa, and also as **Nuclear Paralysis.**—The first of these terms was originally employed to denote those remarkable cases in which, all, or nearly all, of the orbital muscles of both eyes are paralyzed, while the intra-ocular muscles often remain intact. There can be no doubt, however, that these cases do not differ in their nature from many of those in which, in one eye, several orbital muscles supplied by different nerves—*e. g.*, third and fourth—are wholly or partially paralyzed; or where all the orbital muscles in one eye are wholly or partially paralyzed; or where in each eye muscles supplied by the same nerve—*e. g.*, both sixth nerves—are wholly or partially paralyzed; for such cases are often mild forms of the disease, or else stages in its development. At one time it was considered essential for the diagnosis that the intra-ocular muscles should retain their functions, but cases occur in which the sphincter iridis and ciliary muscle are paralyzed.

When these two latter muscles alone are paralyzed, the condition is called Ophthalmoplegia Interna. When both they and groups of orbital muscles are paralyzed, the terms Ophthalmoplegia Interna et Externa, or Ophthalmoplegia Universa, are employed.

The term Nuclear Paralysis indicates any orbital paralysis due to a lesion of the nuclei of the orbital nerves in the pons, and ophthalmoplegia externa comes within this category.

Ptosis, even in cases of complete binocular ophthalmoplegia externa, is often incomplete, and it is remarkable that in some chronic cases, without any improvement in the condition itself, the diplopia, which was at first present, quite disappears.

Occurrence and Progress.—The condition may be congenital, or may make its appearance soon after birth, and may remain permanently without becoming complicated with any further disturbance. Congenital ptosis, which is frequently combined with loss of power in the superior rectus, and is usually binocular, is of this nature. But Nuclear Paralysis is more commonly seen as an acquired condition in childhood, or in adult life, either in an acute

or chronic form. Marked cerebral lethargy is often seen with both forms, and the tendon reflexes may be defective.

Acute Nuclear Paralysis is due either to an acute inflammatory process in the nuclei—comparable to the process which produces poliomyelitis anterior acuta, and hence it is called by Byrom Bramwell poliomyelitis acuta—or to hemorrhagic lesions.

The acute inflammatory cases are apt to have a sudden onset, attended with fever, headache, vomiting, and convulsions, which may subside after a few days, leaving only the ophthalmoplegia behind; and this, too, after a lengthened period, may undergo cure, partial or complete. Sometimes these attacks are complicated with paralysis of the facial nerve, or the diseased process may extend to the spinal cord, and the symptoms of acute poliomyelitis become developed; or, again, acute bulbar paralysis may come on.

Acute peripheral neuritis of the ocular nerves, which is sometimes seen in cases of alcoholic poisoning, may be confounded with acute nuclear palsy. The symptoms of the two states are the same, except that in the case of peripheral neuritis there are no head symptoms at the commencement.

The onset of acute hemorrhagic ophthalmoplegia is sudden, but is unattended by headache, vomiting, or convulsions. It takes different courses. Sometimes it is rapidly fatal; again, it goes on to softening of the nuclei, and becomes chronic; while, again, it undergoes a slow cure.

It is extremely probable that to this hemorrhagic class the paralyses of orbital muscles belong, which sometimes follow on an attack of diphtheritic sore-throat. These paralyses appear in from one to six weeks after the outbreak of the primary affection. The latter need not have been of a severe kind; indeed, sometimes patients are unaware that they have had a sore throat. These diphtheritic paralyses always recover in the course of some weeks.

In diabetes, paralyses of orbital muscles are not very uncommon, and are probably to be classed as nuclear. The same may be said of orbital paralyses in lead-poisoning and in influenza. Other causes are cold, poisoning by nicotin, sulphuric acid, carbonic oxid, and tainted meat.

The Prognosis in all these instances is favorable.

Chronic Nuclear Paralysis (Chronic Poliencephalitis Superior, of Wernicke) is much more common than the acute form. It depends on a degenerative atrophy of the nerve nuclei, analogous to

that which occurs in progressive muscular atrophy and in chronic bulbar paralysis. The onset is gradual, the loss of power in the muscles being at first very slight, but ultimately complete paralysis of the affected muscles results. There is no fever, nor any cerebral symptom. The condition may become associated with chronic bulbar paralysis, with progressive muscular atrophy, or with locomotor ataxy. But this is not so liable to occur in infants as in adults.

In some cases there may be partial paralysis of the orbicularis palpebrarum, which, according to Mendel, is innervated from the third-nerve nucleus through the facial nerve, along with other muscles of the oculofacial group (frontalis and corrrugator supercilii).

Coarse lesions, especially tumors of the pons and of its neighborhood which press on it, may produce orbital paralyses closely simulating those due to nuclear lesions. But here the paralysis is only one of the symptoms in the case, which are likely to include headache, vomiting, optic neuritis, hemianopsia, hemiplegia, etc. Softenings, patches of disseminated sclerosis, and internal hydrocephalus with overdistention of the aqueduct of Sylvius, are other lesions which may give rise to similar orbital paralyses, but which cannot be regarded as true nuclear ophthalmoplegia. The mode of onset, and the concomitant symptoms, of each case must serve as our guides in arriving at a diagnosis, which will sometimes be difficult enough.

Conjugate Lateral Paralysis of the eyes is a symptom which may be caused by a lesion in the pons. We believe that the voluntary motor impulses, coming down from the cortex to produce associated lateral motions of the eyeballs—*i. e.*, action of the external rectus of one eye, along with action of the internal rectus of the other eye—first reach the nucleus of the sixth nerve, and then pass on, through fibers called the posterior longitudinal bands, under the corpora quadrigemina, and join with the fibers of the opposite third pair for the supply of the internal rectus of that side. The sixth pair of one side supplies in this way the external rectus of its own side, and to a slight extent the internal rectus of the opposite side; and it is quite probable that similar decussations may exist in the nerve supply of other orbital muscles. Hence a lesion at, let us say, the left sixth nerve nucleus would paralyze the conjugate lateral motions of the eyes towards the left side; and there would in consequence be conjugate lateral devia-

tion of the eyes towards the right—the eyes looking away from the lesion. In conjugate paralysis, or deviation, whether due to a pontine lesion, or, as in a later paragraph, to a cerebral lesion, the combined action of the internal recti for the purpose of convergence of the eyes is retained.

Paralysis of the orbital muscles from nuclear disease may occur in Locomotor Ataxy, disseminated Sclerosis, General Paralysis, and more rarely in Exophthalmic Goiter and Severe Multiple Neuritis.

Fascicular Paralyses are mainly distinguished by the presence of other symptoms due to involvement of neighboring structures. They are rarely symmetrical. Vertigo is common with fascicular third-nerve paralysis, owing to implication of the red nucleus in the tegmentum which is connected with the superior peduncle of the cerebellum.

In *Myasthenia Gravis* the symptoms include some which are due to derangement in the power of orbital muscles. A very complete account of the disease was published by Buzzard,[3a] and Harry Campbell and Edwin Bramwell [3b] have also published a paper on the subject. As regards the eye-symptoms, these latter authors state: Ptosis is a common symptom; it is usually bilateral and more marked on one side than on the other. It may be constant, or it may only be present towards the latter end of the day, or if the patient looks up for any length of time, when the lids gradually fall. Owing to weakness of the occipito-frontalis, their compensatory overaction, so common in other forms of ptosis, cannot occur. Weakness of the orbital muscles with resulting diplopia is often present. Sometimes one muscle is more affected than others, sometimes there is a general paresis affecting all the orbital muscles, while in some cases complete and persistent ophthalmoplegia externa has been present. An alteration in the relative position of the two images is a striking feature. In some cases irregular nystagmoid movements are induced upon conjugate lateral motion of the eyes. The ocular muscles, as is the case with other voluntary muscles, become readily fatigued, the patients complaining that after reading a few lines the words and letters run into each other. Pupil changes are exceptional, but they are sometimes unequal. In Buzzard's case, after prolonged convergence, the pupils showed a tendency to oscillatory movements. The power of accommodation does not become fatigued.

The general features of the disease are: Weakness of some or

all of the voluntary muscles of the body, which may amount to complete paralysis. After a long rest—*e. g.*, the first thing in the morning—they may respond normally to the will, but become rapidly exhausted after a little use. The affected muscles often exhibit the myasthenic reaction, becoming exhausted by faradic stimulation. The entire system of voluntary muscles may be affected, but those muscles are most apt to be implicated which normally act most constantly, such as the cervical muscles and the extrinsic muscles of the eyeball. The symptoms fluctuate from day to day, or from month to month, and may even disappear for months or even years, and reappear. There are no sensory symptoms. Death occurs in a large proportion of the cases, but no structural changes have been found to account for the symptoms.

Cerebral Paralysis of Orbital Muscles form the fourth and last of the classes enumerated. They include all the orbital paralyses due to lesions above the nuclei—*i. e.*, in the cortex, corona radiata, or internal capsule. They are usually associated with other symptoms, which aid us in localizing, more or less accurately, the lesions which cause them. These paralyses are always physiological, associated, or conjugate, as they are variously and with equal correctness termed—they are, in short, paralyses of motion rather than of muscles.

Conjugate lateral paralysis—loss of power of motion of the eyes to one side or to the other, while the power of convergence of the optic axes is retained—is by far the most common form of this symptom. We do not as yet know where the cortical center for the associated lateral motions of the eyes is situated.* But even if we did know its position, it is not likely that much would be gained so far as clinical localization of the cerebral lesion is concerned; for this center, wherever it may be, is extremely sensitive, and is apt to be thrown out of gear by lesions of many different parts of the cortex. Conjugate deviation is, in short, very apt to be a distant symptom, especially in cerebral hemorrhage,

* This center has been placed by various authors in the inferior parietal lobule (Wernicke, Henschen, Munk, etc.), and in the second frontal convolution (Ferrier, Horsley, and Beevor). Stimulation of the centers of vision in the occipital lobe has also been found to produce conjugate movements (Schaefer, Munk), and these have been regarded as reflex by some; but Knies holds that the visual center contains the motor center for the eye-muscles as well. Moreover, it is stated that the visual cortex contains motor pyramidal cells.

when it is often accompanied by a rotation of the head in the same direction, and lasts only a short time. Moreover, it is thought that, when this center may happen to be actually involved in the lesion, its function, being largely bilateral, is rapidly taken up by the opposite hemisphere; and hence, even when conjugate lateral deviation plays the part of a direct cortical symptom, it rarely can be recognized as such, owing to its evanescent character. In paralyzing lesions the deviation of the eyes is of course towards the side of the lesion—the eyes look at the cerebral lesion, as Prevost has expressed it—while in irritating lesions the spasm of the




FIG. 146.

1. Left Ext. Rectus; 2. Left Int. Rectus; 3. Right Int. Rectus; 4. Right Ext. Rectus; 5. Nucleus left third nerve; 6. Nucleus right third nerve; 7 and 8 Post, longitudinal bands from sixth nerve to opposite third nerve; 9. Nucleus left sixth nerve; 10. Nucleus right sixth nerve; 11 and 12. Left and right cortical centers. An impulse starting from 12 would travel down to 9, and produce an associated movement of the eyes to the left.

affected muscles causes the deviation to be from the side of the lesion. These conditions are the reverse of what happens in conjugate lateral deviation due to lesions in the pons (p. 462), and we are thus enabled to differentiate between lesions in the two positions.

There are four possible cases:

Cerebral Lesions	Destructive.	Eyes turned away from paralyzed side.
	Irritative.	" " towards convulsed side.
Pontine Lesions	Destructive.	" " towards paralytic side.
	Irritative.	" " away from convulsive side.

The cerebral cases show that the center for associated movements is on the opposite side of the brain—*e. g.*, in movements of eyes to the left, the left external rectus and right internal rectus are innervated by the right hemisphere of the brain; consequently, a destructive lesion here would produce paralysis of the left side of the body and of the associated movements of the above orbital muscles, and therefore the eyes would be drawn to the right by their opponents—*i. e.*, away from the paralyzed side. A destructive lesion of the right side of the pons would also, of course, produce paralysis of the left side of the body; but, involving the right sixth nucleus, it would cause paralysis of the associated movements of the right external rectus and left internal rectus, and, consequently, the eyes would be drawn to the left by the opponents—*i. e.*, towards the paralyzed side.

The reverse of the foregoing would occur in irritative lesions. Fig. 146 serves to illustrate the points referred to.

A destructive lesion at 12, the right cortical center, involving also motor centers of the body, would cause left hemiplegia; and since the external rectus of the left eye and internal rectus of the right eye would be paralyzed, the antagonists would turn the eyes to the right—*i. e.*, away from the paralyzed side. A destructive lesion of the right side of the pons, also producing left hemiplegia, if it involve the sixth nucleus, will produce paralysis of the external rectus of the right eye and of the internal rectus of the left eye, and the antagonists would turn the eyes to the left—*i. e.*, towards the paralyzed side. It is easy to see how irritative lesions would produce exactly the opposite effects.

Hemianopsia interferes to a certain extent with the conjugate movement towards the affected side, in so far as this is guided by visual impressions. According to Knies, the difficulty in reading the right hemianopsia is mainly due to this cause.

It seems important here, even at the risk of some repetition, to direct special attention to

The Localizing Value of Paralyses of Orbital Muscles in Cerebral Disease.—*Paralysis of the Third Nerve.*—As regards this nerve we are struck with the fact that ptosis, partial or complete, may be present as a focal symptom in cortical lesions—cerebral ptosis, as it is called—without any other third-nerve branch being paralyzed. That a separate cortical center for this branch of the third nerve exists, and that it innervates the muscle of the opposite side, is very probable. The existence of such a

center would not be inconsistent with the view that, as regards the motions of the eyeballs, associated centers alone are present, for, although as a rule the elevators of the lids are associated in their motions, yet by an effort of the will most people can throw one of them into motion separately, or more than the other. No doubt the power to voluntarily innervate one levator and orbicularis alone varies in different individuals, and in many persons the levator centers are practically associated centers, and probably this is the reason why cerebral ptosis is rather rare. The position of this center is still an open question, but it is believed to be situated in front of the upper extremity of the ascending frontal convolution close to the arm center.

Ptosis, then, has no value as indicating the locality of a lesion in the cortex; but it may be of use in distinguishing a cortical lesion from one situated elsewhere in the brain, for monolateral ptosis, as the only focal symptom, occurs with cortical lesions alone.

It is probable that ptosis, as the result of a cortical lesion, is a distant symptom in not a few of the cases where it is present.

Ptosis on the side of the lesion has occasionally formed a symptom in disease of the pons without paralysis of the other branches of the third nerve—except, sometimes, in so far as conjugate deviation (*vide supra*) is concerned—and without the third nerve being involved in the lesion.

Again, ptosis, by forming a factor of a crossed paralysis, may serve to localize a lesion in the crus cerebri. When the third nerve is paralyzed by a lesion in this situation it is the rule to find it paralyzed as a whole, but paralysis of only some of the third-nerve branches may be produced by a lesion of the cerebral peduncle, and the branch to the levator palpebræ seems to be the one most frequently implicated alone.

In order now to complete this subject of ptosis as a focal symptom I must refer to a rare form of it which has been described by Nothnagel, and which does not depend on a lesion of the third nerve. It may be called sympathetic or pseudo-ptosis, and is accompanied by other eye-symptoms, as well as by symptoms of vasomotor paralysis of one side of the body, such as elevation of temperature, and redness and edema of the skin. In these cases, this author says, there is (1) apparent ptosis on the paralyzed side, owing to the contraction of the palpebral aperture, but the lid can be raised; (2) contraction of the pupil on

the same side; (3) a shrinking back of the eyeball into the orbit, so that it seems to have become smaller; (4) an abnormal secretion of thin mucus from the corresponding nostril, of tears from the affected eye, and of saliva from the corresponding side of the mouth. Nothnagel states he has found this train of symptoms in lesions of the corpus striatum.

A common sign of disease of the crus cerebri is what is known as crossed hemiplegia. Paralysis of the third nerve on the side of the lesion, with hemiplegia, hemianesthesia, often facial, and sometimes hypoglossal, paralysis of the opposite side of the body is a frequent form of it. The lesion may implicate all the branches of the third nerve or only some of them. But the localizing value of crossed hemiplegia, as Hughlings Jackson long ago pointed out, depends chiefly on the hemiplegia and paralysis of the cranial nerve coming on simultaneously. If they occur at different times they may be due to two distinct lesions, neither of which may be in the crus; for the hemiplegia might be due to a lesion in the hemisphere, and the third-nerve paralysis to a basal lesion of earlier or later date. Yet a few cases have been observed where, with a lesion in the cerebral peduncle, the third-nerve paralysis preceded the hemiplegia by a considerable interval.

That basal lesions are by far the most frequent cause of paralysis of the third nerve is beyond a doubt; and here it is usual, but not constant, to find it paralyzed in all its branches. The diagnosis to be made, when direct symptoms are being considered, is, for the most part, between a lesion in the crus and a lesion at the base. We cannot pretend to be able to make this diagnosis with certainty in all cases. Complete paralysis of every branch of the third nerve without any other paralysis is almost always basal; so also are those cases in which, where there is hemiplegia, it is slight as compared with the degree of the third-nerve paralysis; and those cases, too, to which I have already referred, where there is an interval between the onset of the paralysis of the extremities and of the third nerve, are apt to be basal. Of course there may be such a combination of paralyses of the other cerebral nerves with that of the third nerve as to leave no doubt with reference to the basal position of the lesion.

The third nerve may be paralyzed by lesions in the interpeduncular space, in which case the paralysis may be partial (ptosis alone, or abolition of upward and downward motion alone), complete, monocular, or binocular. When both nerves

are affected there is generally also paralysis of the other orbital nerves, or of the facial nerve; and hemiplegia or hemianopsia may also be present.

Thrombosis of the Cavernous Sinus invariably produces paralysis of the third nerve; but all the orbital nerves, as well as the fifth and the optic nerve, may also be involved, giving rise to complete immobility of the eye, with loss of conjunctival and corneal sensation. The pupil is usually contracted at first, but later on dilates. The venous obstruction causes exophthalmos, edema of the lids, and chemosis. Congestion papilla is sometimes found. The general symptoms are rigors, high temperature, and vomiting. Its principal causes are infective inflammation of the orbital cavity; erysipelas of the face; infective inflammation in the buccal, nasal, and pharyngeal cavities, and of the body of the sphenoid; and extension of thrombosis of the sinuses from purulent otitis. The thrombosis in more than half the cases spreads to the other side through the circular sinus. When the invasion occurs from the intracranial direction, pain in some or all of the branches of the first division of the fifth nerve is usually an early symptom.

Third-nerve symptoms—in addition to those included under the headings conjugate deviation, or paralysis, and ptosis—are sometimes distant symptoms. Tumors of the cerebral hemispheres, more particularly if accompanied by violent general head symptoms, indicating probably high intracranial pressure, are the lesions most apt to produce these distant third-nerve symptoms, As a rule the slighter the general cerebral symptoms, the more likely are the third-nerve paralyses to be direct symptoms. This rule, indeed, applies to other as well as to third-nerve focal symptoms.

Paralysis of the Fourth Nerve, when combined with paralysis of other motor eye-nerves, is difficult to recognize; and consequently in such cases it supplies but little aid for localization. Solitary paralysis of this nerve as a symptom of cerebral focal lesion is extremely rare. Nieden has placed a case on record in which paralysis of one-fourth nerve was the only focal symptom to which a tumor of the pineal gland, of the size of a walnut, gave rise. But the isolated fourth-nerve paralysis is more apt to be produced by a basal lesion. Pfungen has pointed out that, in meningitis, exudation in the space between the corpora quadrigemina and the splenium of the corpus callosum may implicate

the fourth nerves in the valve of Vieussens, and believes it is prone to do so in tubercular meningitis. In combination with paralysis of the third nerve it speaks for a lesion in the cerebral peduncle, extending back to the valve of Vieussens, and was, I believe, utilized clinically by Meynert in this sense.

When *Paralysis of the Sixth Nerve* occurs as the only focal sign it is probably due to disease at the base, or it is a distant symptom. There is no cranial nerve so liable to provide a distant symptom as the sixth. Gowers refers this liability to the lengthened course these nerves take over the most prominent part of the pons, which renders them readily affected by distant pressure. One or both nerves may in this way be paralyzed. Wernicke states that sixth-nerve paralysis is most apt to be present as a distant symptom when the lesion, especially a tumor, is situated in the cerebellum, differing in this way from the third nerve, which, as I have said, is more likely to give distant symptoms with a lesion in the cerebral hemisphere.

Paralysis of the sixth nerve, simultaneous in its onset with hemiplegia of the opposite side of the body, indicates a lesion in the pons, usually a hemorrhage, on the side corresponding to the paralyzed nerve. We know that the fifth and facial, and sometimes the auditory, spinal accessory, and hypoglossal nerve, may all, in varying combinations, form one of the elements in a crossed paralysis from a lesion in this position; but, if special localizing value is to be given here to the participation of any one cranial nerve, that nerve is the sixth. The paralysis of this nerve, simultaneously with palsy of the opposite side of the body, while other conditions point to an intracranial lesion, speaks, then, almost certainly for pontine disease.

Basal paralysis of the sixth nerve is frequently double, especially in syphilis. Fracture of the apex of the petrous portion of the temporal may also cause it.

Paralysis of the facial with the sixth is not an uncommon combination caused by a lesion in the pons, which at the same time produces hemiplegia of the opposite side of the body. This combination is a natural one, in view of the close relations of the nuclei of the sixth and seventh nerves. Indeed, according to Lockhart Clarke, Meynert, and others there is one nucleus which is common to both nerves—a view not shared in by Gowers and others. The manner in which the root of the facial nerve winds round the sixth-nerve nucleus must also have an important bear-

ing on the occurrence of associated paralyses of these nerves. (See also p. 207.)

Hemiplegia due to a lesion of the cortical motor region, which might happen to be combined with paralysis of the sixth nerve as a distant symptom, offers no difficulty in its diagnosis from hemiplegia with sixth-nerve paralysis in pontine disease; for, while the latter is a crossed paralysis, the former is homonymous.

Paralysis of the Seventh Nerve.—When lagophthalmos occurs as a symptom in focal cerebral disease, it is useful in localizing the disease by assisting in differentiating a lesion in the internal capsule, or in the facial motor center of the cortex, from one implicating the portio dura in the pons, as it is absent, or very slight, in the former cases, but very often markedly present in the latter. With a lesion in the lower part of the pons we are apt to have lagophthalmos with crossed hemiplegia; but if the lesion be in the upper part of the pons—the fibers from the opposite side having here joined the motor tract—the hemiplegia and lagophthalmos will be homonymous.

Paralysis of the Fifth Nerve, with hemiplegia of the opposite side, points to disease in the pons. Neuroparalytic ophthalmia is said to be the rule in basal lesions of the fifth nerve, and to occur very rarely in nuclear or fascicular lesions.

The *Orbicular Sign* may be noticed in some attacks of apoplexy with hemiplegia after consciousness has returned. It consists in this, that the hemiplegic person, who during health has been able to close each eye separately, and who even now can close both eyes together, or the eye on the sound side alone, is unable to close the eye on the paralyzed side by itself. This sign usually passes away after a short time. Sometimes when both eyes are closed it requires a greater effort to bring the eyelids together on the paralyzed side. I saw the orbicular sign very well marked and persistent in an obscure case of Dr. Wallace Beatty's where a gross cerebral lesion was suspected.

Extensive basal lesions, especially the syphilitic, may produce symptoms due to involvement of widely separate structures, without interfering with those which intervene; hence they tend to implicate several nerves without reference to system or function.

Convergent Concomitant Strabismus.—This is the condition which is popularly known as inward "cast" or "squint." It makes its appearance in children, when they begin to take an interest in small objects, such as toys and pictures; or a little later,

when the first lessons are learned—in short, when they begin to make frequent and prolonged demands on their internal recti and accommodation, most commonly from the age of three to six years.

The term "concomitant" (*concomitatus,* accompanied) is given to it in contradistinction to "paralytic" strabismus; because in it the squinting eye, by virtue of the normal innervation of the associated muscles, accompanies the straight one in all its movements to an equal extent. At the primary position of the eyeballs, in a case of concomitant squint, the parallelism of the visual axes is defective, and, as the eyes are moved from side to side, the defective parallelism continues in the same degree, neither increasing nor decreasing. Moreover, if the straight eye be shaded by the surgeon's left hand, and the squinting eye by this means be obliged to fix the object of vision—*e. g.,* the tip of the index finger of the surgeon's right hand held up two or more feet distant in the median line—it will be found that the straight eye is now squinting inwards. This deviation of the straight eye is called the secondary deviation, and, in these cases of concomitant strabismus, it is equal in degree to the primary deviation of the squinting eye; because the internal rectus of the good eye, being associated in its action with the external rectus of the squinting eye, when the latter muscle is forced, in the foregoing experiment, to roll its eye outwards in order to bring it to fixation, the internal rectus of the good eye, receiving a similar nervous impulse, rolls that eye inwards to the same extent as the squinting eye has been rolled outwards; and the good eye will therefore present, under the covering hand, an internal strabismus of the same amount as that which had previously been present in the squinting eye. This is an important point, for it is an aid in the differential diagnosis of this form of strabismus from the paralytic form, in which the secondary deviation is greater than the primary one (see General Principle No. 2, p. 448).

In order to decide which is the squinting eye it is merely necessary to direct the patient to look at an object held up in the median line on a level with his eyes, and a few feet in front of him.

In concomitant strabismus, of course, both eyes never squint simultaneously, as one hears it sometimes stated by parents.

Causes.—Squint is never due, as is popularly supposed, to fright, imitation, or naughtiness; nor is it ever brought on by

the patient looking at a lock of hair, or other object, which may happen to hang very much to one side.

Donders pointed out that in a large proportion of cases of convergent strabismus the refraction is hypermetropic, and he drew the conclusion that hypermetropia is to be regarded as the cause of the strabismus in the following way: It has been shown (chap. i. p. 22) that with each degree of normal convergence of the optic axes a certain effort of accommodation is associated. The greater the angle of normal convergence the greater is the possible effort of accommodation.

Of this physiological fact, Donders said, the hypermetrope often unconsciously takes advantage, and, in order to brace up his ac-




FIG. 147.

commodation in an excessive degree for the sake of distinct vision with one eye, he increases the angle of convergence of the optic axes by rotating the other eye (*L*, Fig. 147) somewhat inwards. The angle *l'* is thus made larger than the angle *l*, and the effort of accommodation normally belonging to the angle *l'* is obtained for the eye *R*, which consequently receives a clearer image of the visual object *A* on its retina. But, inasmuch as all hypermetropes do not squint, Donders considered that there were contributing circumstances, which caused each hypermetrope to unconsciously decide between distinct monocular vision with strabismus, and indistinct binocular vision. The latter, he said, is likely to be preferred if the condition of the refraction and the

acuteness of vision is the same in each eye; while, if the retinal images differ much, by reason of one eye being more ametropic than its fellow, from nebulous cornea, or from other causes, the desire for binocular vision would be less strong, and the imperfect eye would deviate inwards for the sake of the resulting increase of accommodation in the perfect eye.

It is admitted on all hands that hypermetropia is one of the causes of internal strabismus, but, as Schweigger [5] has pointed out, it is not the only cause, and probably not even the principal cause, for the following reasons: (1) If Donders' theory be complete, convergent strabismus must always appear, whenever there is binocular hypermetropia, along with the conditions which reduce the value of binocular vision. But strabismus is often absent, while the degree of ametropia is markedly different in the two eyes, or while the acuteness of vision is very defective in one eye. (2) According to Donders' theory, the higher the degree of the hypermetropia the greater should be the tendency to strabismus; and yet clinical observation shows that this is not the case. (3) In periodical strabismus, the influence of hypermetropia and of the accommodative effort is very evident; and yet these cases only go to show that, while hypermetropia is very frequently one of the causes of strabismus, it is not the only or most important one; for here, clearly, some factor necessary for the production of a permanent squint is wanting. (4) Donders' theory fails to explain the occurrence of convergent strabismus in emmetropic and in myopic individuals, where, of course, no excessive effort of accommodation is required.

Schweigger considers that a want of equilibrium between the muscles is the chief cause of strabismus (divergent as well as convergent), and that convergent strabismus is mainly due to a preponderance in the power of the internal over the external recti; or, with equal accuracy one might say, to an insufficiency of the external recti. It would seem that in hypermetropia the external recti are apt to be congenitally less powerful than the internal recti; while in myopia congenital insufficiency of the internal recti is the more common condition. The internal recti do, however, sometimes preponderate in emmetropia, and even in myopia; and convergent strabismus does sometimes occur in these forms of refraction. Whatever be the condition of refraction, strabismus is more apt to be developed if the value of binocular vision be diminished by imperfect sight in one eye.

Schweigger does not, however, give any proofs of this preponderance of certain muscles.

Spontaneous cure of strabismus sometimes takes place, most commonly between the tenth and sixteenth year of age. That it may happen with hypermetropia, and with defective vision in one eye, is strongly against Donders' theory.

According to Hansen Grut's view,[6] convergent squint originates in, and is maintained as the result of, an innervation which induces in the interni a shortening greater in amount than that which is desirable.

Single Vision in Concomitant Convergent Strabismus.—For the most part these patients do not complain of double vision, although diplopia is the rule in cases of convergent strabismus due to paralysis of the external rectus. Why is this? The image of the object looked at, it will correctly be said, must be formed in the squinting eye in each of these kinds of strabismus, on a part of the retina not identical with that in the fixing eye, but lying to the inside of it; and hence the image of the object should be projected by the squinting eye to its own side of the true position of the object (homonymous diplopia), and the latter should therefore be seen doubled. It is seen doubled in the paralytic form; why not also in the concomitant form? The only explanation of this circumstance, which, until within the last few years, had been put forward, was, that convergent concomitant strabismus being a quasi-physiological condition, the patient's mind involuntarily suppresses the annoying image belonging to the squinting eye in a manner analogous to that by which, when we are deeply interested in conversation, all extraneous sounds are unperceived, although they, too, must reach the nerve of hearing. This suppression of the image belonging to the squinting eye was believed to be the more easy owing to the indistinctness of the image itself, formed as it is on a peripheral part of the retina, while in the good eye it falls on the macula lutea. We often find, moreover, that the squinting eye is *ab initio* more defective (macula cornea, higher degree of hypermetropia, astigmatism, etc.) than its fellow, and it was held that this, too, rendered suppression of its image more easy. Such a suppression of the image is possible, and it no doubt does occur in many cases of strabismus; but it is certain, as pointed out by Schweigger, that it does not occur in all of them, perhaps not even in most of them. It would be beyond the

scope of this handbook were I to go into the arguments on this point. Suffice it to say that, in those cases where suppression of the image of the squinting eye does not take place, a certain participation in the act of vision on the part of this eye, when not too blind, is implied. One of two events takes place in those cases: Either the region of the retina, on which, in the squinting eye, the image of the visual object is formed, becomes functionally developed into a spot to a great extent physiologically "identical" with the macula lutea of the straight eye, and then something approaching normal binocular fusion of the images comes about, and hence single vision; or else diplopia is actually present, although, as a rule, it passes unnoticed by the patient, owing to its having become habitual to him. In some cases the first of these conditions is the actual state, in others it is the second which exists. I shall mention one fact in support of each, but must refrain from entering more deeply into the subject. In support of the first is the occurrence, not rarely observed, of crossed diplopia after operation for concomitant convergent strabismus, even when there is no divergence produced; and in support of the second, the diplopia which intelligent patients often admit, when they are carefully examined with the aid of a red glass before the good eye.

Amblyopia of the Squinting Eye.—In a large proportion of the cases of internal concomitant strabismus the squinting eye—even where there is no marked astigmatism, and where the media are clear—is amblyopic. Schweigger states the proportion of these amblyopic cases to be 30 per cent., but I believe the percentage to be much larger. It has been a very generally accepted opinion that this amblyopia is due to want of use on the part of the squinting eye, in consequence of the suppression of the image on its retina, and hence it is termed amblyopia ex anopsia. If this view were the correct one, we ought always to find only slight amblyopia of the squinting eye in children soon after strabismus comes on; while it should be of high degree—in fact, the eye should be almost useless—in adults who have not been operated on, and in whom monolateral strabismus had been present since childhood. And yet marked amblyopia may often be found in children in the squinting eye, while in adults the squinting eye often has very good vision—in short, the amblyopia of the squinting eye is not progressive, as it would be were it ex anopsia.

Again, many squinting eyes, when the straight eye is covered, instead of fixing the visual object with the macula lutea, remain unchanged in position, or even turn inwards more than before (amblyopia with eccentric fixation); and in less well-marked cases of the same sort, although there is no eccentric fixation, yet the preference for fixation with the macula lutea is lost, and uncertainty of fixation results, no one part of the retina being more useful for that purpose than another. It is held by many that this form is characteristic of amblyopia ex anopsia, and is the result of the strabismus; but it is identical with a form of congenital amblyopia often present in only one eye without strabismus. A strong argument in favor of amblyopia ex anopsia is the improvement which often seems to take place in the vision of the squinting eye by systematic separate use, or after the strabotomy. But, it is tolerably certain that, where the improvement takes place, the defective vision has been due rather to retinal asthenopia than to amblyopia; and if, at the outset, patients be pressed to discern the test-types, they often succeed in producing a better acuteness of vision than they at first seemed to possess. In many cases separate use fails altogether in improving the vision of the squinting eye, even when it is not very defective—a fact which is unfavorable to the amblyopia ex anopsia theory. The circumstance that in alternating strabismus the sight of each eye is good, cannot be regarded as proof in favor of amblyopia ex anopsia. I myself strongly incline to the views so ably put forward by Schweigger. The explanation which he gives of the very frequent presence of amblyopia in the squinting eye is that it is congenital; and, far from being the result of the strabismus, is a factor in its production, just as opacities of the cornea, or high degrees of ametropia, have always been admitted to be.

Priestley Smith [7] holds somewhat different opinions. To account for the phenomena of convergent strabismus, we have, he says, to consider the natural conditions of vision in early childhood. An infant's vision at birth is a mere perception of light, and his eye-movements are involuntary and purposeless. As the organs develop he learns to control these movements, to direct his eyes simultaneously to a given point, to fuse the two retinal pictures in one mental impression, and to recognize the forms of objects. During the first few years of life these newly acquired faculties are less stable than later on, and more easily

disturbed. Hence the greater liability to strabismus in infancy. Among three hundred and forty-seven cases where the onset age was ascertained, two hundred and fifty-four, = 73 per cent., began before the children were five years old. Three years old was the commonest age.

The hypermetropic child is specially liable to convergent strabismus because he has to overcome a special difficulty; he must learn to converge normally, while he accommodates abnormally. Failing in this, he squints in order to see clearly. Many squints arise in this way. But Priestley Smith says the influence of hypermetropia must not be exaggerated; most young children are more or less hypermetropic; the vast majority of hypermetropes of all degrees and all ages have no squint, and some squinters have little or no hypermetropia.

Infantile disorders—convulsions, whooping-cough, measles, a fright, a fall, etc.—are often the starting-point of strabismus, because the controlling influence of the higher brain centers is weakened at such times.

Priestley Smith thinks that a continuous squint involves weakening or loss of visual function, and that the younger the child the more readily does this occur. The sense of fusion, being no longer exercised, is gradually lost, and may prove irrecoverable a few years later, even though the eyes be put straight. Further, an eye which never fixes the object at which the patient looks, loses the power of true fixation. Such loss is found most often amongst cases of early onset and long duration; it is rarely, if ever, found until the squint has become continuous for at least six months. Again, it is probable that the early onset of strabismus, with complete disuse of the squinting eye, may arrest the development of form-perception in the latter, and thus render it permanently amblyopic.

We should, therefore, he thinks, treat strabismus as early as possible, to prevent secondary impairment of vision, or to correct such impairment by educative measures, if it be already present. By covering the good eye with a pad (almost always possible, with patience on the mother's part) we can oblige the child to use the neglected eye. We can thus preserve its function intact, or stimulate it if it be already weakened. We can even restore fixation-power to an eye which has already lost it, if the loss be not of too long standing. Spectacles to correct refractive errors, if required, are used in conjunction with the pad. At later ages

"bar-reading" and the stereoscope will aid in the recovery of binocular vision.

It is often right, Priestley Smith considers, to operate even at very early ages, when persistent use of the spectacles and pad gives no continuous improvement. To postpone operation in all cases to five or six years of age is often, he thinks, to lessen the likelihood of a complete cure.

There are *Three Clinical Varieties of Convergent Concomitant Strabismus.*—(1) Periodic. (2) Permanent alternating. (3)




Fig. 148.

Permanent monolateral. Periodic strabismus occurs only when some great effort of accommodation is required. It sometimes is the first stage of permanent monolateral or of alternating strabismus; but these two latter forms do not always have their beginning in the periodic form, which often continues as periodic to the end of the chapter. In alternating strabismus the patient squints sometimes with one eye and sometimes with the other.

In permanent monolateral strabismus the squint is confined to one eye.

Measurement of Convergent Strabismus.—The amount or degree of the deviation of the squinting eye from its normal position is not the same in every case, and the size of the squint is measured by one of the following methods. Whichever of them be used, it is important that the patient be directed, during the test, to regard a distant object placed in the median line and on a level with his eyes. If he look at a near object the squint may be overestimated, by reason of its increase with accommodation.

1. By the Linear Method we measure the number of millimeters by which the eye deviates from its normal position. The good eye is shaded, and the squinting eye is caused to fix an object in the median line—by preference a distant object. Close under the margin of the lid a strabometer (Fig. 148) is then placed, so that the 0 point may coincide with a perpendicular let fall from the center of the cornea. The shade being removed



FIG. 149.

from the good eye, the squinting eye is allowed to resume its abnormal position, and the degrees recorded on the instrument, under a perpendicular let fall from the center of the cornea in this position, are read off. They give the amount of the deviation.

2. Hirschberg's Method [8] consists in estimating the degree of deviation by the position of the corneal reflex of a candle flame held straight in front of, and about a foot from, the eye. Where there is no squint this reflex is situated at, or (with large angle γ) slightly to the inner side of, the center of the pupil in each eye. In a convergent squinting eye it is displaced outwards, and Hirschberg recognizes five groups of strabismus. Group 1 (Fig. 149 representing the right eye), in which the reflex is nearer to the center than to the margin of the pupil. This represents a

strabismus of less than 10°, and no operation is indicated. Group 2, in which the reflex is at or about the margin of the pupil, representing a strabismus of 12° to 15°, and indicating a simple tenotomy with occasionally a tenotomy of the other int. rectus. Group 3, in which the reflex is outside the pupillary margin, about halfway between the center of the pupil and the corneal margin. This represents a strabismus of about 25°, and indicates a tenotomy of the internal rectus, com-

FIG. 150.




FIG. 151. FIG. 152.

bined with a moderate advancement of the external rectus. Occasionally, later on, a tenotomy of the other internal rectus will be required. Group 4, in which the reflex is on or near the corneal margin; representing a strabismus of 45° to 50°, and indicating a tenotomy of the internal rectus, along with energetic

advancement of the external rectus, and sometimes a later tenotomy of the other internal rectus. Group 5, in which the reflex is on the sclerotic, between the margin of the cornea and the equator bulbi. This represents a strabismus of 60° to 80°, and requires the combined operation, with strongest possible advancement of the externus. Even this is sometimes insufficient, and a tenotomy of the internal rectus, or even the combined operation on the other eye, may be subsequently required. This is a modification of the linear method, and is a convenient one in routine practice.

3. Priestley Smith measures strabismus by means of a double tape (Fig. 150), used in conjunction with the ophthalmoscope, as shown in the accompanying figures. The patient places the ring *P* on one of his fingers, and holds it to his cheek. The observer places the ring *O* on the forefinger of the hand which holds the ophthalmoscope; this keeps his eye at a distance of one meter from the patient's face. He uses his disengaged hand as a fixation object for the patient, holding it edgewise towards the patient, and letting the graduated tape slide between his fingers. A small weight at the end of the tape keeps it stretched as the hand moves in either direction.

Fig. 151 illustrates the measurement of a convergent strabismus of the right eye. The patient, seated below the lamp and holding the tape as above described, is told to look at the mirror. The observer, holding the ring *O* and the mirror in the right hand, throws the light on the patient's left eye (*L*)—*i. e.*, the fixing eye. He sees the corneal reflex in the center of the pupil, and knows thereby that this eye is fixing properly. He then throws the light on the right eye (*R*), and sees the reflex situated eccentrically outwards, and knows that this eye deviates inwards. Taking the graduated tape between the fingers of his left hand, and telling the patient to watch this hand, he moves it outwards along the tape (see Fig. 150), and meanwhile watches the corneal reflex in the deviating eye. When this latter reaches the middle of the pupil he reads the position of the hand upon the tape. The axis of the deviating eye (*R*) has moved from *R'* to *O*, through the angle *R' R O*. The axis of the non-deviating eye (*L*) has moved through an equal angle (*O L L'*). The angular movement of *L*, as measured by the tape, equals the angular deviation of *R*.

Fig. 152 illustrates the measurement of a divergent strabismus

of the right eye. The hands must be reversed, but the principle is the same as before.

The graduated tape is a substitute for a graduated arc of a circle, but does not exactly correspond with such an arc; the error involved is, however, so small as to be of no importance if the observer keep his two hands at about the same distance from the patient's face. In this mode of measuring a strabismus it is the excursion of the fixing eye which is actually measured, and the excursion of the deviating eye is supposed to be equal to it. If the excursions of the two eyes are unequal, the result would




FIG. 153.

be at fault. The method, though difficult to explain in words, is very quick and satisfactory in practice.

4. The Angular Method.—The object aimed at here is to determine the size of the angle, which the visual axis of the squinting eye makes, with the direction it should normally have. For this purpose a perimeter is employed. Let us suppose that the right eye (*R*, Fig. 153) be the squinting eye, and that *P o P* be the arc of the perimeter. The patient is placed at the instrument, as though the field of vision of his squinting eye were about

to be examined. He is directed to look at a distant object (*A*) with his good eye (*L*). The visual line from *R* should now pass through the point *o,* but it passes through the point *n,* and therefore *o R n* is the angle of the strabismus. The surgeon finds the position of *n* by carrying the flame of a candle along the perimeter, until, with his eye placed behind the flame, he finds that the corneal image of the flame occupies the center of the pupil. The flame itself will then be at *n,* and the size of the squint-angle may be read off there. This gives us the optical axis of the eye; but, to be strictly accurate, we must remember that the position of the visual axis is what we require, and that it lies a few degrees farther inwards, according to the size of the angle γ. The angular method is now in general use instead of the linear method, than which it is more accurate.

5. A good subjective method for determining the dimension of a strabismus, but which can only be used where diplopia is present, is what may be called the Method by Tangents. Upon a wall of the consulting-room, in a horizontal line, and so as to be on a level with the eyes of the patient who is placed about 3 meters from the wall, are, permanently marked out, tangents of angles of 5° each, as seen from the place where the squinting eye is. Exactly opposite to the squinting eye is 0°, while towards the right and left the points are marked up to 45° or more. The flame of a candle being held at 0°, and one eye of the patient being covered with a red glass, he is called on to indicate the position of the image belonging to the squinting eye, and the number on the wall which corresponds to this gives the angle of the strabismus. For the purpose of estimating paralyses of the orbital muscles, a similar row of tangents, or several such, may be marked out in the vertical direction.

Mobility of the Eye Outwards in Convergent Concomitant Strabismus.—This is often defective in the squinting eye, and sometimes also in the fixing eye. The excursiveness of the lateral motions of the eyeball may be measured by the perimeter. Having placed the patient as though the field of vision, say of his right eye, were about to be examined, he is directed to follow with his eye the flame of a candle carried along the perimeter from 0° towards 90° in the temporal direction; and when it is found that the eye cannot be turned any farther in this direction that extreme position is noted by the position of the candle at the perimeter. The corneal image of the flame must, of

course, be central when the position of the flame is read off. In a similar way the mobility of the eye inwards may be measured. In the normal eye the mobility in each direction is about 45°. In strabismus we simply compare the outward mobility of the squinting eye with that of the good eye, to ascertain how much, if anything, the former lacks of its normal amount.

Treatment.—The bearing of hypermetropia on the production of many cases of strabismus long since suggested the idea of curing the deviation by spectacles, which would correct any existing hypermetropia. The accommodation having been paralyzed by atropin, and kept under its influence for some weeks or months, spectacles which completely correct the hypermetropia and astigmatism are meantime constantly worn. Should the patient require to use his eyes for near work while under treatment, it is necessary that he should have suitably higher + glasses for his near work. Occasionally good cures are effected by this means, and even when atropin is not used; and when a periodic strabismus in a child comes under my care I always think it worth while to attempt its correction in this way; but in general it is, by itself, of no use whatever.

Orthoptic Treatment.—To Javal [9] is due the credit of devising this method; but although he did so some years ago it is only recently that the treatment has been introduced into practical ophthalmology.

In order that the treatment may be carried out it is necessary, in the first instance, that the strabismic person should have diplopia. If the latter be not present spontaneously it has to be developed; and it is sometimes possible, when the sight in the squinting eye is not too defective, to give the patient diplopia—*i. e.,* to make him continuously conscious of the presence of the image belonging to the squinting eye. This may be done by means of exercises with a prism, base downwards, before the deviated eye, and a candle flame as visual object. The exercises are to be repeated daily until diplopia without a prism is established. Javal recommends the following exercise to develop diplopia: A screen—*e. g.,* a large sheet of cardboard—is held vertically between the two eyes, while the patient is directed to look at a candle flame about 2 meters in front of him. Double vision may immediately appear; but, if it does not, it may be brought out by now and then covering the good eye for a moment, or by placing before it a red glass, which can soon be done without

Less brilliant visual objects are gradually substituted, until, finally, the double vision will continue even when, at first cautiously, the screen is removed.

Double vision having been established, we proceed to enable the patient to fuse the double images—*i. e.,* to obtain binocular vision—and when we have succeeded in doing this we have cured the squint. The end in view is best effected by means of a stereoscope, into which, in place of the usual prisms, + 6 D lenses have been introduced. The focal distance of these lenses being about the length of an ordinary stereoscope, rays coming from the slides, and passing through them, fall into the observer's eye as parallel rays; the accommodation consequently is suspended, and under normal conditions the visual lines are parallel, as though looking at a distant object. In the normal state the double picture, or diagram, will seem to be single, but to the strabismic patient in whom diplopia is present it will be double. Our duty, then, is to diminish the distance between the pictures, until the patient finds himself just able to fuse the images into a single impression. After a day or two the distance is increased slightly, and so on, until, finally, the normal position is reached. It is needless to say that in these exercises all errors of refraction must be eliminated by the proper glasses.* Priestley Smith's fusion

* The existence or otherwise of true binocular vision may be ascertained by the simple experiment of giving the patient a book to read, and holding a cedar pencil halfway between his eyes and the page at right angles to the lines of type. If binocular vision be present, the pencil will not offer any impediment to the reading; but if it be not present, parts of the page will be hidden behind the pencil. The reader may prove this by performing the experiment on himself, first with both eyes open (binocular vision), and then with one eye shut.

Another method is that known as Hering's Drop Experiment. A cylinder about 25 cm. long, and wide enough to take in both eyes of a person, is provided—at the opposite end from that placed around the eyes —with two strong wires 18 inches long, which jut out in continuation, as it were, of the cylinder, but which are bent outwards sufficiently to keep them out of view of the patient. Between the ends of these wires a fine thread is stretched, with a small bead fastened at its middle point, so that the bead may occupy the center of the field when the patient looks through the cylinder. During the experiment the thread is in the horizontal position, and the bead is used as the patient's fixation point. Small balls of different sizes (peas, beans, etc.) are then let fall from a height, one after another, a couple of dozen times or more, some of them in front of the thread, some of them behind it. If the patient have normal binocular vision, he will be able to say each time with certainty whether the ball falls in front of or behind the thread; but if he have not true

tubes are also useful for bringing about fusion of the double images.

The pictures used in the stereoscope should be geometrical figures, or especially designed pictures, in order that both surgeon and patient may the more readily recognize their fusion.

Only the very slight degrees of strabismus are adapted for the attempt at cure by orthoptic treatment. A marked deviation will not be amenable to it. Moreover, it makes demands both upon the patience and intelligence of the patient, which are rarely fulfilled, especially in hospital practice. A field more fertile in good results for the employment of this treatment is found in the completion of cures which have been commenced by operative measures.

Operative Treatment.—Division of the tendon of the internal rectus muscle, combined, sometimes, with advancement of the insertion of the external rectus, is the measure which has to be applied in most of the cases which come under our notice. I am strongly opposed to operative interference in patients under five years of age, and very much prefer that they should be seven or eight years old, or even older. Early childhood offers a decided obstacle to the careful adjustment of the operation and to orthoptic treatment.

In order that the operative proceeding may be adapted to each case, the following points must have been previously noted with care: (1) The dimension of the strabismus angle. (2) The lateral mobility of the eyes, especially the mobility outwards of the squinting eye. (3) The refraction of the eyes, and the acuteness of vision of the squinting eye, as well as the presence, or otherwise, of diplopia: the first in order that glasses for the correction of any hypermetropia may be worn, if desirable, after the operation; the second, because, *ceteris paribus,* an operation for convergent strabismus will produce a more marked effect if the vision in the squinting eye be good than if it be very defective; and the third, because the presence of diplopia encourages the hope that binocular vision may be restored.

Rules which will insure in every case, with absolute certainty, the desired degree of operative effect cannot be laid down. The following will be found to answer in the majority of cases, and if the effect be now and then too great it can easily be ad-

binocular vision, if only one eye be used, he will merely guess at the position of the falling ball, and will make frequent mistakes.

justed by bringing forward the internal rectus, or by setting back the external rectus, within a few days after the operation. In every instance it should be the desire of the surgeon to leave 2° or 3° of strabismus behind; for the effect of the operation is apt to increase within a year, and, if absolute parallelism be present at first, divergence may ultimately supervene. The establishment of binocular vision when possible would do away with this remnant of strabismus; but under any circumstances the latter does not detract from the cosmetic result.

If the vision of the squinting eye be fairly good, and the deviation amount to not more than 15° or 20°, and the power of the external rectus be sufficient, the correction can be effected by the tenotomy of the internal rectus of the squinting eye. A strabismus of 20° will require the free separation of the delicate connections between the anterior surface of the tendon, or capsule of Tenon, and the conjunctiva as far back as the caruncle, in order that the tendon may be free to contract. For a deviation of 15° or less this separation should not be so free, or should be quite omitted; or, if a very slight effect be desired, it can be produced by drawing the conjunctival wound together, after an operation which has been confined strictly to the insertion of the tendon.

If the vision of the squinting eye be fairly good, and the power of the external rectus sufficient, and if the squint be more than 20°, it is advisable to divide the proceeding between the eyes—*e. g.*, if it be 30°, about 20° are corrected by tenotomy of the internal rectus of the squinting eye, and the remainder by tenotomy of the internal rectus of the fixing eye. If desired, the effect of the tenotomy in one or both eyes may be increased by a suture passed through a fold of conjunctiva at the outer side of the globe, and tied tightly.

If, although the vision of the squinting eye be good and the deviation not more than 20° or 25°, there be marked loss of power of the external rectus muscle, tenotomy of the internal rectus alone will often lead to disappointment, and a good result will require this tenotomy to be combined with advancement of the external rectus, the operative measures being confined to the squinting eye. But advancements in such cases as this must be very cautiously carried out, as an excessive effect may easily be produced. The external rectus should be but slightly brought forward.

If the deviation exceeds 35°, even when there is good vision

in the squinting eye, and no loss of power in the external rectus, tenotomy of the internal rectus of each eye is rarely sufficient, and as a rule advancement of the external rectus of the squinting eye must be combined with these measures.

With a deviation of 30° to 35°, and loss of power in the external rectus, the demand for advancement of the external rectus becomes more imperative. The correction of squints of 40° and more are, in every instance, to be effected by tenotomy with vigorous advancement in the squinting eye, and subsequent tenotomy of the internal rectus in the good eye.

In cases where the vision of the squinting eye is much reduced, the deviation great, and the insufficiency of the external rectus marked, the combined operation in one or both eyes is the proper proceeding.

Mode of Operating for Strabismus. *Tenotomy.*—The instruments required for this operation are a spring-stop speculum, a small-toothed forceps, a blunt scissors somewhat curved on the flat, and two strabismus hooks (Fig. 154).

The eye having been thoroughly cocainized, the patient is placed on his back, the surgeon standing in front of him and on his left-hand side if the left eye is to be operated on, or behind him if it be the right eye. The speculum is then applied, and the conjunctiva over the insertion of the tendon of the internal rectus is seized with the forceps, and incised with the scissors between the forceps and the eye. Into the opening thus made the points of the closed scissors are inserted, and, with a snipping action, a passage is made through the subconjunctival tissue—from the conjunctival aperture to the upper border of the tendon in case of the left eye, or to its lower border in the right eye. The scissors are now laid aside, but the conjunctiva is still held in the forceps, and with the right hand, the point of the hook is passed through the opening and along the passage, until the edge of the tendon is reached. The point of the hook being kept in contact with the sclerotic, the instrument is then turned rapidly round and under the tendon, and is brought close up to the insertion of the latter into the sclerotic, care being taken that the whole breadth of the tendon lies on the hook. The forceps are now laid aside, and the hook carrying the tendon is transferred to the left hand. One blade of the scissors (held in the right hand)



Fig. 154.

is now inserted between the globe and the tendon, and the latter is completely divided at its insertion. The second hook is then employed for searching, above and below, for any strands of the tendon which may be left undivided, the test for complete division being that the hook can be brought up without obstruction to the margin of the cornea. If the smallest segment of the tendon be left undivided, the result of the operation is apt to be unsatisfactory. Immediately after the operation a marked diminution in the mobility of the eye inwards should be looked for, as this motion can now only take place by aid of any remaining connective tissue attachments of the muscle to the eyeball and capsule of Tenon. If this defect in motion be not present, or in only a slight degree in comparison with the supposed extent of operation, it may be concluded that the tendon is imperfectly divided, and a new search with the hook for undivided filaments must be made. To estimate this loss of motion it is necessary before the operation to note the degree of mobility of the eyeball inwards, and to compare it with the inward motion of the other eye.

The effect of the operation may be diminished, if found necessary, by drawing the edges of the conjunctival wound together with a suture, the tendon being thus prevented from uniting with the globe so far back. The more conjunctiva we include in the suture at each side of the wound the more will the effect of the tenotomy be reduced. This restricting suture should be applied when the immediate result of the tenotomy is greater than expected or desired.

As the edges of the conjunctival wound cannot be accurately adjusted with sutures, none are applied for that purpose. They are used, as above, to diminish the operative effect; or, when an extensive loosening of the subconjunctival tissue has been performed, to prevent sinking of the caruncle.

The Subconjunctival Operation for Strabismus, proposed by the late Mr. Critchett, is performed as follows: A fold of conjunctiva is seized close to the lower margin of the insertion of the muscle, and incised with a blunt-pointed scissors, so as to expose the tendon. A strabismus hook is passed through the opening and under the tendon. The scissors is now inserted and opened slightly, one point being kept close to the hook, while the other is passed between the tendon and the conjunctiva, and the tendon is divided at its insertion. This method is very generally

adopted by English surgeons. For myself I prefer the operation (von Graefe's) previously described, as it much more readily admits of modifications of the effect.

In von Arlt's Method, instead of a hook being passed under the tendon in the first instance, it is seized with the forceps with which, just previously, the conjunctiva had been raised. In other respects the proceeding is the same as von Graefe's, than which it is said to be less painful.

The immediate and ultimate effects of a tenotomy are by no means identical. Immediately after the operation the effect is very marked, owing to the loosening of the tendon from its insertion. In a few days, when it becomes reattached, the effect diminishes, and in the course of some weeks there is again an increase in the effect, and this increase continues for about a year, as above stated.

The ultimate result may, with tolerable certainty, be estimated immediately after the operation by testing the power of convergence. If the patient be directed to look with both eyes at the surgeon's finger held in the middle line, and it be approached to within 12 or 15 cm. of his nose, and if the convergence of the eyes can be maintained at that distance, the effect will not be too great. But if at a distance of from 18 to 20 cm. the operated eye ceases to converge, or begins to diverge, or if even at 12 cm. the convergence, although accomplished, cannot be maintained for more than a few moments, and if then the operated eye deviates outwards, ultimate divergence may be expected, even though the actual position of the visual axes be correct. A restricting suture must be applied in such cases.

Sometimes, although the patient converges up to 12 cm. satisfactorily, and maintains the convergence at that distance for some moments, the eye will then rotate inwards. In such cases there is apt to be a recurrence of the strabismus.

Advancement.—In cases of convergent squint, in which it is desirable to combine advancement of the external rectus with tenotomy of the internal rectus, the latter is done first, as above described, at the same sitting.

An opening is then made in the conjunctiva immediately over the insertion of the external rectus, and as long as the breadth of the tendon. The band of conjunctiva between the opening and the cornea is separated up with the scissors from the sclerotic, for to it the tendon has to be fastened later on. A strabismus

hook is now passed under the tendon, and brought well up to its insertion, care being taken that the whole width of the tendon is held on the hook. A needle carrying a fine silk suture is introduced from its upper margin between the tendon and sclerotic, and passed through the tendon at its middle line. In the same way another suture is passed behind the tendon from its lower margin, and through it, close to the first suture. Each of these sutures is knotted firmly on the tendon, a long end being left to each suture (Fig. 155). The tendon is separated off with the scissors from the sclerotic close to its insertion. The sutures are passed through the conjunctival flap in the direction of the muscle, and are respectively tied with their own ends. A greater or less effect is produced according as the sutures are placed farther or nearer to the insertion of the tendon, and according as they are




Fig. 155.

drawn more or less tightly. I have found this method perfectly satisfactory.

Immediately after the combined operation is finished there should be no divergence, nor should there be marked loss of motion of the eyeball inwards. In either case the effect is too great, and must at once be diminished by an adjustment of the advancing sutures, or a bringing forward of the internal rectus.

In my opinion, even if it lie in the plan of the treatment to supplement the tenotomy (or combined operation) on the squinting eye by a tenotomy (or combined operation) on the fixing eye,

both eyes should not be operated on at one and the same sitting. An interval of a fortnight or more should elapse, in order that the true effect of the first proceeding may be accurately gauged, and then the surgeon will be in a position to know how to regulate his operative measures for the other eye.

After a strabismus operation, a light dressing is applied, and is changed morning and evening for forty-eight hours, when, if no suture has been used, it may be discarded. If sutures have been employed, the dressing is retained until they come away.

Dangers of the Strabismus Operation.—I have never seen any inflammatory reaction after strabismus operation, not even after an advancement, nor have I ever seen any serious accident during the operation. Puncture of the sclerotic with the scissors while the tendon was being divided has occurred in the hands of some operators; but I confess I cannot understand how such an accident could happen, unless the operator had his own eyes shut. It is also stated that eyes have been lost after squint operations through orbital cellulitis, which, beyond doubt, must have been brought on by the introduction of septic matter upon the instruments.

Occasionally a small arterial branch may be divided during the operation, and this, bleeding into the capsule of Tenon, may cause rather alarming exophthalmos. The protrusion goes back in a few days with use of a pressure bandage. I have only seen the occurrence twice.

Sinking back of the caruncle, some months after the tenotomy, when it does rarely occur, can be remedied in the following way: The conjunctiva is divided vertically about 6 mm. from the caruncle. The inner lip of the wound is raised, a scissors curved on the flat passed in, and the subconjunctival tissue as far as under the sunken caruncle separated. The subconjunctival tissue under the outer lip of the wound, and as far as the corneal margin, is loosened in the same way, and the two flaps are brought together with a suture, which includes a sufficiency of conjunctiva to draw the caruncle well forwards.

Treatment subsequent to Operation.—It is generally necessary for the patient to wear the correcting spectacles for his hypermetropia either constantly or for near vision only, according as the result of the operative measures makes it more or less desirable to suspend the accommodation. After some months it is usually possible to leave off the spectacles, except for near vision.

A cure of the strabismus, in the sense of removal of the deformity, can be attained by operation, and by itself affords ample indication for the operation. But a cure, in the true sense of the term, involves restoration of binocular vision,* and this is very rarely obtained by operative measures alone. To this end the operation must be followed up by orthoptic treatment.

Insufficiency of Convergence, or Insufficiency of the Internal Recti Muscles, and Divergent Concomitant Strabismus.—In the normal condition the orbital muscles are in a state of equilibrium, no one muscle or pair of muscles having more power over the eyeballs than its fellow.

Insufficiency of the Internal Recti Muscles, or Insufficiency of Convergence, as it is more correctly called, implies a disturbance of this equilibrium. The converging power of the internal recti, in these cases, is so much weakened that they are obliged to make a constant effort to prevent the eyes, or one of them, from becoming divergent, and it is only the demand for binocular vision which stimulates the muscles to this effect.

Muscular Asthenopia is the symptom caused by this insufficiency. The patients complain that, after reading, writing, sewing, or employment at other near work for a time, they begin to find the objects spreading, becoming indistinct, and perhaps doubled. Pain in and about the eyes comes on. These symptoms gradually increase, until the work has to be discontinued.

A great deal has been written within recent years upon the relationship of some nervous diseases, especially epilepsy, to want of power in one or more of the orbital muscles. It has been thought that "eye strain," from want of co-ordination in these muscles, sometimes aggravated, if it did not actually cause, epilepsy; but the outcome of the whole discussion seems to be that there is no such connection.

The diagnosis of insufficiency of convergence can be made by the following methods:

(*a*) The patient is directed to look at the tip of the surgeon's finger held up in the middle line. The finger is brought slowly closer to the eyes until a certain point is reached where the internal rectus of one eye ceases to act, the other eye still remaining in

* The importance of binocular vision consists in the fact that it is chiefly by its aid we estimate distances finely and observe the shape of objects. Even plane surfaces are seen much more accurately with binocular than with monocular vision.

fixation. The first eye, upon the finger being advanced a little more, usually becomes divergent.

(*b*) If the tip of the finger be held some 20 cm. from the patient's eyes, and if, with his other hand, the surgeon cover one of the eyes, say the right, while the left is caused to fix the finger-tip, it will be found that the eye under the hand is diverging, and, when the hand is removed from it, it makes an inward motion, in order again to fix the finger-tip. The explanation of this is that when one eye is covered there is nothing to be gained in the way of single vision by an excessive exertion of the weak internal recti; and consequently the eye which is excluded from the act of vision is abandoned to the control of the external rectus, and only returns to its normal position when, being restored to participation in the act of vision, diplopia would otherwise be present.



Fig. 156.

(*c*) The following is Von Graefe's Test for insufficiency of the Internal Recti: A dot with a fine line drawn vertically through it (Fig. 156) on a sheet of white paper is given to the patient to look at, at his usual reading distance. Before one eye, say the right, a prism of about 10° with its base downwards is held vertically. This, in the normal condition, would produce a double image of the dot, so that the figure would seem to be a line with two dots, the upper dot being the image belonging to the right eye. In insufficiency of the interni, the image of the right eye would not only be higher than that of the left, but it would also stand to the left (crossed double images) more or less, so that here the picture is that of two lines, each with a dot, the upper line and dot standing to the left-hand side (Fig. 157). This crossed diplopia indicates divergence. The explanation of the experiment is as follows: When a prism is held before the right eye, the possibility of binocular vision is removed, and, insufficiency existing, the weak internal rectus of the right eye has no object in greatly exerting itself, and, consequently, abandons the eye to the traction of the external rectus. Hence the divergence and the projection of the image of this eye to the opposite side. The degree of insufficiency existing may be determined by this same experiment. If a weak prism be held with its



Fig. 157.

base inwards before the left eye, in the above case, the images of the lines will appear to be brought closer. By gradually proceeding to higher prisms, one will be found which brings the lines together, so that the picture will now be that of two dots over each other on one line. This prism is the measure of the insufficiency.

(*d*) Landolt estimates the amount of insufficiency of convergence by means of the meter angle and amplitude of convergence. For an account of the method I must refer the reader to his valuable work.[10]

(*e*) Maddox's Rod Test is an admirable method for ascertaining the condition of the muscular equilibrium of the eyeballs, and for estimating any existing derangement of it.

The apparent lengthening of a flame into a line of light, when looked at through a strong cylinder, is utilized to make the two images so dissimilar that no desire to unite them remains. The chief advantage of this principle is that slight malpositions do not,




Fig. 158.

as with prisms, vitiate the result materially. A glass rod mounted in a circular metal disc, as in Fig. 158, may be used; or a plano-cylinder with a radius of about 20 mm.; or a piece of corrugated glass; or a flat series of thin glass rods side by side. The best flame to employ is that of a gas-jet turned low, at a distance of 5 mm. or 6 mm., and the appearance is improved by a piece of blue glass before the other eye, to equalize the illumination of the two images. The line of light is at right angles to the axis of the cylinder. If it pass through the flame, the balance is perfect; if not, the defect is measured by the deviating angle of the prism which is found to bring them together, or, preferably, by a litho-

graphed scale, placed with its zero just behind the flame, so that the figure crossed by the line of light gives the deviation in degrees. For vertical diplopia the scale should be vertical, and for horizontal diplopia horizontal. In either case the axis of the cylinder should be parallel to the scale. When the cylinder is vertical it should be shaded from the light of the window. By placing the patient's head in different positions the diplopia can be measured in all parts of the motor field. Vertical and horizontal scale should, for this purpose, be fixed on the wall, with their zeros coinciding at the position of the flame. For near-vision tests a flame is too large. A scale should be used on a black background, with a small silvered hemisphere or bead fixed to its zero, to be a source of reflected light from the window or from a flame.

This test is also very serviceable in overcoming the suppression of the false image in old squints, and for discovering the latent paresis of an ocular muscle.

Insufficiency of the internal recti is a common attendant upon myopia, but it is also found with emmetropia, and even with hypermetropia.

Concomitant divergent strabismus is a further development of the same condition.

Treatment.—In moderate degrees of myopia the use of such concave glasses as will permit the patient to read at 35 cm. distance may relieve the asthenopic symptoms.

Decentration of these glasses may give further aid. If the glasses be so set in the spectacle-frame that their centers are on the outer side of the visual lines, the inner half of the glasses act as prisms with their bases inwards, and by them the rays are broken inwards—*i. e.,* towards the macula lutea in each eye, so that a slight divergence may take place without diplopia, etc. In this way the internal recti are relieved. Should the case be one demanding the use of convex glasses (hypermetropia, presbyopia) the decentration must be inwards.

A more perfect and accurate method is that of prescribing prisms, bases inwards, to be worn for reading and other near work. These may be combined with concave or convex glasses, where such are indicated. The prism which is the measure of the insufficiency (see above) is divided between the two eyes. If it be 4°, a prism of 2° is placed, base inwards, before each eye for near work. Very high prisms cannot be ordered, owing to the

color effects they produce; and in cases where they would be required the insufficiency can be only partially corrected by prisms.

Operative Treatment.—This consists in weakening the too strong external rectus by tenotomy. The danger of the method is that convergent strabismus with homonymous diplopia for distant objects may result, unless the case be suitable for operation. Only those cases are suitable in which absolute divergent strabismus is present; or those in which, with a prism of not less than 10°, base inwards, before one eye, the flame of a candle at 3 m. distance is seen single, or if it be perhaps doubled for a moment, then becoming again single. When with such a prism single vision is present, the external rectus by an effort must have overcome the effect of the prism, and it is admissible to deprive the muscle of the power represented by that effort or prism. If diplopia be produced by a prism of 10°, the tenotomy is contraindicated, for the effect of the latter could not be modified to the slight power of abduction indicated by a weaker prism. A source of error in the ascertaining of this abduction prism, which must be guarded against, is that the patient may suppress the image of one eye, and that his single vision may be merely due to the fact that he is seeing with the other alone. The higher the abduction prism, the more extensive may be the division of the subconjunctival tissue, etc., while with weak abduction, the effect must be diminished by a conjunctival suture.

Immediately after the operation there should be a certain amount of convergence, as shown by homonymous diplopia in the middle line for the flame of a candle at 3 m. distance. This convergence, or diplopia should not be greater than can be corrected by a prism of 10°. Moreover, if the candle be moved from the middle line 15° to the opposite side from the operated muscle (to the right if the left external rectus has been tenotomized), there should be no convergence (no diplopia), and a vertical prism before one eye should only cause double images placed directly over each other. If, by these experiments, it be shown that the operation has produced an excessive effect, the latter must be diminished by a suture drawing the lips of the conjunctival wound together, and including more or less conjunctiva, according to the excess to be corrected. Or, if a suture have already been applied, and the result be still in excess, it must be withdrawn, and a still more restricting suture inserted. In all these cases, convergence must necessarily be present, when the candle is carried over to the

side of the operated muscle; but this disappears—except perhaps at the very most extreme position on that side—as also the convergence in the middle line, by reason of cicatricial contraction at the new insertion of the tendon; always provided that the indications for the operation and its performance, as above set forth, have been accurately attended to.

Nystagmus.

This term indicates an involuntary oscillation of the eyeballs laterally (the most common form), vertically, or with a rotary motion (caused by the oblique muscles).

It is most commonly found with congenitally defective vision—microphthalmos, coloboma of the chorioid, in albinos, etc.; but it may be acquired, and is often seen, in those employed in coal-mines. It occurs in about one-half the cases of disseminated sclerosis.*

In the congenital cases it is probable that the absence of the stimulus which accurate retinal impressions afford interferes with the functional development of the co-ordinating centers for the orbital muscles. In coal-mines, the very defective light, and the blackness of the surroundings, deprive the miners of any defined retinal impression, and hence their co-ordinating centers are apt to become deranged. But as it is chiefly those who work in one constrained position on their sides, with eyes directed obliquely upwards, who become affected, it seems probable [11] that this unnatural, and long-continued direction of the eyeballs is an important factor in the production of the affection; indeed, it may be to a great extent a professional cramp, like writer's cramp. In fact, a case of acquired nystagmus in a compositor, due to working in a strained position, has been recorded by Snell.[12]

Those patients, in whom nystagmus is due to a congenital defect of vision, do not complain of oscillation of the objects they look at; but individuals, who become affected with it in later life, are much troubled with that symptom, especially at the onset.

Treatment.—In congenital cases, which admit of improvement of vision, a cure, partial or complete, is sometimes brought about

* According to Gowers ("Diseases of the Nervous System," vol. i. 2d Ed.), nystagmus occurs often in ataxic paraplegia, primary spastic paraplegia, and hereditary ataxia, sometimes in severe multiple neuritis and syringomyelia, but rarely in progressive muscular atrophy.

when the vision improves. If strabismus be present it should be cured, after which a diminution in the oscillations may result. In miner's nystagmus, the all-important measure is a permanent relinquishment of mine work; and this is frequently followed by satisfactory results.

References.

[1] Landolt and Wecker, "Traité d'Ophtal.," vol. iii. p. 782.
[2] "Ophthalmic Review," v. p. 65.
[3a] "Trans. Ophth. Soc., U. K.," ix. p. 191.
[3b] "Brit. Med. Jour.," March 3, 1900.
[3] "Brain," Summer, 1900.
[4] First proposed by Professor J. Michel, "Klin. Monatsbl. f. Augenheilk.," 1887, p. 373.
[5] "Ueber das Schielen" and "Handbuch der Augenheilkunde," 5th ed., p. 146.
[6] "Bowman Lecture," 1889.
[7] "Bowman Lecture," 1898.
[8] "Centralblatt. f. p. Augenheilkunde," 1886, p. 5.
[9] "Annales d'Oculistique," Juillet et Août, 1871. See also Mars et Avril, Mai et Juin, and Nov. et Déc., for the same year.
[10] "The Refraction and Accommodation of the Eye," p. 501.
[11] *Vide* S. Snell, "Brit. Med. Journ.," July 11, 1891.
[12] "Trans. Ophthal. Soc., U. K.," xi. p. 102.

CHAPTER XVIII.

DISEASES OF THE ORBIT.

Orbital Cellulitis.—*The Symptoms* of this affection are: erysipelatous swelling of the lids, especially of the upper lid; serous chemosis; pain in the orbit, increased on pressure of the eyeball backwards; violent facial neuralgia; exophthalmos, with impairment of the motions of the eye in every direction; and high fever.

Vision is not generally affected, but sometimes it is so from optic neuritis, and then, too, mydriasis is seen. The cornea is often completely or partially anesthetic.

The surgeon, by pressing the tip of his fourth finger between the eyeball and the margin of the orbit, may feel a more or less resistant tumor. This gradually increases in some one direction, the integument in that position becomes redder, fluctuation becomes pronounced, and the abscess finally opens through the skin, or into the conjunctival sac, the pointing being usually at the upper and inner angle of the orbit. Restoration to the normal state, as a rule, comes about; but in some cases complete atrophy of the optic nerve supervenes.

Causes.—(1) Idiopathic (*e. g.,* cold); (2) traumatic (perforating injuries, foreign bodies); (3) extension of inflammation from surrounding parts (erysipelas, diseased tooth, ethmoidal cells); (4) metastasis, (pyemia, metria); (5) sequelæ of fevers (scarlatina, typhoid, purulent meningitis, influenza).

Treatment.—Locally, poultices or warm fomentations; and when pus has formed, its earliest possible evacuation—by preference from the conjunctival sac. The general constitutional treatment suitable to each case need not be discussed here.

Thrombosis of the Cavernous Sinus gives rise to symptoms which may be mistaken for those of an orbital process. The affection is described at p. 469.

Periostitis of the Orbit.—Acute periostitis has many symptoms in common with phlegmonous inflammation of the orbital connective tissue which generally accompanies it; but may usually

be distinguished from the latter inflammation occurring independently by the fact, as first pointed out by the late Mr. John Hamilton, of Dublin,[1] that in it pressure on the orbital margin is painful. The absence of this tenderness, however, is not always conclusive of the absence of periostitis, especially when the latter is restricted to the deep parts of the orbit. In periostitis, the eyelids are not usually so swollen as in inflammation of the orbital tissues. Suppuration may take place, necrosis in consequence of detachment of the periosteum may come on, and communications with the neighboring cavities may be formed.

In secondary syphilis, or in later stages of the disease, a syphilitic gumma of the orbital wall may form. This is accompanied by violent frontal neuralgia or headache, increasing at night. Proptosis (*πρό, forwards; πτῶσις, falling*), or protrusion of the eyeball occurs, with marked loss of motion in the eyeball in one or more directions. This loss of motion is a very characteristic symptom, and serves to assist in the diagnosis between this affection and other orbital tumors. It is probably due to an extension of the inflammation to the connective tissue of the orbit, and to the muscles themselves.

Very suggestive of gummatous periostitis of the orbit are: A rapidly increasing proptosis, with displacement of the globe downwards and forwards, and much loss of motion of the eye, and on palpation the sensation is given to the finger of a tumor in the roof of the orbit, where gummata most commonly are situated. Also, thickening of the upper margin of the orbit, with pain on pressure of the roof of the orbit, and radiating peri-orbital pain at night.

Again, periostitis of a chronic form, and without tendency to suppuration, occurs most commonly in persons with a constitutional rheumatic tendency. It is accompanied by pain in and about the orbit, and there is increased tenderness on pressure of the eyeball backwards. Exophthalmos, and all other outward signs, are here usually wanting.

The Prognosis depends much on the seat of the inflammation. If this be in the deep parts of the orbit, thickening of the periosteum may cause permanent protrusion of the eyeball; extension of the inflammation to the optic nerve may result in optic atrophy; the orbital muscles, or the nerves which supply them, may be implicated, with consequent paralysis; or, finally, the inflammation of the periosteum may strike into the meninges of the brain.

When the inflammation is near the margin of the orbit, early evacuation of pus, if it has formed, reduces the process within safe bounds; and this position is one of less danger in respect of its surroundings, than if the process be deep in the orbit.

Causes.—Periostitis of the orbit may be caused by blows or other traumata, by extension from neighboring cavities, by syphilis, or rheumatism.

Treatment.—Warm fomentations. Exit given to pus, if possible. Constitutional measures.

Caries of the Orbit is very frequently the result of periostitis, but often commences in the bone, and in either case is usually due to tubercular disease. It is also seen in very late syphilis. A trauma is sometimes the immediate cause of its onset.

It may attack any part of the orbital walls, its favorite seats being the margin above and to the outside, or below and to the outside. When it is seated deeply in the orbit, it often causes exophthalmos and pain. At the margin of the orbit it produces edema and swelling of the eyelids, with conjunctivitis; suppuration comes on, and the abscess finally opens through the integument or conjunctiva. A fistula is apt to remain for a length of time, and, the skin being drawn into this, ectropion of the lid is produced. If a portion of dead bone comes away, the resulting cicatrix is liable to maintain the ectropion (p. 228).

Treatment.—The evacuation of purulent collections at the earliest possible moment—if deep in the orbit, by the careful introduction of a long bistoury—the insertion of a drainage-tube, and the regular washing out of the cavity with antiseptic solutions, until no more rough or bare bone can be felt with the probe.

Injuries of the Orbit.—Wounds of the soft parts in the supra-orbital region, involving the supra-orbital nerve, are held by some to be capable of producing a reflex amaurosis (p. 554), and many such cases have been recorded under the name of supra-orbital amaurosis. By the light of modern physiology and ophthalmology it is not probable that any such reflex could take place. It seems more likely that the blindness in the cases recorded was brought about in some other way—*e. g.*, injury to the optic nerve in the optic foramen by the concussion, or by a fracture of the margin of the foramen, orbital periostitis, concomitant injury to the eyeball itself, facial erysipelas, intracranial lesions, and so on.

It may be, however, that a functional amblyopia, or amauro-

sis, similar to that occasionally seen, after long-continued blepharospasm (p. 136), has been present.

Perforating injuries of the orbit through the eyelids by prods of walking-canes, etc., and the lodgment of foreign bodies in the orbit are serious accidents. They are liable to be followed by phlegmonous inflammation; or, if a pointed weapon (stick, sword-cane, etc.) has been pushed into the orbit with some force, it may pass through the bony wall and perforate the brain, with fatal result.

It is remarkable what large foreign bodies may be concealed in the orbit. I once saw a case in which a bit of wood, 3-4 inch long by 1-2 inch wide, lay unsuspected in the orbit for many weeks, without causing any marked displacement of the eyeball.

Treatment.—Foreign bodies should be removed by dilatation of their wounds of entrance, or by the formation of a new passage through the conjunctival fornix—and great care should be taken to prevent the onset of inflammation, or to keep it within safe bounds.

Enophthalmos, or sinking of the eye back into the orbit, occurs to a certain extent in extreme emaciation, in Asiatic cholera, in paralysis of the sympathetic and in facial hemiatrophy, but it has been observed to an extreme degree as a result of injury. Beer [2] attributes it to atrophy of the retrobulbar cellular tissue; Lang [3] explains it by fracture or depression of a portion of the orbital wall; while Schapringer [4] refers the condition to paralysis of Müller's muscle from injury of the sympathetic nerve.

Orbital Tumors.—In the *Diagnosis of an Orbital Tumor* three questions present themselves:—First, Is a tumor of the orbit present? Secondly, Is the new growth confined to the orbit, or does it extend to neighboring cavities? and thirdly, Of what kind is the new growth? The diagnosis as regards any of these points does not often occasion much difficulty in advanced stages of the disease, especially where the growth occupies the anterior part of the orbit, or protrudes from it. It is rather in the early and middle stages that difficulties in diagnosis are apt to present themselves, and attention will here be mainly directed to those stages.

Exophthalmos is, of the signs by which the presence of a tumor is diagnosed in its early stages, by far the most important, because it is the most constant. In the earliest stages of

a growth which commences in the deepest part of the orbit there may be, it is true, no exophthalmos, while other symptoms—defects of sight, pain, loss of motion—may already be present; but when the growth attains to certain dimensions, or if in the anterior part of the orbit there be even a small tumor, the eyeball must be pushed out of its place.

An important diagnostic point in connection with the exophthalmos caused by a tumor—other than one within the muscular cone—is that its direction is almost always oblique and not straight forwards, for orbital tumors commonly tend to develop more along some one wall of the orbit than along the others, and hence the eyeball becomes pushed towards the opposite side as well as forwards. In cellulitis, edema of the orbital tissues, Graves' disease, and paralytic proptosis the exophthalmos has a direction straight forwards. Tumors growing from the apex of the orbit may, in their early stages, cause no obliquity of direction in the displacement of the globe, and some tumors do not do so even in an advanced stage of their growth; but these cases are exceptional. Tumors, too, situated altogether within the muscular cone, of which the most common are tumors of the optic nerve, need not cause any lateral displacement of the globe. Again, the exophthalmos caused by an orbital tumor usually increases in degree slowly and gradually, differing in this respect from exophthalmos due to most of the other causes, in which either a sudden or a rapid development of the exorbitism is the rule. While tumors are sometimes present in both orbits, especially lymphoma or lympho-sarcoma, yet it is infinitely more common for one orbit alone to be diseased; and hence monolateral exophthalmos is suggestive of orbital tumor.

Palpation in the Orbit often provides a valuable sign, should the new growth have come within reach in the anterior part of the cavity. In many cases, indeed, there is no difficulty whatever in recognizing the presence of an orbital tumor by this means, the sensation obtainable by the tip of the surgeon's finger pressed into the orbit being very definite; but in other cases the evidence is not so clear, and a reasonable doubt may exist as to whether any abnormal resistance is met with. By palpation we may gain some knowledge of the position, extent, shape, and consistence of the tumor, and whether it be adherent either to the walls of the orbit or to the eyeball. It is important, when practicable, to compare the result of examination of the dis-

eased orbit with the condition of the sound orbit; and this can be done to greater advantage, if palpation of the orbits be performed simultaneously with a finger of each hand.

The Röntgen Rays have been successfully employed in some cases for the diagnosis of retrobulbar growths.

Derangements of Vision are often, but by no means always, present in the early and middle stages of the growth of an orbital tumor. Their occurrence depends frequently on the rapidity of the growth of the tumor rather than upon its size. In an early stage of a rapidly increasing tumor the sudden stretching of, and pressure on the optic nerve may produce absolute blindness; while in another case, with an equal degree of exorbitism, but which has been brought on by a slowly growing tumor, vision may be unaffected by reason of the optic nerve becoming gradually accustomed to the change. Yet slowly growing tumors, which spring from the optic nerve or its neighborhood or from the deepest part of the orbit, are competent, by direct pressure on, or by implication of the optic nerve, to cause serious loss of sight, even in an early stage, and with but little exophthalmos. Optic neuritis, and, later on, optic atrophy, are occasionally discovered with the ophthalmoscope. Diplopia is often present when the globe is at first displaced, but disappears when the exophthalmos becomes extreme or the vision defective.

Pain is a symptom sometimes, but by no means always present in cases of orbital tumors. It is especially liable to be complained of when the growth is increasing rapidly in size, even though it may not have attained to great dimensions. The pain is then often of a neuralgic kind, and very severe, from the unaccustomed pressure on branches of the fifth nerve in the orbit.

Loss of Power of Motion of the Eyeball is a very common symptom in cases of orbital tumors. It is caused in some cases by the mechanical obstruction offered by the tumor, as a result of which motion of the eyeball becomes defective towards the side of the orbit on which the new growth is situated. In other cases the loss of motion is caused by stretching of the muscles from the exophthalmos, or by implication of them in the new growth, or by atrophy of their tissue, or by paralysis of the orbital nerves from pressure. When there is little or no loss of motion, while the exorbitism is marked, the conclusion may be drawn that the tumor lies within the muscular cone.

In every case the history, the rapidity of growth, the age and

general condition of the patient are important items for consideration.

Diagnosis of the Nature of an Orbital Tumor.—As regards the nature of the growth which may be present, it must be admitted that in many instances, in the early stages of a deeply seated tumor, we have to rest content with an indefinite or provisional diagnosis, unless an exploratory operation, with puncture or harpooning of the mass, is practicable; and such a procedure is often called for, in order to decide not only the nature of the tumor, but also its extent and origin.

Orbital Cysts.—Dermoid cysts are those most frequently found, and they are usually congenital. Indeed, if an orbital tumor be congenital, it is, as a rule, either a dermoid cyst or an encephalocele. Dermoid cysts, although usually congenital, do not often grow to any size until the age of puberty or later, and may then for the first time give rise to troublesome symptoms. They grow slowly, and finally reach very considerable size, and then bulge out between the eyeball and margin of the orbit. Pressure upon this protruding part causes it to diminish, while the exophthalmos is at the same time increased, and distinct fluctuation in the protruding part can be felt. The growth of the cyst is unaccompanied by pain or other inconvenience. The contents are generally either serous or honey-like, and occasionally hairs and other epidermic formations have been found in them.

Hydatid cysts also occur in the orbit, and several of these cases have been observed in England.

Treatment.—The cysts should be freely opened at the most prominent point, evacuated by gentle pressure backwards of the eyeball, and the sac syringed out two or three times daily with an antiseptic solution, until all discharge has ceased. The opening will then close, while the eyeball will already have returned to its place. If the contents of the cyst are solid, or nearly so, it becomes necessary to extirpate it *in toto*. To do this, Krönlein's operation (p. 515) must be resorted to. Or, a horizontal incision may be made along the orbital margin through the eyelid, in order that the cavity of the orbit may be reached, or two perpendicular incisions at either canthus through the upper lid may be made, and the latter turned upwards. With hooks or forceps, and scalpel or scissors, the cyst wall must then be carefully separated from all adhesions.

Exostoses occur as the result of inflammation of the bone and of periostitis, and also without any apparent cause, and are usually of the kind known as ivory exostoses. They spring most commonly from the ethmoid or from the frontal bone.

All the bony tumors give, of course, the sensation of dense hardness to the touch; but there are some malignant growths of such hardness that it may not be easy to tell them from the osteomata by palpation, and an exploratory puncture becomes necessary in order to decide the point. The growth of an orbital osteoma is excessively slow, in many instances commencing in infancy, and lasting into advanced life. In addition to the dense hardness of these tumors, the deciding points in the diagnosis are their usually globular and somewhat nodulated surface, and their immobility and direct connection with the walls of the orbit ascertainable by touch.

Operative interference in cases of exostosis of the orbit is only justifiable when the tumor does not grow from the roof of the orbit (as it then often involves the cranial cavity), and when there is reason to think it is attached to the orbital wall by a narrow base or pedicle. Several instances are on record in which the growth has become spontaneously separated by necrosis of its pedicle. Beyond destruction of the eyeball there is no danger associated with these tumors, even if their growth takes an intracranial direction; but they cause serious disfigurement and much pain.

Carcinoma and Sarcoma.—The first of these tumors takes its origin in some neighboring cavity, or from within the eyeball, and grows into the orbit; it never originates in the orbit. Sarcoma may originate in many different positions, most frequently, perhaps, in the periosteum and in the connective tissue about the lacrimal gland. These malignant tumors, after destruction of the eyeball by pressure, or by phthisis following ulceration of the cornea, attack the bony walls of the orbit and its surroundings.

The early extirpation of the tumor with complete evisceration of the orbital contents affords, in general, the only prospect, and that a slight one, of saving the patient's life.

Many forms of sarcoma, however, are non-malignant, especially those which lie free in the orbit and arise from the connective tissue. Indeed, Panas [5] held that many cases of sarcoma, as also of lymphadenoma of the orbit, are due to infectious

principles, toxins, or microbes, and are amenable to medical treatment by mercury, iodin, arsenic, or toxitherapy. So much certainly must be admitted—namely, that cases now and then present themselves, with all the signs and symptoms of orbital tumor, which ultimately undergo a purely spontaneous cure, or one unexpectedly brought about by iodid of potassium.

Symmetrical Tumors of the Orbits.—With the exception of tumors of the lacrimal glands, and possibly of rare instances of metastatic tumors, symmetrical tumors of the orbits are almost invariably lymphomata or lymphadenomata, occurring in leukemia or in pseudo-leukemia.

Pulsating Exophthalmos.—This title covers a great variety of vascular tumors, the majority of them having their origin within the cranium, while the remainder are truly orbital. Symptoms common to all these are: proptosis; the presence of peculiar bruits, which can be heard over the orbit, and usually, also, over a more or less extensive portion of the skull; and pulsation, apparent in the eyeball, or at some point of the orbital aperture. The last symptom may occasionally be absent during the whole, or part, of the progress of the case. The intracranial vascular tumors with which we are most likely to meet are: aneurysm of the ophthalmic artery at its point of origin from the internal carotid; aneurysm of the latter vessel; and, most commonly, arterio-venous aneurysm from communication of the internal carotid with the cavernous sinus—this latter of traumatic origin. In the orbit the following occur: True aneurysm of any of the arterial branches; diffused or circumscribed traumatic aneurysm; arterio-venous aneurysm, of traumatic origin; aneurysm per anastomosis; and telangiectic tumors.

Hemorrhage is liable to prove fatal in these cases.

Treatment.—Ligature of the common carotid affords the best prospect of cure. Digital compression of the same vessel has produced cure in some cases. Spontaneous cure has been observed occasionally in cases of arterio-venous aneurysm.

Implication of Neighboring Cavities.—As regards the question whether the tumor is confined to the orbit, or involves one or more of the neighboring cavities, it may be assumed that it is confined to the orbit, unless there are symptoms or signs which point in the opposite direction; and in each case these symptoms and signs ought to be looked for. Tumors may either originate in one of these spaces and grow into the orbit, which is the

more common event; or, originating in the orbit, they may at a later stage spread to a neighboring space; and it is often the history or progress of the case alone that can inform us which of these events has taken place.

Tumors which originate in the Frontal Sinus are usually either mucocele or exostosis. Mucocele of the frontal sinus frequently extends to the ethmoidal sinus and thence first encroaches on the orbit, pushing the eyeball downwards and outwards. Sometimes there is supra-orbital pain, and sometimes, when the nasal meatus has become involved, there is discharge from the nostril. The diagnosis in these cases is often obscure. Osteoma of the frontal sinus shows itself as a slowly growing and densely hard tumor almost free from pain, situated along the superior margin of the orbit, extending into the latter and pushing the eyeball downwards and forwards. It may subsequently extend to the orbital plate of the ethmoid. An error in diagnosis is, I think, liable to be made sometimes when a tumor of the frontal sinus drives the outer table downwards and forwards, and when the latter gives to the touch the sensation of a bony growth. If the tumor also involves the ethmoid cells, the lacrimal bone is apt to be similarly driven forwards, and the liability to the error I have mentioned is further increased. Bony growths originating in the orbit may invade the frontal sinus, and, whether originating there or in the sinus, are liable to produce absorption of the tables of the skull without any cerebral symptoms to indicate the occurrence.

Tumors of the Ethmoid Cells, which encroach upon the orbit, are likewise most commonly either mucocele or osteoma. Mucocele of the ethmoid cells presents itself in the orbit as a tumor, gradually increasing in size, on the inner wall of the orbit, and pushing the eyeball outwards and forwards. When it has grown sufficiently large, palpation of it will discover fluctuation. The source of error just referred to, when the lacrimal bone is pushed in front of a slowly growing tumor of the ethmoid cells, must be borne in mind. The sharp posterior edge of the lacrimal bone is easily felt for and found, and will direct the diagnosis into the proper channel. Mucocele of the ethmoid cells encroaching on the orbit must also be distinguished from a dermoid cyst, but to this I shall return later on. Osteoma of the ethmoid appears in the orbit as a hard round swelling at the inner canthus, followed by a swelling of the cheek and displace-

ment of the eye outwards and forwards. It is apt also to extend into the nasal meatus, driving the septum out of place, and pushing the hard palate downwards, so that examinations of the nose and of the mouth should be made in aid of the diagnosis. Enchondromata and fibromata also sometimes spring from the ethmoid, and extend into the orbit, and malignant growths may be met with here.

Tumors that spring from the Body of the Sphenoid Bone, or from the Antrum of the Sphenoid, and encroach upon the orbit are rare, and the diagnosis of their origin in an early stage is practically impossible. Here, again, the examination of the naso-pharynx is important. It is stated (Stedman Bull) that an orbital tumor which soon causes blindness, commencing in the temporal side of the field, and leaving the fixation point unaffected to the last, while at the same time a growth appears in naso-pharynx, is likely to be one having its origin in the sphenoid antrum. Bony tumors—osteoma, hyperostosis and extostosis—polypi, and sarcomata, are the growths most frequently found to originate in the sphenoid antrum.

Tumors of the Maxillary Antrum sometimes erode the floor of the orbit, and grow into that cavity, driving the eyeball upwards and inwards, or upwards and outwards. The breadth of the cheek is increased, the nose becomes pushed towards the opposite side, and the roof of the mouth is pushed downwards. Tumors of the antrum of Highmore sometimes cause pain in the teeth, or in the region of distribution of the infra-orbital nerve, and there may be a dull pain in the region of the antrum. In some cases there is a discharge of pus or of blood from the nostril.

Intracranial Tumors do not often invade the orbit, and then it is tumors of the middle fossa which gain access through the sphenoid fissure and optic foramen. The diagnosis of the origin of the disease can only be made if cerebral symptoms have existed prior to any sign of a new growth in the orbit. Tumors of the pituitary body may encroach upon the orbit by way of the sphenoid fissure, and are apt to be associated with polyuria and bitemporal hemianopsia, which serve to aid the diagnosis.

A more common event, although not in an early stage of the growth, is the extension of a primary orbital tumor to the brain either along the optic nerve, through the sphenoid fissure, or through the roof of the orbit by erosion of the bone. This oc-

currence is usually evidenced by the presence of cerebral symptoms; but cases have been met with where no such symptoms existed, although the orbital growth had encroached upon the anterior or middle fossa of the skull.

Tumors of the Lacrimal Gland.— Slowly increasing exophthalmos, the eyeball being gradually pushed forwards and inwards, and its motions curtailed in the upward and outward direction, is a constant symptom here. In the region of the gland the upper eyelid seems to be swollen; but palpation shows this to be caused by a growth situated behind the lid, and not in it, and, further, that the tumor originates in the orbit. The upper fornix of the conjunctiva is found, on eversion of the upper lid, to be pushed downwards. After a time the blood-vessels of the upper lid become congested and tortuous, and when the tumor has grown very large the eyelids cannot be closed, the eyeball becomes injected, and the cornea dry and opaque.

Adenoma, or Adeno-sarcoma, and Fibro-adenoma are the most common forms of tumor of the lacrimal gland, and the gland on each side may be affected.

Extirpation of the growth at as early a stage as possible is indicated. The tumor is reached, either through an incision made through the lid parallel to the outer half of the upper orbital margin; or, the external commissure having been divided, and the upper lid turned up, the growth can be removed through an incision made in the conjunctival fornix; or Krönlein's operation (p. 515) may be indicated.

Tumors of the Optic Nerve. (See p. 436.)

Hernia Cerebri, either in the form of meningocele or of encephalocele, may invade the orbit. Its most common situation is the upper and inner angle of the orbit, to which it gains access through the suture between the frontal and ethmoid bones. It appears as a fluctuating, often transparent, pulsating congenital tumor. Pressure upon it causes it to disappear, but gives rise, at the same time, to symptoms of cerebral irritation, or pressure.

A congenital tumor in the upper inner angle of the orbit must always be regarded with suspicion, lest it be a cerebral hernia, even though it do not pulsate, or on pressure cause cerebral symptoms. In the large cerebral hernia death in the first few days of life is, we know, the rule.

Shrinking of the Conjunctiva (Xerophthalmos) and of the

Subconjunctival Tissue of the Orbit, Subsequent to Enucleation of the Eyeball.—In some cases where the eyeball has been excised, and in due course a prothesis fitted, the conjunctiva and subconjunctival tissues shrink to such a degree, after some months or years, as to reduce the size of the orbital cavity so that the wearing of a glass eye becomes impossible. This is especially liable to occur amongst those hospital patients who are careless in removing the prothesis at night, and in keeping the socket thoroughly clean at all times. The attempt is then often made to restore the orbital cavity, so as to render it possible to wear at least a small glass eye, by means of skin grafts or of mucous membrane grafts after the method either of Thiersch or of Wolf. The success attendant on these procedures is usually a very moder-




Fig. 159.

ate one, and often not permanent, owing to subsequent renewed shrinking of the subconjunctival tissue.

In these cases the lower sulcus is the most important part of the cavity, and if it can be made sufficiently deep, a small artificial eye will be retained. With this object in view, Mr. P. W. Maxwell, of Dublin, has devised the following operation:

An incision is made in the floor of the socket, and carried downwards behind the lower lid. A semilunar flap, about 8 mm. in width at its widest part, is marked out on the skin of the lid, its upper concave border being about 5 mm. below the palpebral margin. The incision along the upper border of the flap is made to communicate with the bottom of the wound in the socket. The flap is now dissected up from the subcutaneous tissue, except an area represented by the dotted line in Fig. 159. The two ends of the flap (a' and b') are passed through the

opening into the socket, and sutured to each end of the socket incision (*a* and *b*); and the borders *A'* and *B'*, being also passed through, are sutured to *A* and *B* respectively. The space on the cheek is closed, and the operation completed by inserting into the socket a temporary glass eye or shell. This should be, as nearly as possible, of the size and shape of the eye to be ultimately worn; it prevents the new sulcus from being obliterated by contraction, and gives it a suitable shape. It cannot safely be taken out for at least a week, as the skin incision might perhaps be opened in so doing. If there be secretion, the space behind may be flushed out by a lacrimal syringe armed with a fine curved nozzle, which can be introduced under the edge of the eye at the inner or outer canthus. A glass shell with a hole in front is preferable to a glass eye, for it allows a syringe to be more easily used, and, being transparent, a view of the parts behind can be obtained.

To obtain a good result the following points should be attended to:

1. Make the incision in the socket as long as the space will permit, and see that this length is maintained throughout its entire depth.

2. Make the skin flap considerably longer than the incision in the socket.

3. When dissecting up the skin flap, leave undisturbed a portion (dotted line in figure) equal in length to the socket incision. This subsequently forms the fornix, or sulcus. If a shorter portion is left, the sulcus is apt to become V-shaped, which would require a specially made glass eye.

4. When closing the space on the cheek, as the lower border is longer than the upper, great care should be taken to equably distribute the excess, so as to avoid puckering. When this has been neatly done, the line upon the face becomes quite invisible after a few months.

In none of the cases, so far, has it been necessary to make a sulcus above. The same operation could, however, be performed on the upper lid, provided that, after dissecting up both the borders of the skin flap, the tendon of the levator were secured with one or two sutures before dividing it. After the skin flap is in its new position, the cut end of the levator could be attached to the tarsus. In closing the skin wound, the ends of these deep sutures should be allowed to project outwards, so that they may be pulled out

when they ultimately become loose. If it were possible to obtain really aseptic catgut, the ends of the sutures might be cut short and buried.

In addition to providing a sulcus, the operation adds half the width of the flap—viz., 4 mm.—to the vertical diameter of the socket.

Temporary Resection of the Outer Wall of the Orbit (Krönlein's Operation).—This operation was devised by Professor Krönlein,[6] of Zurich, for the removal of tumors of the optic nerve, and other new growths and cysts in the posterior part of the orbit, as well as foreign bodies, without sacrificing the eyeball, or perhaps even the sight. It may also be employed to reach purulent foci in the orbit, and has been used, with what result I cannot say, to remove some of the retrobulbar fat in cases of exophthalmic goiter.

The eyebrow and the scalp in the temporal region are shaved, and the skin of the whole region of the operation is rendered aseptic.

The First Stage of the operation consists in making a curved incision on the temple through the skin and soft parts. This incision commences on the temporal ridge, at a point where the latter would be intersected by a horizontal line running 1 cm. above the supra-orbital margin. The middle point, or apex, of the incision lies in the center of a horizontal line, which unites the external canthus with the outer orbital margin. The end of the incision lies on the zygoma, in the center of a horizontal line uniting the external canthus with the tragus. The length of the incision in adults is 6 to 7 cm., and the direct distance between its two ends is about 5 cm. Smaller incisions are inconvenient. In that portion of the incision which runs along the margin of the orbit it goes to the bone, through the periosteum.

The Second Stage consists in raising the periosteum from the inner surface of the outer wall of the orbit with a slightly curved and somewhat pointed elevator, which is introduced at the exposed outer orbital margin. The periosteum is separated upwards as far as 1 cm. above the fronto-malar suture, downwards as far as the spheno-maxillary fissure, and posteriorly until well behind the spheno-zygomatic suture. This proceeding is not difficult, as the periosteum is closely adherent along the orbital margin only, and at the sutures. The point of the elevator is now passed directly downwards, and carefully introduced into the spheno-

maxillary fissure a few millimeters behind the spheno-zygomatic suture. The handle of the instrument is then turned over gently towards the nose, thus pressing the periosteum and all the contents of the orbit somewhat inwards, and exposing the bared inner surface of the outer orbital wall. The object of passing the point of the elevator into the spheno-maxillary fissure—where it remains during the next stage of the operation—is to fix the point towards which the osseous incisions are to be made to converge. Some surgeons prefer to omit this act, and the proximity of the infra-orbital nerve, and of the infra-orbital vessels must be borne in mind.

The Third Stage is the resection of the bony wall by three incisions, two horizontal and one oblique. The upper horizontal bony incision is made with a thin, sharp chisel, and divides the external angular process of the frontal bone close to its base. The soft parts have been previously drawn aside, the periosteum over the seat of the proposed bony incision divided and, along with the lacrimal gland, drawn aside.

The oblique bony incision passes from the deepest part of the previous incision downwards and backwards behind the spheno-maxillary suture, through the greater wing of the sphenoid bone, to a point about 1 cm. behind the anterior end of the spheno-maxillary fissure, where the point of the elevator has been kept all through.

The lower horizontal bony incision divides the frontal process of the malar bone close to its base, the soft parts having been drawn aside, and the periosteum divided. The incision ends at the anterior extremity of the spheno-maxillary fissure.

In making the bony incisions there is the danger of splintering to be contended with, and in the oblique incision there is some danger of luxating the spheno-maxillary suture. The chisel must be very sharp and thin, and it is well to apply its corner rather than its full edge to the bone, while only light taps with the mallet are used. It is important to make the bony incision in the above order; or, at any rate, the oblique incision should not be the last to be made, for, if it be, the thin outer wall of the orbit is liable to become severely splintered during the chiseling of the second bony process.

The Fourth Stage is the turning backwards of the flap of bone and soft parts, and the exposure of the interior of the orbit. It is necessary, after the flap has been turned well back, to divide the

separated periosteum with a blunt-pointed scissors from before backwards. It is sometimes necessary, in order to reach the focus of disease, to divide the tendon of the external rectus near its sclerotic insertion, and possibly other orbital muscles must be severed; but this should be avoided, if possible.

When all manipulations required in the orbit have been completed, any muscles which may have been divided are sutured to their insertions, the periosteum is replaced in its normal position, the flap of bone and soft parts turned forwards into its place, and secured there by a few catgut sutures through the periosteum. A drain is then placed in the lower part of the wound, and the rest of the wound is accurately closed with fine silk sutures, and an aseptic dressing and bandage applied. The catgut sutures through the periosteum, and the drain, are regarded by several operators as unnecessary. The only recorded case of this operation in the United Kingdom is one successfully performed by Dr. Louis Werner [7] for a tumor of the optic nerve. (See also p. 436.)

Exophthalmic Goiter (Graves' Disease, Basedow's Disease).

Symptoms.—The three cardinal symptoms of this disease are: increased rapidity of the heart's action, which may reach two hundred beats per minute; tumefaction of the thyroid gland; and exophthalmos. Of these the cardiac symptom is the most constant, and usually the first to appear; either, or both of the others, may be wanting. There is often also great emaciation, with outbursts of sweating and diarrhea. A venous murmur may be heard in the neck; and in females there is very commonly irregularity or suppression of menstruation.

The disease has been observed at all ages, but is most common in early adult life.

Von Graefe's Sign is a very early, tolerably constant, and almost pathognomonic one; it consists in an impairment of the consensual movement of the upper lid in association with the eyeball. When, in the normal condition, the globe is rolled downwards, the upper eyelid falls, and thus its margin is kept throughout in a constant relation to the upper margin of the cornea. In Graves' Disease the descent of the upper lid does not take place, or only in an imperfect manner; and, consequently, when the patient looks down, a zone of sclerotic becomes visible between the margin of the lid and the cornea. This symptom is often present prior to any exophthalmos, and hence its great diagnostic

value. It may also continue after the latter disappears,—although it is perhaps more common for it to disappear before the proptosis, —and it is not seen, or but very rarely so, in protrusion of the globe from other causes. But the sign is not so absolutely pathognomonic as it was held by von Graefe to be; for it may be absent in Graves' Disease, although very rarely so, in the early stages, and it is sometimes present in other diseased states, and even in health.

Stellwag's Sign is also very constant. It is incompleteness and diminished frequency of the act of involuntary nictitation.

This act occurs sometimes only once in a minute; or several rapid nictitations take place, and then a lengthened pause. The nictitation each time is incomplete, the margins of the lid not being brought together. The result may be that the lower third of the cornea becomes covered with pannus vessels, owing to the constant exposure; for even during sleep the eyelids remain partially open.

Dalrymple's Sign consists in an abnormal widening of the palpebral aperture, due to retraction of the upper eyelid. It is this gaping of the eyelids which gives the characteristic staring aspect to the patient. This sign is often erroneously attributed to Stellwag, or is included in his sign. The error is due to the fact that in the same paper [8] in which Stellwag first drew attention to what is above described as his sign, he discussed this other previously observed sign. According to White Cooper [9] it was Dalrymple who first pointed out the latter.*

Probably each of these "signs" is due to the one cause suggested by Sharkey [10]—namely, loss of power in the orbicularis rather than overaction of the levator.

Otto Becker stated that in a majority of the cases spontaneous pulsation may be seen in the retinal arteries, but I have only found it sometimes. The vision—unless when corneal complications supervene—and condition of the pupil are unaffected by the disease. In some cases there is an increased flow of tears, but most of the patients complain of a dryness of the eyeballs. The sensibility of the cornea is diminished. Ulcers of the cornea are

* Other conditions which produce widening of the palpebral aperture or "Staring Eye," are: (1) Orbital Tumor (mechanically). (2) Stimulation of the Cervical Sympathetic. (3) Cocain (in slight degree, probably by reason of 2.—Jessop). (4) Women after child-birth (hysteria). (5) In tetanus (spasm of occipito-frontalis). (6) In complete amaurosis.

not common, but are said (von Graefe) to be more frequent in men than in women, although Graves' Disease is more common in women. The exposure of the eye and dryness of the cornea are the chief causes of ulceration, when it occurs; but Sattler inclines to the belief that it is also largely due to paralysis of the nervous supply of the cornea.

The patients are often hysterical; and even marked psychical disturbances have been noted, such as a peculiar and unnatural gayety, rapidity of speech, and great irritability; or, on the other hand, extreme depression, and even attempts at suicide have been observed; also loss of memory and inability to make a mental effort. The motions of the eyeball have in some cases been defective—a fact for which the exophthalmos does not account. Trousseau's Cerebral Macula is often well marked.

The Progress of the Disease is, as a rule, very chronic, extending over months or years, but liable to fluctuations in the intensity of its symptoms. A few cases have been recorded in which it became fully developed in the course of some hours or days. After a lengthened period, and many fluctuations, the symptoms usually slowly disappear. Occasionally a slight permanent swelling of the thyroid may remain, and very often more or less exophthalmos. About 12 per cent. of the cases go from bad to worse, and end fatally from general exhaustion, organic disease of the heart which may have come on, cerebral apoplexy, hemorrhage from the bowels, or gangrene of the extremities.

Causes.—Anemia and chlorosis are general conditions very often present, as are, also, irregularities of menstruation; but it is probable that the latter should be regarded rather as a concomitant symptom than as a cause. Severe illnesses are recorded as having gone before the onset in many cases, and also excessive bodily or mental efforts. Great sexual excitement has been known to be followed by Graves' Disease, and depressing psychical causes are not unfrequent forerunners of it. In many instances, however, the patients have been perfectly healthy, and no cause could be assigned.

The Enlargement of the Thyroid is due in the first instance to dilatation of its vessels; but in a late stage hypertrophy of the gland tissue may be produced, and increase of its connective tissue, and even cystic degeneration. The Exophthalmos is due to hyperemia of the retrobulbar orbital tissues, as is demonstrated by a vascular bruit often present, and the fact that steady pres-

sure on the globe diminishes the protrusion. Hypertrophy of the orbital fat may be found *postmortem,* but is, doubtless, secondary to the hyperemia.

With regard to the nature of the disease, very many theories have, from time to time, been put forward, but none is quite satisfactory, and a discussion of them does not come within the scope of this work.

Treatment.—A principal part of this consists in the careful regulation of the patient's general health and functions. Freedom from mental anxiety and excitement, regular hours, moderate exercise, and change of air are the most important items.

The fluctuations, which occur in the intensity of the symptoms, render it difficult to arrive at definite conclusions with regard to the efficacy of remedies, a vast number of which have been tried and lauded from time to time. In mild forms of the affection, and especially if the anemia be well marked, iron internally is beneficial, but in severe cases it has the opposite effect. Quinin in moderate doses has been employed with benefit in some cases. Trousseau recommended digitalis in large doses, but its effect must be watched. The beneficial action of iodid of potassium in ordinary goiter has suggested its use in this disease; but under its influence the symptoms are sometimes aggravated, and it is doubtful whether they are ever relieved by it. Hulke praised aconite highly, and Sir Samuel Wilks has no doubt as to the value of belladonna. Ergotin internally has been tried, and with advantage in some instances. Sattler warmly recommends a well-regulated hydropathic treatment, when the patient is not too excitable. Paroxysms of cardiac palpitations, etc., are best combated by ice applied to the head, heart, and goiter. The sympathetic theory has induced the trial of a galvanic treatment of the cervical sympathetic.

Gauthier recommends antipyrin before everything else. Extract of the thymus gland has been occasionally employed, and with encouraging results.

Extirpation of the thyroid has been performed in recent years with success in some cases.

The great number of remedies which have been proposed for it demonstrate the intractable nature of this disease. Yet a considerable proportion of the cases do undergo cure, in so far as quieting of the heart's action, and reduction, or possibly sometimes complete disappearance, of the goiter and exophthalmos, are

concerned. It is common, however, even in the best recoveries, to see some exophthalmos remain permanently.

In cases where the exophthalmos is so great that the cornea is exposed even during sleep, it is desirable to perform tarsorrhaphy (p. 207), and the same operation is indicated when, the disease having subsided, the exophthalmos still remains to a degree which gives the patient a disagreeable expression.

References.

[1] "Dublin Journal of Medical Sciences," 1836.
[2] "Archives of Ophthal.," xxii. p. 98.
[3] "Trans. Ophthal. Soc., U. K.," vol. ix. p. 41.
[4] "Klin. Monatsbl. f. Augenheilk.," September, 1893.
[5] "Brit. Med. Journal," October 19, 1895.
[6] "Beiträge zur Klinischen Chirurgie," iv. 1; see also Domela-Nieuwenhuis, "Beiträge zur Klinischen Chirurgie," xxvii. 2. p. 558.
[7] "Trans. Ophthal. Soc., U. K.," vol. xxiii.
[8] "Wiener Med. Jahrbücher," xvii. p. 25, 1869. See also "Klin. Monatsbl. für Augenheilkunde," 1869, p. 216; and "v. Graefe und Saemisch's Handbuch," vi. pp. 955 and 956.
[9] The "Lancet," May 26, 1849, p. 553.
[10] "Trans. Ophth, Soc., U. K.," vol. xi. p. 204.

CHAPTER XIX.

Part I.—Ocular Diseases and Symptoms liable to accompany Focal Disease of the Brain.

Part II.—Ocular Diseases and Symptoms liable to accompany Diffuse Organic Disease of the Brain.

Part III.—Ocular Diseases and Symptoms liable to accompany Diseases and Injuries of the Spinal Cord.

Part IV.—Nervous Amblyopia, or Asthenopia.

Part V.—Various Forms of Amblyopia.

Part I.

OCULAR DISEASES AND SYMPTOMS LIABLE TO ACCOMPANY FOCAL DISEASE OF THE BRAIN.

Hemianopsia (*ἥμισυς, half; ἀ, priv.; ὤψ, the eye*). This term implies a loss of sight in one-half of the field of vision, usually of each eye, consequent upon a lesion at the center of vision, at the chiasma, or at some point in the course of the visual fibers between these two places. It is not used for cases in which one-half of the field is lost, owing to disease within the eye itself.

The line dividing the seeing from the blind half passes vertically down the center of the field, as in Fig. 160. Sometimes this line lies a little to one side of the center of the field, so as to admit of the latter being included in the seeing part, as in Fig. 161; and sometimes—although in other respects the dividing line lies in the center of the field—the fixation point is circumvented by it, so as to leave that point free, as in Fig. 162; and probably this is the most common arrangement. This subject will be further discussed later on. Again, although rarely, the dividing line may have an oblique direction, as in Fig. 163. It is probable that such a field as Fig. 163 is due to some peculiar arrangement in

the decussation of the nerve fibers in the individual case. Furthermore, cases occur which are properly regarded as hemianopsia, and yet in which only a sector of one side of the field is wanting, as in Fig. 164. Figs. 160, 161, 162, and 163 would be called complete hemianopsia, while Fig. 164 would be termed incomplete or partial hemianopsia. Finally, if all three visual per-




Fig. 160.*




Fig. 162.




Fig. 161.

ceptions be lost, the hemianopsia is called absolute; but if only one (color) or two (color and form) be wanting in the defective part of the field, it is termed relative hemianopsia. The vast majority of cases of hemianopsia are absolute.

Homonymous Hemianopsia is the most frequent form. In it the corresponding half—the right half or the left half—of the field of each eye is wanting, as in Fig. 165, in which the left side




Fig. 163.




Fig. 164.

of the fields, from the patient's point of view, is blind, implying a loss of function in the right half of each retina.

Temporal Hemianopsia is loss of vision in the outer side of each field, in consequence of loss of power in the median half of each retina (Fig. 166). It is by no means so common as the homonymous form.


Superior or Inferior Hemianopsia, also called Altitudinal Hemianopsia, in which the upper or lower half of the field is blind, is very rare; and it is doubtful whether Nasal Hemianopsia has

* Figs. 160, 161, 162, 163, and 164 are diagrammatic representations of the left field of vision.

really been observed, although it has been described. In the latter the inner side of the field of one eye only is lost, owing to defective function of the temporal side of the retina.

It will be convenient here to set forth the prevailing views as to

The Arrangement of the Cortical Visual Centers, their relations



Left Field. Right Field.

Fig. 165.

to the Retina, and the Course of the Optic Fibers between these two points.

Pathological anatomy leaves little doubt but that in man the visual center is situated on the mesial surface of the occipital lobe, rather than in the angular gyrus or elsewhere; and the evidence goes to show that the absolute optical center chiefly occupies the



Left Field. Right Field.

Fig. 166.

cortex of the cuneus and of the superior occipito-temporal convolution.

Henschen,[1] as the result of clinico-pathological investigations, believes it to be situated in the middle part of the calcarine fissure, which lies between these structures; and that the upper, or cuneic, lip represents the homonymous dorsal retinal quadrants; while the lower, or lingual, lip represents the homonymous ventral quadrants of the retina. Vialet,[2] on the other hand, thinks that the visual center embraces all the mesial surface of the occipital lobe included between the occipito-parietal fissure and the lower border of the third occipital convolution, and that it extends above and behind

as far as the free border of the hemisphere. The calcarine fissure he also regards as of great importance *; and I think, indeed, he rather proves than disproves Henschen's view of its midddle third being the actual cortical center for vision.

It is universally recognized that the nerve fibers from the homonymous half of each retina, *e. g.*, from the temporal half of the right retina and from the median half of the left retina, pass wholly through the corresponding optic tract,—in this case the right tract,—to the corresponding cortical center for vision (Fig. 167).

A case published by Hun,[3] in which the left lower quadrant in each field was blind, and where the autopsy showed a lesion (atrophy) strictly limited to the lowed half of the right cuneus, renders it probable that there is in man a correlation between parts of the retina and of the occipital lobe, as Munk had already proved to be the case in dogs, and that the optic fibers from the right lower quadrant of each retina terminate in the adjacent part of the right superior occipito-temporal convolution, the left halves of the retina and left optic centers being of course similarly correlated. If this view be correct, as seems probable from Henschen's investigations, it is evident that altitudinal hemianopsia can hardly occur as the result of a central lesion, as nothing short of disease confined to the lower half of each cuneus would produce it.

It is now generally believed that relative hemianopsia (*e. g.*, color hemianopsia alone) is the result of a lesion of less intensity that which causes absolute hemianopsia. Cases of apparently pure hemiachromatopsia may, with careful tests, show some diminution of the form-sense in the half fields which are defective for color-sensations. Non-cortical lesions, even at the chiasma, may also give rise to hemiachromatopsia. Thus it would seem that the color-sense is more easily affected by disease than the form or light senses, and that, too, irrespective of the position of the lesion in the visual path.

It is now generally conceded that the macula lutea is specially represented in the cortical center. But there are at least two very distinct views as to the arrangement of these macular centers,

* One of the most important cases which has been published in connection with this question is that of Dejerine and Vialet (Société de Biologie, Paris, December, 1893), in which both eyes became suddenly blind, without loss of consciousness or other symptoms. The patient lived for a short time, and after death from pneumonia the post-mortem showed lesions of the structures bordering the calcarine fissure on both sides.




FIG. 167.—EXPLANATION OF FIG. 167.

FIG. 167.—Diagram of Course of Optic Fibers, with the Cortical Centers and Relations to Fields of Vision, illustrating one theory of the Macular Supply; according to which the macula is supplied on the same plan as the rest of the retina—*i. e.*, each side of it from the corresponding side of the brain.

R. F., Right field of vision; *L. F.*, Left field of vision; *R. E.*, Right eye (retina); *L. E.*, Left eye (retina); *m. l.* and *m. l.*, Macula lutea; *O. N.* and *O. N.*, Optic nerves; *Ch.*, Chiasma; *Tr.* and *Tr.*, Optic tracts;

R. C. C. and *L. C. C.*, Right and left cortical centers; *M.* and *M.*, Macular fibers; *P.* and *P.*, Peripheral fibers.

1. Lesion of right cortical center = left homonymous hemianopsia, the line of demarcation passing round the left side of the fixation point in cases of embolism and thrombosis, but through the fixation point in cases of hemorrhage (*vide infra*).
2. Lesion of the right optic tract = left hemianopsia, the line of demarcation passing through the fixation point.
3. Lesion of the chiasma = bitemporal hemianopsia, the line of demarcation passing through the fixation point.
4. Lesion involving fasciculus lateralis only to right eye, causing nasal hemianopsia in the right field.

Diagram 167 also illustrates the fact that, as regards its relation to the optic tracts, the field of each eye is divided unequally, and not in halves—*e. g.*, the right tract governs about one-third of the field of the right eye, while the other two-thirds is governed by the left optic tract.

and as to the course of the macular fibers. These different views have been called into existence by the desire to explain the fact that in hemianopsia the line of demarcation sometimes passes through the fixation point in the field, and sometimes leaves it in the seeing half. It seems to me that neither of these theories is quite satisfactory, and I regret that I cannot offer one that is more so.

According to one theory, illustrated by Fig. 167, the macular region of the retina is invariably supplied on the same plan as the rest of the retina—*i. e.*, each side of it from the corresponding side of the brain. In order, then, to explain why it is that in some cortical lesions the line of demarcation passes through the fixation point in the field, while in others it deviates towards the blind side, the supporters of this view state that the cortical center for the macular region is more richly supplied with blood-vessels than the rest of the visual center; as is the macula lutea itself in relation to the rest of the retina. Hence, when the lesion is an embolism, or thrombosis, of the vessels supplying that part of the brain, this special region, by reason of abundant anastomoses, preserves its functions, and then fields as in Fig. 165 are produced. But, if the lesion be a hemorrhage, the macular region of the cortex would be apt to be involved in the lesion with the rest of the visual center, and loss of function in the corresponding half of the macula lutea, with the line of demarcation passing through the fixation point, results.

According to the other theory, the whole of the macular region —and in some instances even more than this—of each retina being innervated from each hemisphere, there is an overlapping, as it is called, of nervous supply to these retinal regions. Consequently, if there be a lesion at the center for vision in one occipital lobe, the center for vision in the other occipital lobe being sound, the functions of the whole of each macula—or even of more than this—of the defective side of each retina will be preserved. Cases where, occasionally, in cortical lesions, the line of demarcation in the field does go through the fixation point would be accounted for according to this theory, by an individual variation in the supply of the maculæ, which in these instances would be similar to that of the remainder of the retinæ.

But any such theory, to be satisfactory, must be capable of explaining the phenomenon in question, not only when the lesion is in the cortex, but also when the hemianopsia is caused by a lesion in the tract or chiasma. Yet an examination of Fig. 167 will show that, according to the theory it represents, in lesions of the tract (2), or of the chiasma (3), the line of demarcation would pass through the fixation point. And, according to the other theory, a lesion either at the tract or at the chiasma would always cause the dividing line to circumvent the fixation point. It happens, however, that with lesions at either of these situations the dividing line sometimes passes through the fixation point, and sometimes to one side of it. Consequently, I do not think we have yet solved the problem of the nervous supply of the macula lutea.

Some ophthalmologists hold that the line of demarcation always passes through the fixation point, and that it is merely imperfect fixation on the part of the patient which makes it seem to pass round it. This I believe to be an erroneous view; but there are no doubt cases in which it is difficult to determine the question, and where the line of demarcation approaches very close to the fixation point.

The Localization of the Lesion in Cases of Hemianopsia is a subject of interest, and, in view of the advances made within recent years in cerebral surgery, it is of great practical importance.

Lesions of the center of the *Chiasma* injuring the crossed fibers produce as their characteristic symptom bitemporal hemianopsia, which may be relative at first, beginning, for instance, as a hemi-

achromatopsia, but later on becoming absolute. In some cases (basal meningitis, periostitis, hyperostosis) the diseased process comes to a standstill, and the bitemporal hemianopsia remains. But the disease generally extends to the uncrossed fibers, ultimately causing complete blindness. Even when the disease is non-progressive, central vision is impaired; whereas in homonymous hemianopsia the latter is not always affected. Optic atrophy, often commencing on the inner side of the papilla, is nearly always present at some period of the disease. Syphilitic gummata may cause transient recurrent attacks of bitemporal hemianopsia.

In Altitudinal Hemianopsia the lesion must also, as a rule, be at the chiasma, encroaching on it from above or below. Symmetrical cortical lesions might, and optic neuritis sometimes does, produce it.

In Nasal Hemianopsia, too, the lesion must be at the chiasma, and must be so situated in its outer angle as to involve only the fasciculus lateralis of the affected eye. The occurrence of binocular nasal hemianopsia is evidently almost impossible, implying, as it does, symmetrical lesion of the fasciculus lateralis of each tract. According to Henschen, a tumor in the external angle of the chiasma is apt to affect the crossed fibers as well as the uncrossed, and to produce a form of bilateral homonymous hemianopsia. Other symptoms which may be present in lesions of the chiasma are anosmia, paralysis of orbital nerves, and anesthesia of the conjunctiva and cornea. The causes are: fractures of the body of the sphenoid, cysts, tubercle, tumors, exostoses, distention of the infundibulum of the third ventricle, and, most frequently, tumors of the pituitary body. In the latter case proptosis, discharge of fluid from the nostril, and diabetes may be present.

Bitemporal Hemianopsia is a very common and early symptom in *Acromegaly,* a disease characterized by great hypertrophy of the face and extremities, associated with enlargement of the pituitary body, and other conditions which are not so constant.

In Homonymous Hemianopsia—the commonest form of the symptom—localization of the lesion is a more difficult matter than in any of the other forms; for here the disease cannot be situated at the chiasma, but may be in the optic tract, or in the visual center, or anywhere in the lengthened course of the fibers which connect these two parts.

Can we distinguish a complete and absolute hemianopsia, due to a lesion confined to the occipital lobe, from a similar defect in the field, due to a lesion in the optic radiations, internal capsule, pulvinar, or optic tract? We may conclude that the hemianopsia depends upon an occipital lesion, if it be unaccompanied by hemiplegia, motor aphasia, or paralysis of cerebral nerves, as direct symptoms; but be it remembered that one and all of these are liable to accompany lesions of the occipital lobe as distant * symptoms.

Aphasia, too, occasionally accompanies right cortical hemianopsia (*i. e.,* due to a lesion in the left occipital lobe), although it is not easy to offer a satisfactory explanation of the fact.

A diagnostic symptom is what is known as negative vision, "vision nulle"—that is to say, the patient, though he may be aware of the loss of half of his visual field, has no sensation of darkness in it, and is just as unconscious of the defect as a healthy person is of his blind spot.

Cortical hemianopsia may be a distant symptom. Gowers has observed that, at the onset of many attacks of cerebral hemorrhage, hemianopsia is present as a distant symptom of very fleeting character—so transitory, indeed, that it does not complicate attempts at localization; but I have seen it to last as long as three weeks. Except under this condition, distant hemianopsia seems to be rare—a fact which enhances the localizing value of the symptom.

Cortical hemianopsia may be incomplete, inasmuch as the homonymous quadrant only of each field may be wanting. The explanation of this was given when discussing (p. 524) the correlation of the visual cortical centers to parts of the retina.

So much for absolute hemianopsia. But the lesion may be such as to destroy only the color-sense, without reaching those for form and light. Eight cases of hemiachromatopsia are on record.

* I suggest the term "distant symptom" in preference to those in common use—namely, "indirect symptom" and "pressure symptom." We cannot assume that these symptoms are less the direct result of the lesion than any of the others which are present; and, in many instances at least, it is certain that they cannot be due to pressure. In short, we do not yet know what produces these symptoms,—they may be caused by inhibition,—we only know that they are the result of interference with functions of parts of the brain not involved in the lesion; and the term "distant symptom" conveys this idea—although perhaps not quite grammatically—without committing us to any theory. The corresponding German term is "Fernwirkung."

Again, the form-sense may be lost in the half field along with the color-sense, while only the light-sense is retained. Furthermore, cases of hemianopsia are on record in which, in part of the defect, both the color and form senses were absent, but the light-sense present, while in the remainder of the defect all three vsiual prceptions were lost.

It is generally held that all lesions of the *Optic Radiations* cause homonymous hemianopsia, but it has not yet been proved that these are all true visual fibers. Henschen indeed believes that only a small portion of them can be regarded as visual fibers, while Vialet's investigations seem to show that the visual path includes the whole of the optic radiations. A lesion here would be distinguished from one in the cortical center by the possibility of hallucinations of vision occurring in the former and not in the latter; and further, there would not be "vision nulle" in the hemianopic defects from lesion in the optic radiations. Lesions of the posterior third of the posterior limb of the *Internal Capsule* (Charcot's "Sensory Crossway") are still believed by some to cause hemianopsia and hemianesthesia of the opposite side of the body; but analysis of clinical cases affords no support to this view, for there are no recorded cases which furnish any definite evidence in this respect. Yet, anatomically, fibers have been traced from the occipital cortex through the optic radiations and internal capsule to the basal ganglia, and thence into the optic tract. The fibers passing through the internal capsule from the external geniculate body are probably reflex fibers simply.

The symptoms due to lesion of the *Primary Optic Ganglia* (Pulvinar, Anterior Corpus Quadrigeminum, and External Geniculate Body) have not as yet been ascertained, the clinical evidence being indefinite. In lesions of the pulvinar alone two typical symptoms occur—viz., hemianopsia and athetosis—and sometimes hemianesthesia may be present.

Hemianopsia from lesions of the *Optic Tract* is characterized by the absence of such symptoms as mind-blindness, word-blindness, etc., which are apt to occur in cortical affections, and by the presence, probably, of other symptoms pointing to a basal lesion. The defects in the fields may be relative (hemiachromatopsia) or incomplete (only homonymous quadrants being lost). Lesions of the optic tract are, of course, apt to implicate the crus cerebri, but do not necessarily do so; and then we would have hemiplegia of the opposite side of the body associated with the hemianopsia.

Symptoms may also be caused by implication of cranial nerves, especially of those which supply the orbital muscles.

Atrophy of the optic nerve, and sometimes neuritis, according to the nature of the lesion, are frequently present.

The characteristic sign which enables us to localize a lesion in the optic tract from one elsewhere causing hemianopsia, is the Hemianopic Pupil (Wernicke's pupil-symptom). Illumination of the amaurotic half of the retina produces a more sluggish reaction than when the light is thrown on the sound side, because the lesion is on the distal side of the corpora quadrigemina, and, consequently, the impulse cannot reach Meynert's fibers to be conducted to the center for the third nerve (see pp. 303 and 311). It must be stated that some observers deny the occurrence of the hemianopic pupil. But, on the other hand, many observers have obtained the symptom. A great obstacle in observing it lies in the difficulty of concentrating the light on the blind side of the retina without allowing it to fall on the good side. If present, this is a valuable sign; but its absence is not decisive, owing to the difficulty of obtaining it.

Wilbrand [4] has proposed an aid in deciding whether the seat of lesion in a case of homonymous hemianopsia is above the primary optic centers (*i. e.*, in the optic radiation or cortex), or in the optic tract. He terms this the Hemianopic Prism Phenomenon, and states that he has found it of practical clinical value. The patient faces a black wall on which a small white mark is made. One eye is closed with a bandage, and the patient is directed to look at the mark with the other eye. A prism of about 12° or 14° is brought suddenly before the eye, its base being so directed that the retinal image of the white spot may be thrown on the half of the retina which does not see (*e. g.*, if the experiment be performed with the right eye, in a case of right homonymous hemianopsia, the inner half of the retina being blind, the prism must be placed opposite the eye with its base inwards). At the same moment the surgeon has to observe whether or not the eye makes such a movement as would tend to bring the retinal image again on the macula lutea (*e. g.*, in the example above chosen the motion would be outwards); and, again, whether or not at the moment of rapid removal of the prism the eye returns to its former position. The prism must be brought rapidly before the eye, in order that the patient may not be able to observe the path of the moving retinal image from the macula lutea towards the boundary line between

the seeing and blind halves of the retina. Those cases in which the boundary line is at or close to the macula lutea are the most favorable for the experiment. If a compensatory movement of the eye takes place when the prism is held before it, then the path through the optic tract to the nuclear oculo-motor centers is free, and the lesion must be situated above these centers, in the optic radiations or cortex. On the other hand, if there is no compensatory motion, the path for the movements of the eye from the retina to the nuclear centers must be interrupted. The explanation of this phenomenon is very similar to that of the hemianopic pupil symptom. Although the motions of the eye for the purpose of fixing visual objects are not, strictly speaking, reflex motions, yet, in each individual they to a great extent become so by long usage, and are controlled mainly by the lower rather than by the cortical centers. Hence, so long as the path for these motions to the nuclear center is uninterrupted, if the retinal image of the visual object be thrown by a prism on a non-seeing part of the retina, the necessary compensatory motion of the eye will be made to bring the image again on the macula lutea.

The Forms of Diseased Process causing a Lesion of the Optic Tract are: syphilitic gummata and syphilitic meningitis; new growths, including tubercle; softening and hemorrhage are rare. Tumors of the optic thalamus, lenticular nucleus, or temporo-sphenoidal lobe may also injure the tract by extension or pressure.

The Prognosis for recovery of vision in the defective halves of the fields depends, of course, upon the nature of the lesion. But recovery is rare, specially in the most common class of cases—those, namely, which are due to cerebral apoplexy.

In Right Homonymous Hemianopsia, wherever the position of the lesion may be, a greater difficulty in reading is experienced than in left hemianopsia. This is partly due to the fact that we read from left to right and that, owing to the defect being on the right side, the word immediately following that at which the patient is looking cannot be seen at the same moment. Knies offers another explanation, namely, that, owing to the right-sided defect, there is loss of the fine co-ordinated movements of the eyes to the right.

Alexia (*ἀ, priv.; λέξις, speech*), or **Word-Blindness**, is the term given by Kussmaul to an inability to understand written or printed characters, although they and other small objects can be distinctly

seen. Other visual objects are named with ease (no visual aphasia). The patient can express his ideas in writing, or write from dictation, yet cannot understand what he has just written, nor can he copy written or printed words. He does understand the meaning of spoken words, and the use of all objects around him (no mind-blindness). He can generally recognize individual letters with some difficulty. This is "pure word-blindness," or "subcortical alexia." When combined with inability to write spontaneously or from dictation, it is known as "cortical alexia" (Wernicke). The condition has been occasionally complicated with hemianopsia. In those cases where an autopsy was obtained the lesion was found in the left occipital lobe. Word-blindness with agraphia or cortical alexia is due, according to Dejerine and Wernicke, to a lesion in the center for visual memory for words, which, in right-handed people, is the left angular gyrus, and inferior parietal lobule.

Visual Aphasia consists in inability to name objects seen, the use of which is known. The objects can be named, if the patient be allowed to feel them, even with his eyes closed. A few cases of this affection have been recorded, and in all there was right homonymous hemianopsia. Alexia and agraphia sometimes coexisted.

Dyslexia.—This symptom was first described by Berlin.[5] In a wide sense it belongs to the aphasic group. It consists in a want of power on the patient's part to read more than a very few—four or five—words consecutively, either aloud or to himself. The difficulty is not caused by dimness of sight, nor by pain in the eye or head, but simply by an unconquerable feeling of dislike or disgust, due to the mental effort. After a few words which can be well understood have been read, the book is pushed away, and the head drawn backwards and turned aside; and then in a moment or two the patient may be tempted to repeat the effort, but with the same result after a very few words have been read. The symptom comes on suddenly, and has been usually the first sign of the presence of cerebral disease. Although in most of the cases the dyslexia disappeared in the course of a few weeks, either permanently or to recur later on, yet other symptoms soon followed its first onset, such as headache, giddiness, aphasia, hemianopsia, paralysis of the tongue, hemianesthesia, hemiplegia, twitching of the facial muscles, etc. Seven or eight cases are on record, and all have ended fatally. The lesion was situated, in all but one of

those cases where an autopsy was obtained, in the neighborhood of Broca's lobe. In one case the left hemisphere was normal, while the right hemisphere was extensively diseased.

Amnestic Color-Blindness is a symptom which is most probably due to a lesion in the occipital lobe, interrupting the paths between the center for vision and the speech center. It has always been accompanied by right homonymous hemianopsia. In this condition the patient sees colors and can recognize them, and he can perform Holmgren's tests, but he is unable to name each color.

Visual Hallucinations may occur in cases of homonymous hemianopsia in the blind side of the field only, being due to irritation of the visual-memory center. Homonymous hemianopic hallucinations, persisting for years without hemianopsia, have also been observed. Visual hallucinations also occur very occasionally in connection with glaucoma, and of this I have seen an example. Hallucinations differ from illusions in that the former are completely subjective, while the latter are perverted sensory impressions.

Mind-Blindness, also called **Optic Amnesia,** is a symptom first observed by Munk [6] in his experiments upon animals. It consists in the loss of power of recognizing objects, while the power of seeing them continues. A whip is seen by the animal, but inspires no terror; a tempting morsel is seen, but excites no desire. The symptom was caused by destruction of a region situated chiefly in the posterior division of the second external convolution of the dog's brain. It has also been observed in man. The patient fails to recognize the most familiar objects by sight. In a case of my own the patient could not recognize his own wife until she spoke. There are two forms of mind-blindness—the cortical and the transcortical. In the former, the lesion is in the center for memory; and the patient has lost the power of visual imagination, and cannot describe visual objects from memory. In the latter, the connecting path between the center for vision and the visual memory center is interrupted, and the patient, though he can describe an object from memory, is unable to recognize it when looking at it. Hemianopsia is present in the majority of cases of mind-blindness; and color-blindness complete or hemianopic is not unusual. The lesion has been found in the occipital lobe sometimes involving the parietal convolutions. It usually consists in hemorrhage or softening and the symptom is consequently sudden in its onset; but it also occurs from tumors.

Exhausting illnesses, by reducing the mental energy, may produce a condition of mind-blindness.

Some authors localize the center for visual memory in the angular gyrus, whilst others take for it the whole of the occipital lobe, except the cuneus and its neighborhood.

The localizing value of orbital paralyses has been explained (p. 466).

Part II.

OCULAR DISEASES AND SYMPTOMS LIABLE TO ACCOMPANY CERTAIN DIFFUSE ORGANIC DISEASES OF THE BRAIN.

There are organic diseases of the brain which are not focal, and which, as they attack extensive regions of the brain substance, may be called diffuse. Under the same heading may be placed some diseased cerebral states which we cannot doubt are organic, although their pathology is as yet unascertained. I propose here to describe the points of ophthalmological interest which accompany some of these diseases.

Disseminated Sclerosis of the Brain and Spinal Cord.—Central Color Scotoma is the most usual defect of sight in this disease, and in a few cases absolute central scotoma is present. Irregular defects in the periphery of the fields—sometimes only for color—or regular concentric contraction may be found. These defects may be in one or in both eyes; they most commonly come on very rapidly, and they may get better, or, after a time, get quite well. Even complete blindness, lasting as long as several months, may occur; but permanent complete blindness is rare. The ophthalmoscopic appearances do not always coincide with the state of the vision; for with marked defect of sight the fundus oculi may be normal, or the vision may be normal, while the optic papilla looks diseased, or both sight and ophthalmoscopic appearances may be abnormal. The most common ophthalmoscopic change is a not very intense atrophic appearance of the whole surface of the papilla, or its temporal third alone may be affected in this way. But in these latter cases, where the temporal third alone shows atrophy, a central scotoma is not necessarily present, nor are the papillo-macular fasciculi in the nerve diseased. In a very few cases optic neuritis is present. The ophthalomoscopic changes may

be in both eyes or in only one. Hemianopsia has not been noted, and therefore the defects of vision are evidently always due to disease in the optic nerve, and not in the chiasma or tract. Uhthoff has shown that in disseminated sclerosis there may be disease in the trunk of the optic nerve, without any abnormal ophthalmoscopic appearances, or defect of sight. Sometimes defects of vision and ophthalmoscopic changes precede all other symptoms by long periods, or they appear in the very early stages of the disease; but more commonly they do not come on until other symptoms have been present for some time.

Isolated paralyses of orbital muscles, nuclear paralysis, and nystagmus are derangements of the oculo-motor apparatus, which are liable to be present in disseminated sclerosis. Marked exterior ophthalmoplegia is rare; but the paralyses of nuclear origin of which there can be no doubt are loss of conjugate motion to one or other side, and defective power of convergence. Nystagmus is present in about 50 per cent. of the cases, and is either of the ordinary kind or consists merely in nystagmic twitchings, more particularly at the extreme lateral positions of the eyeballs. Very slight twitchings in these extreme positions are of no import, as they occur even in the healthy state. As true nystagmus is an uncommon symptom in other diseases of the general nervous system, it is of value in this diagnosis. Nystagmic twitchings, while they do occur in other general nervous diseases, are more common in disseminated sclerosis than in any other of these diseases.

Disseminated sclerosis in its early stages is apt to be mistaken for hysteria, owing to the presence of such symptoms as transitory loss of power in limbs, aphonia, convulsive seizures, hysterical manner, and so on, and here the eye-symptoms may come to our aid. In hysteria the ophthalmoscopic appearances are normal; the fields of vision, if deranged, are contracted, central scotoma being rare, and when the fields are contracted the color boundaries often do not recede in their regular order—the field for red, for example, may be wider than that for the other colors. In hysteria, again, it may be found impossible to examine the color fields at all, all colors being named dark or black; and finally oculo-motor disturbances rarely occur.

Diffuse Sclerosis of the Brain.—In some rare cases of this disease headache, vomiting, and double optic neuritis may lead to the diagnosis of cerebral tumor, and error in diagnosis which, with our present knowledge, it is impossible to avoid, unless there be also

focal symptoms that would point with certainty to a tumor. The mistake will not often occur, as the cases here indicated are exceedingly rare.

General Paralysis of the Insane.—Derangements of the intrinsic muscles of the eyeball, orbital paralyses, atrophy of the optic disc, and mind-blindness are the eye symptoms which may be found in this disease.

The Pupil, etc.—The pupils are usually contracted in the early stages, and dilated at later periods. An early symptom is slight inequality in the pupils, with somewhat sluggish reaction of the wider one, and, also at an early period, there is apt to be loss of the pupil-reflex to sensory stimuli. Later on the larger pupil does not react to light at all, while its fellow does so normally, and sight is good. The so-called paradoxical pupil-symptom is an early augury of coming paralysis, and consists in this, that when a strong beam of light is thrown into the eye with the focal illumination the pupil at first contracts fairly well, then dilates slightly, contracts again, and after a few such oscillations finally dilates widely, although the strong light still shines into the eye. The Argyll Robertson pupil is only found in some cases, and then usually in the late stages, but it does occasionally present itself in the initial stages. Sometimes the pupil is irregular in shape.

Paralyses of Orbital Muscles.—These are of rarer occurrence than paralysis of the pupil; but the third and sixth nerves are occasionally paralyzed even in the early stages, and in these stages, too, ptosis and transient nystagmus and twitchings of the eyelids may be seen.

Optic Atrophy.—This is rare in general paralysis, and is then seen for the most part in the late stages. But it has sometimes come on in a very early period, and has even preceded every other symptom by several years.

Mind-blindness occurs in cases of general paralysis, usually in the advanced stages.

Amaurotic Family Idiocy (Sachs[7]) or **Infantile Cerebral Degeneration with Symmetrical Changes at the Macula Lutea** (Kingdon and Risien Russell[8]).—This disease occurs in children during the first year of life, and most, if not all, of the cases recorded, which may number about fifty, occurred in Jewish families. Family predisposition is strongly marked, as many as five children in a family of seven having been attacked. The causes which have been assigned are neurotic taint, blood relation-

ship between the parents, and traumatism of the mother during pregnancy. Syphilis does not seem to play any part in the etiology. The children are born sound and healthy, and continue to be so for some months. They then cease to develop mentally, and idiocy is gradually established. At the same time paresis or paralysis, either flaccid or spastic, of the greater part of the body appears, while the reflexes may be deficient or increased. Hyperacusis is often present. A chief and very early symptom of the disease is loss of sight, ending in absolute blindness, with certain characteristic ophthalmoscopic appearances, and nystagmus and strabismus are sometimes present. A slowly increasing marasmus leads to a fatal termination before the end of the second year, as a rule. Waren Tay [9] first observed the peculiar ophthalmoscopic appearances, and Sachs described the clinical history, general symptoms, and morbid changes in the brain. The ophthalmoscopic appearances are as follows: There is at first no change in the optic discs. At the macula lutea in each eye there is a large white spot, rather diffuse, with softened edges, and about twice the size of the optic papilla. In its center there is a brownish-red, fairly circular spot, which contrasts strongly with the white around it. This central spot, as Tay says, has not the appearance of a hemorrhage, nor of pigment, but suggests a gap in the white patch through which the healthy structures are seen. In short, the appearance reminds one of that seen in cases of embolism of the central artery of the retina. At a later period, with complete amaurosis, atrophy of the optic nerve is found.

In the brain the diseased appearances consist in degeneration of the pyramidal cells of the cerebral cortex. In the pons and medulla oblongata, degeneration of the pyramidal fibers and of the fillet has been found; and in the spinal cord, degeneration of both the crossed and direct pyramidal tracts was seen. Treacher Collins found edema of the retina around the yellow spot, and Holden discovered disease of the ganglion cells of the retina similar to that in the cerebral cortex. The normal absence of the ganglion cell layer at the macula lutea, and the fact that it is thickest just around the latter, go far to explain the ophthalmoscopic appearances.

Meningitis.—Inflammation of the cerebral meninges, of whatever form, and whether at the base or on the convexity of the brain, is liable to be accompanied by optic neuritis. When the meningitis is at the base, ocular paralyses, pain or anesthesia of

regions supplied by the fifth nerve, and defects in the fields of vision from pressure on the optic tracts or commissure, may be found.

Acute Tubercular Meningitis.—In a small percentage of the cases of this form of meningitis miliary tubercles in the chorioid are present. Optic neuritis is more common than in any other form of meningitis, as are also orbital paralyses, in consequence of the tendency of this form to attack the base of the brain.

Cerebrospinal Meningitis.—Eye-symptoms are often present both in the epidemic and sporadic forms of this disease. Swelling of the eyelids, conjunctivitis, and photophobia are frequent even in the early stages. The pupils may be unequal, contracted, or dilated. There may be ulceration of the cornea, parenchymatous keratitis, or deep purulent infiltrations. Retinitis and plastic irido-chorioiditis, followed by retinal detachment, may be found, or there may be purulent irido-chorioiditis, with purulent infiltration of the vitreous humor, going on to panophthalmitis. If the fundus can be examined, optic neuritis or neuro-retinitis will often be seen, or thrombosis of the central vein, with retinal hemorrhages. Each epidemic of cerebrospinal meningitis is apt to be associated with some one of these conditions as its special type of eye-affection. The eye-affections in cerebrospinal meningitis then are very grave; but some of the cases of irido-chorioiditis do recover, with retention of good sight.

Traumatic Meningitis.—Falls and blows on the head which do not fracture the skull are held by many to be capable of causing meningitis, and occasionally, the inflammatory process, reaching the optic nerve, creeps down it to the optic papilla, where it may be diagnosed with the ophthalmoscope.

Hydrocephalus.—Well-marked papillitis, or neuritic atrophy, is sometimes found in congenital hydrocephalus, or in the hydrocephalus which makes its appearance in infancy; and it would probably be more common, but for the compensation for the increased intracranial pressure, which distention of the sutures and fontanels must provide. In the acquired hydrocephalus of later life, optic neuritis, passing over the optic atrophy, is the rule; and such cases may closely simulate an intracranial tumor in all their other symptoms as well. Bitemporal hemianopsia is apt to be present, owing to pressure on the optic commissure by the distended floor of the third ventricle.

Infantile Paralysis.—Hemianopsia has been noted in a very

few cases of this affection; and papillitis, with some orbital paralysis, has also been seen, but usually there are no eye-symptoms.

Paralysis Agitans, or Parkinson's Disease.—In some cases a fine vibratory tremor may be noticed along the margin of the upper lid, especially when the eyes are closed, and the lids will be found to be unusually rigid on an attempt being made at passive opening of them. The slowness of muscular action in other parts does not affect the motions of the eyeballs. If a patient be called on to look in any direction, the eyes are instantly turned, while the head slowly follows them.

Encephalopathia Saturnia.—Even in the milder cases, transient hemianopsia or amaurosis, which may last for several hours, is sometimes met with. There need be no renal disease, and the visual defect must be taken as the result of the lead-poisoning on the brain. In those cases in which acute cerebral disturbances sets in (convulsions, delirium, coma) it is often attended by optic neuritis, with considerable swelling of the disc, and retinal hemorrhages.

Sometimes the fields are contracted without ophthalmoscopic appearances, as in hysteria. The pupils may be unequal. As headache, vomiting, and convulsions are symptoms of the more serious cases of lead-poisoning, it is evident that when intense optic neuritis is added, the diagnosis between this disease and cerebral tumor has to be considered. The characteristic blue line on the gums, anemia, colic, constipation, drop wrist, and lead in the urine, are the aids to the diagnosis, along with the previous history, including the patient's occupation.

Epilepsy.—A visual aura is more common than any other special sense aura in idiopathic epilepsy. It may take the form of subjective sensations of lights, color, flames, megalopsia or micropsia, etc.; or visual hallucinations may occur; or there may be simple homonymous hemianopsia. Where epilepsy is due to organic brain disease a visual aura, occurring always in homonymous sides of the fields, is important, as indicating the occipital lobe as the region of the brain in which the discharge originates. At the onset of an epileptic fit there is often conjugate lateral deviation of the eyes to the opposite side of the body from that on which the convulsions commence, with rotation of the head in the same direction, while subsequently the eyes may suddenly be turned in the opposite direction. The condition of the pupils vary, often even in one and the same fit. At the onset they are usually normal or

contracted; but during the tonic spasm they become dilated, and remain so until consciousness returns. The pupillary light-reflex is lost—a point of importance in the diagnosis of a true epileptic fit from an hysterical attack, in which latter it is retained. After a fit, rapid changes in the size of the pupil may sometimes be seen, and these are valuable as evidence of the fit having been a genuine one. The ophthalmoscopic appearances during a fit vary in different cases. In some they are normal, in others there is marked pallor of the disc and contraction of the blood-vessels, and, again, in others the papilla is hyperemic and the retinal veins enlarged. Optic neuritis and optic atrophy do not belong to epilepsy; and if found they can be regarded only as complications. Between attacks the fundus may be normal; but it is not unusual to find a high degree of hyperemia of the retina and papilla, which may continue for some days or hours, or may even become chronic. The fields of vision after a fit, and sometimes as a permanent state, are concentrically contracted; or there may be color-blindness, and the central acuteness of vision may be reduced. The state of the fields is a valuable aid in the detection of simulation. Transitory amblyopia (migraine, scotoma, etc.) is more frequent in connection with epilepsy than under any other condition. It may precede the true attack by years, or it may occur with, or for an hour or so before, the fits, or it may be substituted for them. Inasmuch as this transitory amblyopia is often attended by disturbances in speech, in the intelligence, or by passing paralysis, and as both eyes are usually attacked by it, frequently in the form of homonymous hemianopsia, it is obvious that its cause resides in the visual cortex. Occasionally the blindness is monocular, and must then be referred to disturbance in the circulation of the retina or optic nerve. It is held by some authorities that, given a predisposition to epilepsy, irregularities in refraction may at times prove the exciting cause of the disease; and that cases of epilepsy occur, in which the attack is induced by the undue strain put upon the muscular apparatus of the eye by reason of an abnormality of refraction. They also hold that, if correcting glasses be worn by these patients at a sufficiently early period, the fits will cease, or at least in a considerable proportion of the cases. Further investigations on this subject are required, especially as concerns the permanence of cures.

Chorea.—It is probable, I think, that in some cases, at least, of this affection cerebral embolism may be taken as the cause. Sev-

eral instances of embolism of retinal vessels have been seen in immediate connection with the onset of chorea.

In chorea the eyes participate in the irregular jerky motions and the spasm may be so unequal in the two eyes as to cause brief diplopia; although, not being constant, it is little heeded by the patients, and is rarely mentioned by them.

Part III.

OCULAR DISEASES AND SYMPTOMS LIABLE TO ACCOMPANY CERTAIN DISEASES AND INJURIES OF THE SPINAL CORD.

Tabes Dorsalis.—Amongst the ocular complications to be found in this disease *Atrophy of the Optic Nerve* is the most serious. It occurs in about 20 per cent. of the cases, and commences more frequently in the preataxic period than subsequently. Rarely it is the first symptom, preceding all spinal symptoms by from two to twenty years, and it does sometimes commence in the later stages of locomotor ataxy. Coming on in the preataxic stage, optic atrophy seems very often to have, as Benedikt first pointed out, a favorable influence on the spinal disease, the spinal symptoms already existing becoming ameliorated or disappearing, while the further progress of the disease is retarded or averted. It is indeed rare for tabetic patients who go blind at an early stage of the disease to become ataxic later; but if the ataxy is once well marked, it does not improve with a subsequent development of optic atrophy. It sometimes occurs that the onset of optic atrophy in one eye precedes that in the other by a long interval, even by many years; but usually the eyes are affected simultaneously or with a very short interval. The relation between the optic atrophy and the spinal disease is not as yet well understood. The atrophy is probably merely a manifestation of a diseased process in the optic nerve, similar to that which attacks the posterior columns of the cord.

Paralysis and Ataxy of the Orbital Muscles.—Paralyses of orbital muscles in locomotor ataxy occur in about 30 per cent. of the cases. They usually appear in the preataxic stage, and even as an initial symptom, and are of two kinds—namely, the transient paralysis, which lasts a few days or weeks, and may recur; and

the permanent paralysis of one or two muscles. Diplopia is produced by these paralyses, and is often the symptom which first induces the patient to see his doctor. The sixth nerve is the one most commonly paralyzed; but the third nerve is also often paralyzed, including the branch to the levator palpebræ, with resulting ptosis. Loss of power of convergence is often present in commencing tabes, and double exterior ophthalmoplegia, as well as double sixth-nerve paralysis, is sometimes seen; and there can be no doubt but that all these three conditions, and probably also some of the other oculo-motor disturbances in tabes, are of nuclear origin. But the orbital nerves may, it is found, undergo atrophy without their nuclei being altered, and probably, therefore, some of the ocular paralyses here are due to peripheral neuritis.

Ocular ataxy is another not infrequent symptom in tabes. It is sometimes erroneously called nystagmus; but nystagmus is a constant oscillatory motion of the eyeballs, both while the eyes are at rest, and when they are looking at an object, and is extremely rare in tabes. In ocular ataxy, so long as the eyes are at rest, there is no oscillation or twitching; but as soon as an object is carefully looked at, and especially if followed when in motion, and more particularly at the end of the latter, a slight twitching of the eyeballs is seen. It may be found in any stage of tabes.

Pupillary Alterations.—Myosis is the usual state of the pupil in tabes, and is held to be due to paralysis of the pupil-dilating fibers from disease in the front part of the aqueduct of Sylvius. The myosis is often extreme, or "pin-hole," as it is then termed; yet the pupil may react to light and on convergence. The pupil may be of normal size in tabes; but mydriasis, except as part of a third-nerve paralysis, is rare. Again, both in the early and later stages the pupils may be of different sizes.

The Argyll Robertson pupil is an important symptom of tabes. It consists in this, that the pupil, although as a rule contracted, does not respond to the stimulus of light by further contraction, or, if so, but slightly, yet does become more contracted on convergence of the visual axes (or on accommodation). Myosis need not necessarily be present with the Argyll Robertson pupil; the pupil may be of normal size or dilated. The symptom is one of those most regularly found in tabes. It is often an early or initial symptom, and it continues through all the stages of the disease. It is occasionally present in one eye only, and is sometimes quite wanting. (See also p. 310.)

Neither the Argyll Robertson pupil nor primary optic atrophy occurs in peripheral neuritis, a disease which is liable to be sometimes mistaken for tabes.

Paralysis of Accommodation without paralysis of the sphincter iridis is a rare symptom in tabes. It is more common in the late than in the early stages.

Narrowing of the Palpebral Fissure, due to a slight drooping of the eyelids, hardly to be called ptosis, sometimes occurs in tabes along with the myosis. It is held to be due to paralysis of the sympathetic (sympathetic ptosis), is usually binocular, and the frequency of its occurrence increases as the disease advances.

Twitchings in the Orbicularis Muscle for some Moments after Closure of the Eyelids may sometimes be observed in tabes. Similar twitchings may occasionally be seen in some other nervous diseases, and even in health, but less well marked. Probably their marked character in tabes is due to very slight facial paralysis, and the consequent imperfect power of closing the eyelids.

Epiphora is stated by some authors to be not rare in tabes, but others deny this, and I have not myself observed it to be so.

Reduction of Intra-ocular Tension is a symptom in tabes to which as yet Berger alone has drawn attention. He found it present in thirty-five out of one hundred and nine cases examined.

Hereditary Ataxy (Friedreich's Disease) has few eye-symptoms, a fact of some diagnostic importance. Ataxic nystagmus, as Friedreich pointed out, is the only one which occurs with any constancy. Optic atrophy is of such rare occurrence in the disease that it can hardly be reckoned as one of its symptoms. Paralyses of orbital muscles do not occur, nor does any pupil-symptom.

Myelitis.—Apart from the inflammation of its meninges (cerebrospinal meningitis), of which I have already spoken, acute inflammation of the cord may be associated with optic neuritis. The optic nerve seems usually to become inflamed before the spinal cord, but the myelitis may precede the optic neuritis, or optic nerve and spinal cord may be simultaneously attacked. The relation of the optic neuritis and myelitis to each other is, doubtless, nothing more than that each is a manifestation of the presence in the system of one and the same toxic influence, whatever it may be. Rheumatism, epidemic influenza, and syphilis are amongst the causes assigned in some cases, while in others no cause could be assigned. If the cervical portion of the cord is in-

flamed, pupillary symptoms—irritation mydriasis or paralytic myosis—are apt to be present.

Syringomyelia and **Morvan's Disease.**—One eye-symptom is common to both of these diseases (which are, indeed, held by many to be one and the same disease)—namely, a concentric contraction of the field of vision without ophthalmoscopic changes. It is not quite certain whether this abnormality of the field is due, at least sometimes, to attendant hysteria, or is always a symptom of the organic disease as such. Inequality of the pupils has sometimes been noted.

Myotonia Congenita (Thomson's Disease).—In some cases of this rare disease the external musculature of the eyes affords symptoms, although the intrinsic muscles are never disordered. The opening and closing of the eyelids may be difficult—they cannot be closed or opened at one stroke, successive jerky motions being required to effect closure or opening. As in Graves' disease when the eyes are open the upper lid is apt to be retracted, and the upper lid does not readily follow the downward motions of the eyeball. Transitory amblyopia, or even amaurosis, has been noted in some cases.

Acute Ascending Paralysis (Landry's Disease).—Eye-symptoms are rare in this disease, but there may be paralysis of some of the orbital muscles, paralysis of accommodation, mydriasis, or loss of the light-reflex.

Injuries of the Spinal Cord.—The condition which used to be known as railway spine, but which is now better styled traumatic neurosis, and is due to mental shock rather than to organic lesions of the brain and spinal cord, is accompanied frequently by certain functional eye-symptoms of which the chief one is a contraction of the field of vision similar to that found in some cases of hysteria. In those much rarer cases of organic injury to the cord, or of myelitis, or of hemorrhage in, or inflammation of, its membranes, following on railway and other accidents, organic eye-disease seldom results, although optic neuritis and optic atrophy used to be held to be frequent consequences of these injuries. If the lesion be in the lower cervical region of the cord, the pupils are apt to be contracted from sympathetic paralysis.

Part IV.

NERVOUS AMBLYOPIA, OR NERVOUS ASTHENOPIA.

We find Nervous Amblyopia, or Nervous Asthenopia, for the most part in connection with three functional disorders of the nervous system—namely, Neurasthenia, Hysteria, and Traumatic Neurosis. Many observers, it is true, hold that these three conditions ought to be regarded and treated of as hysteria, that the term neurasthenia is quite superfluous, while traumatic neurosis is merely hysteria caused by shock. This is not the place to enter into a discussion on this question; and it need only be said that while these various states of the nervous system are admitted on all hands to have much in common, and also to merge insensibly into each other, yet typical cases of each are sufficiently differentiated to make it justifiable, and convenient, for the present, to retain all three in our minds, as separate clinical entities.

It may in general be stated that Neurasthenia is abnormal susceptibility of the system to fatigue from mental or bodily exertion; while in hysteria the symptoms depend upon idea, the essence of hysterical conditions being that ideas too easily excite abnormal changes in the organism.

The defects of vision which accompany these disorders are, like all their other symptoms, purely functional—*i. e.*, they do not depend on any organic disease in the retina, or other portions of the visual apparatus, but merely upon derangement of the functions of these parts. Consequently, there are no ophthalmoscopic changes in the fundus oculi.

In the following, the derangements of vision most liable to be found in each condition will be pointed out, but here it is desirable in the first instance to state them in a general way. Complete blindness of one or both eyes may be found, but is rare; a diminished, but fluctuating, acuteness of vision is more common, the effort or desire to see well being often the signal for the acuteness of vision to fail, and objects disappear from sight if looked at long. Attacks of defective sight, too, may come on suddenly without any provocation, accompanied by positive scotomata, and may last for some minutes. But the most remarkable, important, and characteristic symptom is concentric contraction of the fields of

vision. It is almost always necessary, in order to ascertain the presence of this symptom, to examine the fields with the perimeter —no rougher method will answer—and it is most important to use a test-object of not more than 5 mm. square. Concentric contraction of the field is, we know, a symptom in optic atrophy and in glaucoma; but, while in those diseases the contraction usually advances with more or less deep re-entering angles directed towards the fixation point, in nervous amblyopia the contraction is about equal in degree in each meridian, and hence the seeing portion of the field which is left presents a somewhat circular shape. This shape of the field with normal ophthalmoscopic appearances is pathognomonic of the condition. The contraction may be but slight, or it may approach to within 10° or 5° of the fixation point. It is almost invariably present in both eyes; but it is often much more marked in one eye than in the other.

Associated sometimes with this concentric contraction, and sometimes without it, is a phenomenon known as the fatigue field. It consists in this, that if the test-object be brought from the periphery towards the fixation point in each meridian successively, it will be found that the outside limit of the field is nearer the fixation point on each successive meridian examined, without regard to the part of the field in which the examination is commenced. Or, if the test-object be brought in the horizontal meridian from the periphery on, say, the temporal side across the field until it disappears on the nasal side, and the points of entrance and of exit noted, and the object be immediately carried back on the same meridian until it disappears on the nasal side, and the entrance and exit again noted, and this maneuver repeated five or six times; should fatigue be present, it will be shown by the points of entrance and exit coming nearer and nearer to the fixation point on each journey—in short, the field is becoming more and more contracted. This method of taking the field in these cases has been proposed by Wilbrand, and is useful, too, as showing whether at the beginning there is any concentric contraction of the field.

These two modes of examination are practically the same; and the reason for the form of fields they are intended to bring out is that the longer in each case the examination is continued the more fatigued does the nervous visual apparatus (be it cerebral center, or retina, or both) become, and this exhaustion is most marked in the periphery of the field. In the normal state the boundary of

the field is not much affected by the length of the examination. Ring-form and island-like defects in various parts of the field, and which come and go, are recognized as functional defects, and cannot be confused with the continuing central scotoma of toxic amblyopia due to disease in the papillo-macular fibers. In addition to the defective sight, or contraction of the fields, or fleeting scotomata, there are often other eye-symptoms present, such as weakness of accommodation, or of the internal recti, or some derangement of the fifth or facial nerves.

While functional derangements of vision, as distinguished from those due to organic disease, are what are here under consideration, yet it is very necessary to mention that visual defects due to organic disease may sometimes be aggravated by functional blindness. In tabes with optic atrophy, for instance, the contraction of the field may become suddenly increased with the occurrence of some mental worry or intercurrent general illness, and become restored again to its former dimensions with the return to a calmer state of mind or to improved health. In homonymous hemianopsia there is often a peripheral contraction in the seeing side of the field, which can only be due to diminished functional activity in the opposite side of the brain from that in which the disease is situated.

In the three disorders of the nervous system mentioned, the symptoms may in a given case remain confined to the nerves which are associated with the various functions of the eye; but this is rare. It is more common to find also symptoms provided by the derangement of functions in other parts of the nervous system.

Nervous Amblyopia in Neurasthenia.—School-children and those of that age are very liable to become neurasthenic. They are brought to the physician with the complaints that the sight is confused, that print disappears as they look at it, that reading causes the eyes to smart and run with water, and that it brings on headache. If the patient be required to read aloud he soon stops, complaining that the words are running into each other, and the book is then brought closer to the eyes; then a few more words are read, and the book is brought still closer, until, finally, it is nearly in contact with the nose; and then further attempts to see are made by twisting the head about, turning the book towards the light, frowning, and so on. Obviously what causes this difficulty in reading is a rapid exhaustion of the accommodation. Insufficiency of the internal recti is also often present, and would

contribute to the difficulty of use for near work. The eyes are often emmetropic, and the amplitude of accommodation is normal. Examination of the fields will often discover them to be concentrically contracted, and the fatigue field, too, is frequently present. With these asthenic symptoms there are often symptoms of exalted sensibility of the visual apparatus, such as photopsiæ (bright spots, colored balls, glittering surfaces, etc., before the eyes), a prolonged continuance of the after-images of objects, increased sensitiveness to daylight, and still more so to artificial light, and visual hallucinations (heads, animals, passing shadows, etc.). In the neurasthenia of school-children eye-symptoms often predominate, but other nervous symptoms are nearly always present, such as hallucinations of hearing, states of uncalled-for joyous excitement, or of mental depression, or of irritability of temper. Vertigo, a tendency to weep, some loss of memory, and insomnia may all, or any, of them be present. The patellar reflex is usually increased. Patches of diminished sensation may be found here and there over the surface of the body, although completely anesthetic patches, or hemianesthesia, are rare.

In school-children complaints of difficulty in reading suggest malingering in many instances, but it is not wise to adopt this view without good grounds for it. An examination of the fields may set the question at rest, for neither the concentrically contracted field nor the fatigue field can be malingered.

The neurasthenia of adults manifests itself, so far as eye-symptoms are concerned, less in the use for near work than is the case with school-children. In them, moreover, the contraction of the fields is usually slight, while the fatigue field is well-marked. These patients come complaining of unpleasant and painful sensations in and around the eye, such as creeping sensations and boring pains in the orbit, stabbings in the eyeball, a sensation as if the eye were turned round in the head, uneasy feelings attending the motions of the globe. The eye may be very painful on pressure at some one spot without apparent cause; and there are often uncomfortable sensations of cold, burning, or dryness under the lids. If there be an error of refraction it is difficult to find glasses with which the patient will be content, the bridge and wings of the frames annoying them with their slight pressure, while the reflection of light from the margins of the eye-pieces causes dazzling. The patients are very sensitive to any bright

light. The central acuteness of vision is usually normal, but use of the eyes for near work causes headache, often in the form of a hammering in the temples, or a sensation of pressure on the vertex.

Treatment.—Tinted protection spectacles. Abstinence from use of the eyes for near work. A general tonic treatment, including cold sponge baths when they can be borne, bracing air, plenty of exercise in the open air short of fatigue, early hours, and easily digested diet. As regards drugs, strychnin and iron are those from which most can be expected.

Nervous Amblyopia in Hysteria.—Nervous amblyopia, or nervous asthenopia, in hysteria is often very similar to that in the neurasthenia of school-children, except that the difficulty for near work is even greater. Tonic blepharospasm and partial paralysis of orbital muscles may accompany it. The field of vision is commonly more contracted in one eye than in the other, or the contraction may be very marked in one field, while the other field is normal or nearly so. In neurasthenia the contraction is usually about equal in each eye. Orientation is rendered more difficult by the hysterical than by the neurasthenic field. A high degree of blindness, or even complete amaurosis, may attack a neurasthenic school-child for a few minutes; but in hysteria such attacks, which may occur in both eyes, but are usually confined to one eye, are likely to last for weeks, or months, or longer. In the amblyopia of hysteria, we may find that an eye which cannot see moderately sized type is enabled to do so by placing any plane glass as spectacles before the eye. Such an occurrence by no means proves that the patient is malingering; it shows, rather, that the psychical inhibition to the function of sight in the eye has been withdrawn by the suggestion provided by the spectacles.

With monocular amblyopia or amaurosis, there is usually hemianesthesia of the same side of the body as the blind eye; or, if there be merely contraction of the fields, there is often hemianesthesia of the side of the most contracted field.

The pupils vary much in these cases, and even in one and the same case from time to time. They may be normal, or wide and immovable, contracted, or of different size in each eye.

Nervous Amblyopia in Traumatic Neurosis.—In traumatic neurosis, one of the most important and most constant of the symptoms is concentric contraction of the field of vision. Yet it is often absent, and, when present, is not always sufficiently typical

in form to enable it to be utilized in the diagnosis. It is not often so pronounced as to interfere with orientation, and must be sought for with the perimeter to determine its presence. The boundaries for the color-fields are affected even more than that for white, and consequently the tests for these boundaries may discover the contraction more readily than examination of the boundary for white. The relative position of the color boundaries is seldom altered, and color-blindness is seldom present. The defect in the field is usually to be found in both eyes, and if there be hemianesthesia it is on the side of the most contracted field. It is an important fact that the contraction of the field may be the only derangement of sensation, either special or general. The contraction is liable to continue for months or years, and to become more marked for a time, as the result of any passing mental disturbance. The fatigue field, too, is present in some cases of traumatic neurosis.

As regards other ocular symptoms in traumatic neurosis: the pupil-reflex is usually normal, but is occasionally wanting, and a difference in size of the pupils may sometimes be noted; paralyses of orbital muscles are rare, but insufficiency of the internal recti is not uncommon; sensations of sparks, colors, and waviness before the eyes are sometimes complained of; photophobia, and sensations of dazzling with their resulting blepharospasm, may be present.

It is not desirable to rest content with one examination of the field of vision which may prove negative in its result, for it is only shown thereby that on that particular occasion the field was normal. At a later period a defect may be found.

Part V.

VARIOUS FORMS OF AMBLYOPIA.

Transitory Hemianopsia, or Scintillating Scotoma.—This affection is characterized by (1) symmetrical defects in the fields of vision, usually of the hemianopic type, and (2) vibrating or scintillating luminous sensations, which after a short time disappear, and are followed by an attack of (3) migraine. In fact, the visual troubles belong to the symptoms of migraine.

The scintillations and defects in the fields, either of which may occur first, commence over a small area, generally near the center

of the field, and gradually widen out; the flashing increases in intensity, and often assumes a zigzag shape, like fortifications, at the periphery of the defect in the field. And this defect may exist as symmetrical scotomata, complete or partial homonymous hemianopsia, or even altitudinal hemianopsia. In some cases the scintillation may be absent, while in others the attack of migraine does not follow. The ocular symptoms, which last from a period varying from a few minutes to half an hour, are not accompanied by any changes in the fundus oculi, and always end in complete recovery. Vertigo, nausea, or sickness, and even slight aphasia sometimes accompany the headache.

This affection occurs most frequently in intellectually active individuals; fatigue, long reading, and hunger have been known to bring on attacks. The symptoms are most probably due to disturbances in the cerebral circulation.

Treatment should be directed to the cause of the migraine. Lying with the head low, or stimulation of the circulation by wine or nitro-glycerin, sometimes cuts short an attack.

Congenital Amblyopia.—This condition is not very uncommon. Ophthalmologists, in the course of their practice, come across people in whom the vision of both eyes is below the normal standard, even with perfect correction of any error in refraction, and who declare that they never have seen better, and that their sight is not getting worse. Still more common is congenital amblyopia in one eye. As a rule the field of vision and the color-vision are normal, but cases are seen in which there is contraction of the field, with defective color-sight.

The *Ophthalmoscopic Appearances* are normal.

Reflex Amblyopia is said to have been observed, and chiefly in connection with irritation of the fifth pair, especially its dental branches; but I have not seen these cases, and I am rather skeptical as to their occurrence. Carious molar teeth are reputed to be its frequent cause, usually with severe toothache, but sometimes without it. The defect of vision may be confined to the side of the carious tooth, and is nearly always most marked on that side. It is said that it may be of extreme degree, vision being reduced even to the merest perception of light.

More generally recognized than amblyopia, as the result of toothache, are: hyperesthesia of the retina, photophobia, subjective sensations of light, and diminution in the amplitude of accommodation.

All these symptoms, even amblyopia of the severest type, disappear when the dental affection is relieved.

Many cases are on record in which wounds of the supra-orbital nerve were looked on as the cause of amblyopia or of amaurosis; but it is by no means certain that an ophthalmoscopic examination would not have afforded another explanation in many of these cases. Yet even nowadays many hold that wounds of the supra-orbital region can produce amblyopia, as cases are said to have been cured by division of the nerve involved in a cicatrix that was tender on pressure.

Sympathetic Irritation (p. 266) is to be included under this heading. It is seen in the sound eye in some cases of cyclitis, and must not be confounded with sympathetic ophthalmitis, which comes about in quite a different way. Its symptoms are: diminution of the amplitude of accommodation, asthenopia, hyperesthesia of the retina, lacrimation, and subjective appearances of light.

Removal of the exciting eye, if otherwise indicated, always relieves sympathetic irritation; but where this is not admissible the dark room, atropin, dry cupping at the temple, with bromid of potassium internally, may be employed.

The *Ophthalmoscopic Appearances* in reflex amblyopia are normal.

Night-Blindness.—This is a well-recognized symptom of the disease known as Retinitis Pigmentosa (p. 407). I have seen an instance of congenital night-blindness in five members of a family of ten children without ophthalmoscopic signs; and Richter, quoted by Lawrence, observed a similar instance. But the condition of which I have here to speak is Acute, or Idiopathic, Night-Blindness.

The patients can see well in good daylight; but on a very dull day, or in the dusk of evening, or by indifferent artificial light, their vision sinks very much more than that of persons with normal eyes. They are then unable to see small objects, which are quite plain to other people, and in a still worse light they fail even to recognize large objects visible to everyone else. This peculiar visual defect is due to imperfect adaptation power of the retina, and not to defective light-sense, as is sometimes stated.

Conjunctivitis and xerosis of the conjunctiva are often present in acute night-blindness (p. 125). Some observers have found micrococci and bacilli in the conjunctiva in these cases, and have regarded these organisms as the cause of the conjunctival affec-

tion. It seems now more probable that they are merely secondary to the xerosis.

The connection between night-blindness and xerosis conjunctivæ remains to be explained; but it is likely that they are both results of the one cause.

Acute night-blindness is often the result of long-continued dazzling by very bright sunlight, or of lengthened exposure to bright firelight (*e. g.,* in foundries), and it is probable that in many, if not in most, instances of this affection, defective nutrition of the system plays the chief *rôle* in rendering the patients liable to it. Thus, in scorbutus, acute night-blindness has been frequently seen, when the patients have been exposed to strong glares of sunlight.

Treatment consists in protection from light,—in short, in complete darkness for a time,—and then gradual return to ordinary daylight; while the system is to be strengthened by careful dietary and suitable tonic medicines.

Uremic Amblyopia.—This is most commonly seen in connection with the nephritis of pregnancy and scarlatina, but may occur in any case of uremic poisoning. It is met with in the acute forms of nephritis, in which albuminuric retinitis is not so liable to occur. The blindness is usually absolute, and may come on suddenly or with a short previous stage of dimness of vision. It lasts from twelve hours to two or three days, and may recover completely, but in some cases a central scotoma remains.

The *Ophthalmoscopic Appearances* are negative.

Treatment can only be directed to the general condition.

The Prognosis for vision is good, as it always recovers if the patient's life be spared.

Pretended Amaurosis.—Malingerers rarely pretend total blindness of both eyes, and such cases can often only be detected by constant observation of their actions.

Presence of pupillary reflex is no proof that the patient sees, for this would be quite compatible with a cortical lesion causing total loss of sight (p. 312).

The crossed diplopia test (*vide infra*) may be employed in these cases; for if both eyes see, the one with the prism will rotate inwards for the sake of single vision, while if both eyes be blind, of course no such motion will take place. Again, if the malingerer's own hand be placed in various positions, and he be asked to look at it, he will in all probability look in some other direction; whereas

a truly blind man usually makes a fair attempt at directing his eyes towards his own hand.

Pretended monocular amaurosis can generally be detected by the Diplopia Test. If the malingerer be made to look, with both eyes open, at a lighted candle placed some feet off, while a prism with its base downwards is held before the admittedly good eye, he will say he sees two images of the light one over the other. Were he blind of one eye he would not see two images.

Another method—the Crossed Diplopia Test—consists in holding a prism of some 10° or 12° with its base outwards before the pretended blind eye, when, if it sees, it will make a rotation inwards for the sake of single vision, an effort which a blind eye would not make.

Alfred Graefe's Method.—In this test the pretended blind eye is covered with the surgeon's hand from behind the patient, while with the other hand a prism (about 10°) is held base down before the good eye, so that its edge may pass horizontally across the center of the pupil. Monocular double vision results, as the rays pass through the upper part of the pupil normally, while through the lower part of it they are refracted downwards by the prism. The double images stand over each other. If now the hand which excludes the pretended blind eye be rapidly removed, while at the same moment the prism is moved upwards, so that the entire pupil is covered by it, a malingerer will still see double images standing one over the other; for now the diplopia must be binocular.

Harlan's Test consists in placing a trial frame on the patient's nose with a very high + lens—say + 14 D—opposite the good eye, by which means it is excluded from distant vision, and a plane glass—or a 0.25 D convex or concave lens, which of course would not materially interfere with its distant vision—opposite the pretended blind eye. The patient then, believing there is much the same kind of glass before each eye, will read the test-types; and if it be now desired to expose the deception, the pretended blind eye is excluded from sight, and the malingerer will then be unable to read the test-types.

Snellen's Colored Types may also be used for this purpose. These types are printed in green and red. If a person be really blind of one eye, he will, of course, see both the green and the red letters with the good eye. But if a green glass be held before the good eye, the rays from the red letters will be excluded, and he will now only see the green letters; or with a red glass the red

letters alone will be seen. A malingerer may be detected by holding before his admittedly good eye a green glass; and if he now still see the red letters, it must be that he does so with the so-called blind eye.

It is well to have this variety of tests, in order that they may be used to corroborate each other.

Erythropsia (*ἐρυθρός, red*)—**Red Vision.** A large number of cases of this remarkable affection are on record; indeed, it will have come under the notice of nearly every ophthalmic surgeon of any experience. Two-thirds of the cases have been subjects of successful cataract operations, whilst the remainder have possessed normal eyes. In some cases the red vision remains only a few minutes, and does not again return; whilst in others it appears every day for a short time, for weeks or months; and, again, in others it continues for several days, and then disappears for good or recurs at intervals. In the aphakic cases it does not usually appear for weeks or months after the removal of the cataract, and in one case the interval was two years. During the attacks the patients see all objects of a deep red color, and occasionally of a purple or violet hue. In no instance is the acuteness of vision affected either during or after the attacks.

A satisfactory explanation for the affection has not yet been offered. It seems probable that it is due to overexcitation of the visual nervous apparatus—some believe of the visual center, others of the retina—set a-going by exposure of the retina to strong light, along with other favoring circumstances, especially general over-excitement of the body or mind. More than this cannot at present be said. Why aphakic eyes should be so much more liable to erythropsia than eyes which possess their crystalline lenses is an enigma.

Treatment seems to have but little effect. Protection of the eyes from light has not been of use. Bromid of potassium internally seems to have done some good in those cases where it was tried.

References.

[1] "Klinische und Anat. Beiträge zur Pathol. des Gehirns." Upsala, 1890-92.

[2] Vialet, "Les Centres Cérébraux de la Vision." Paris, 1893.

[3] "American Journal of the Med. Sciences," January, 1887.

[4] "Zeitschrift für Augenheilkunde," i. p. 125.

[5] "Archiv. f. Psych.," vol. xv. p. 276, and in his monograph, "Eine besondere Art der Wortblindheit (Dyslexie)." Wiesbaden, 1887.

[6] "Zur Physiologie der Grosshirnrinde," "Archiv f. Anat. und Physiol.," v. and vi. pp. 162 and 547.

[7] "Journal of Nervous and Mental Disease," 1887, p. 541; and "New York Medical Journal," May 30, 1896.

[8] "Trans. Med. Chir. Soc.," lxxx.

[9] "Trans. Ophthal. Soc., U. K.," i. p. 55, iv. p. 158.

APPENDIX I.

HOLMGREN'S METHOD FOR TESTING THE COLOR-SENSE.

For the purposes of this method a selection of Berlin worsteds is made, including red, orange, yellow, yellow-green, pure green, blue-green, blue, violet, purple, pink, brown, gray; several shades of each color being present, and at least five gradations of each tint, from the deepest to the lightest. Green and gray, several kinds each of pink, blue, and violet, and the pale gray shades of brown, yellow, red, and pink must be well represented. But no two samples are to be of precisely the same shade of the same color. This large number of colors and shades is used because the color-blind person escapes detection with more difficulty, and the diagnosis therefore is all the more certainly made, the greater the variety of colors. The normal-eyed individual readily selects the right ones from the mass; whilst the color-blind person, although the right ones are directly before him, picks out wrong ones, thereby disclosing the character of his defect.

The test-color with which the examination invariably begins is a pale pure green, because green is the whitest of the spectral colors, and, consequently, the one in which the presence of color is most difficult to recognize —the one, in short, most easily mistaken for gray (= no color). We all experience the most difficulty in deciding whether there be any color at all present in the very deepest shades (nearly black), and in the very palest shades (nearly white); therefore it was plainly either a very dark or a very pale shade of green that should be employed, and Holmgren's experience made him decide for the pale shade, as providing the most delicate test.

As a test for the diagnosis of the particular kind of color-blindness, Holmgren recommends a purple (deep pink) sample—that is, the whole group of colors in which red (orange) and blue (violet) are combined in nearly equal proportions, or at least in such proportions that no one of them sufficiently preponderates over the others, to the normal color-sense, so as to give its name to the combination. Purple is of especial importance in the examination of the color-blind, for the reason that it forms a combination of two fundamental colors (red and blue)—the two extreme colors—which are never confounded with each other.

The Method of Examination and of Diagnosis is as follows: The worsteds are placed in a pile on a table in broad daylight. The test skein (pale pure green) is taken from the pile, and laid at a short distance from it, so as not to be confounded with the other skeins during the trial; and the person examined is then requested to select other

skeins most resembling this in color, and to place them by the side of the sample. It is necessary he should have clearly understood what is required of him—namely, that he should search the pile for the skeins making an impression on his chromatic sense similar to that made by the sample, and independently of any name he may give the color. Indeed, it is not desirable that he should be asked to name the colors, and he should be discouraged from doing so. The examiner should explain that resemblance in every respect is not necessary; that no two specimens are just alike; that the only question is the resemblance of color; and that, consequently, he must endeavor to find something lighter and darker of the same color. If the person examined cannot understand this verbal explanation, the examiner must resort to action. He must himself make the trial by searching with his two hands for the skeins, thereby showing what is meant by a shade, and afterwards restoring the whole to the pile, except the sample-skein. Or, when a large number of persons have to be examined together, it will be more rapid to begin at once with such a demonstration before the assemblage. There is no loss of security in this, for no one with defective chromatic sense finds the correct skeins in the pile any the more easily from the fact of having a moment before seen someone else looking for and arranging them.

On the card which is attached to the inside of the back cover of this book there are two classes of wool-samples. (1) The Test-Samples, which are placed horizontally. (2) The Colors of Confusion—that is, those which the color-blind person selects from the heap of wools, because he confuses them with the color of the sample—and these are arranged vertically under their respective test-samples. The object of this card is merely to illustrate this text. It cannot itself be used for testing the color-sense, nor does it contain all the colors and shades necessary for the purpose.

The test is conducted as follows: *Test I.* The green sample is presented. This sample, as already explained, should be of the palest shade of very pure green, which is neither yellow-green nor blue-green to the normal eye, but fairly intermediate between the two. The examination must be continued until the person examined has selected all the other skeins of the same color, or else, with these or separately, one or several skeins of the class corresponding to the "colors of confusion" (1 to 5), until he has sufficiently proved, by his manner of doing it, that he can easily and unerringly distinguish the confusion colors, or until he has given proof of unmistakable difficulty in accomplishing his task. He who places beside the sample one of the colors of confusion (1 to 5)—that is to say, finds that it resembles the test-sample—is color-blind. He who, without being quite guilty of this confusion, evinces a manifest disposition to do so, has a feeble chromatic sense.

If we want to know the kind and degree of the color-blindness which the failure to perform *Test I.* shows to be present, we must proceed to

Test IIa.—A purple skein is shown to the person being examined. The trial must be continued until he has selected all, or the greater

part of, the skeins of the same color, or else, simultaneously or separately, one or several skeins of "confusion" (6 to 9). He who confuses selects either the light or deep shades of blue and violet, especially the deep shades (6 and 7), or the light or deep shades of one kind of green or gray, inclining to blue (8 and 9). (1) He who is color-blind by *Test I.*, and who, upon *Test IIa,* selects only purple skeins, is termed "incompletely color-blind." (2) He who, in *Test IIa,* selects with purple only blue and violet, or one of them, is "completely red-blind." (3) He who, in *Test IIa,* selects with purple only green and gray, or one of them, is "completely green-blind." The red-blind never selects the colors taken by the green-blind, or *vice versa.* Often the green-blind places a violet or blue skein beside the green, but only the brightest shades of these colors. This does not influence the diagnosis.

The examination may end here, and the diagnosis be regarded as settled. But to convince railway employers, and shipowners, and their employees, a still further trial may be made. It only serves to corroborate the diagnosis.

Test IIb.—The red skein is presented. It is necessary to have a vivid red color, like the red flag used as signals on railways. This test, which is applied only to those either "completely red-blind" or "completely green-blind," should be continued until the person examined has placed beside the specimen all the skeins belonging to this shade, or the greater part of them, or else, separately, one or several "colors of confusion" (10 to 13). Alongside the red the red-blind person places green and brown shades, which (10 and 11) to the normal sense seem darker than red. On the other hand, the green-blind person selects opposite shades —those which appear lighter than red (12 and 13). Every case of complete color-blindness discovered does not always make the precise mistakes just mentioned with *Test IIb.* These exceptions are either persons with comparatively inferior degrees of complete color-blindness, or of color-blind persons who have been exercised in the colors of signals, and who try not to be discovered. They usually, but not always, confound at least green and brown. Total color-blindness is extremely rare, but such a case would be recognized by a confusion of every color having the same intensity of light.

Violet-blindness will be recognized by a genuine confusion of purple, red, and orange in *Test IIb.*

If further information on the subject be desired, the reader should consult Professor Holmgren's original monograph, "De la Cécité des Couleurs," Stockholm, 1877, or Dr. Joy Jeffries' "Color-Blindness," Boston, 1879.

APPENDIX II.

REGULATIONS AS TO DEFECTS OF VISION WHICH DISQUALIFY CANDIDATES FOR ADMISSION INTO THE CIVIL, NAVAL, AND MILITARY GOVERNMENT SERVICES, THE ROYAL IRISH CONSTABULARY, AND THE MERCANTILE MARINE.

By an Army Circular issued by the War Office—

Candidates for Commissions in the Army are required to possess the following visual powers. These regulations apply to all branches of the service, including the Medical Department.

Snellen's Test Types are used for determining the acuteness of vision.

1. If a candidate can read D = 6 at 6 meters (20 English feet), and D = 0.6, at any distance selected by himself, with each eye, without glasses, he will be considered fit.

2. If a candidate can only read D = 24 at 6 meters (20 English feet) with each eye without glasses, his visual deficiency being due to faulty refraction, which can be corrected by glasses which will enable him to read D = 6 at 6 meters with one eye, and D = 12 at the same distance with the other eye, and can also read D = 0.8 with each eye without glasses, at any distance selected by himself, he will be considered fit.

3. If a candidate cannot read D = 24 at 6 meters (20 English feet) with each eye without glasses, notwithstanding that he can read D = 0.6 he will be considered unfit.

Squint, inability to distinguish the principal colors, or any morbid conditions, subject to the risk of aggravation or recurrence in either eye, will cause the rejection of a candidate.

The Royal Navy.—A candidate is disqualified unless both eyes are emmetropic. The candidate's acuteness of vision and range of accommodation must be perfect.

2. A candidate is disqualified by any imperfection of his color-sense.

The author has it on good authority that no absolute rule as to vision is laid down with regard to candidates for entry into the Navy Medical Service. Each case is determined at the physical examination at the Naval Medical Department, which takes place shortly before the competition.

Full normal vision is not necessarily essential in all cases for Naval Medical officers.

3. Strabismus, or any defective action of the exterior muscles of the eyeball, disqualifies a candidate for the Royal Navy.

The Home Civil Service.—Blindness or defective vision, except a moderate degree of ordinary shortsight, disqualifies. But candidates for the Customs Outdoor Service must not be shortsighted.

The Indian Civil Service (*Covenanted and Uncovenanted*).—1. A candidate may be admitted into the Civil Service of the Government of India if ametropic in one or both eyes, provided that, with correcting lenses, the acuteness of vision be not less than 6-9 in one eye and 6-6 in the other, there being no morbid changes in the fundus of either eye.

2. Cases of myopia, however, with a posterior staphyloma, may be admitted into the service, provided the ametropia in either eye do not exceed 2.5 D, and no active morbid changes of chorioid or retina be present.

3. A candidate who has a defect of vision arising from nebula of the cornea is disqualified if the sight of either eye be less than 6-12; and in such a case the acuteness of vision in the better eye must equal 6-6, with or without glasses.

4. Paralysis of one or more of the exterior muscles of the eyeball disqualifies a candidate for the Indian Civil Service. In the case of a candidate said to have been cured of strabismus by operation, but without restoration of binocular vision, if with correcting glasses the vision reach the above standard (1), and if the movement of each eye be good, the candidate may be passed. The same rule applies to the case of unequal ametropia without binocular vision, both eyes having full acuteness of vision with glasses, and good movement.

The Indian Marine Service (*Including Engineers and Firemen*).—1. A candidate is disqualified if he have an error of refraction in one or both eyes which is not neutralized by a concave, or by a convex 1 D lens, or some lower power.

2. A candidate is disqualified if he cannot distinguish the primary colors and their various shades—red, green, violet or blue, and yellow.

3. Strabismus, or any defective action of the exterior muscles of the eyeball, disqualifies a candidate for the Marine Service.

Royal Irish Constabulary.—Candidates for cadetships in the Royal Irish Constabulary and recruits must be able to read with each eye separately, and without glasses, Snellen's metrical test types numbered D = 10, at 20 English feet, and those numbered D = 0.8 at any distance selected by the candidate himself.

Squint, inability to distinguish the principal colors, or any morbid condition, liable to the risk of aggravation or recurrence in either eye, will involve the rejection of the candidate.

The British Mercantile Marine.—An Appendix (Exn. 1, Appendix T) to the regulations relating to the Examinations of Masters and Mates in the Mercantile Marine has been issued by the Board of Trade, and came into force on September 1, 1894.

It is entitled "Form Vision, Color Vision, and Color Ignorance Tests," and enacts the following rules:

1. Examinations for Form Vision, Color Vision, and Color Ignorance are open to all persons serving or intending to serve in the Mercantile Marine, and all such persons are recommended to take the earliest opportunity of ascertaining by means of these examinations whether their vision is such as to qualify them for service in that profession.

2. The examination consists of three parts:

(*a*) Form Vision Test; (*b*) Color Vision Test; (*c*) Color Ignorance Test.

No candidate will be examined in the Color Vision Test until he has passed the Form Vision Test, or in the Color Ignorance Test until he has passed the Color Vision Test.

3. Any person serving or intending to serve in the Mercantile Marine, if desirous of undergoing the Form Vision, Color Vision, and Color Ignorance Test *only*, must make application to the Superintendent of a Mercantile Marine Office on the Form Exn. 2a, and must pay a fee of one shilling.

4. Every candidate for a Certificate of Competency who is not already in possession of such a Certificate will be required to pass the three tests mentioned in Rule 2 before he can proceed to the examination in Navigation and Seamanship for the Certificate which he desires to obtain, even though he may have passed the tests on some previous occasion.

5. Every candidate who is already in possession of a Certificate of Competency, and who desires to obtain a Certificate of a higher grade. will be required to pass the three tests mentioned in Rule 2 before he can proceed to the examination in Navigation and Seamanship for the Certificate of a higher grade.

That is to say, no candidate will be permitted to proceed with the examination in Navigation and Seamanship for a higher Certificate if he fail to pass the three tests.

6. If a candidate fail to pass any of the three tests, a note of the fact of his having done so will be written across the face of the Certificate which he already possesses before the Certificate is returned to him.

7. If a candidate who undergoes the Form Vision, Color Vision, and Color Ignorance Tests *only* (see Rule 3) be in possession of a Certificate of Competency, he must hand in his Certificate before the examination commences, and if he fail to pass any of the three tests, a statement of his failure will be written on the Certificate before it is returned to him.

8. Candidates who fail to pass the Form Vision Test or Color Ignorance Test can be re-examined at intervals of three months, but candidates who fail to pass the Color Vision Test cannot be re-examined. It is open, however, to any candidate who has failed to pass that test to appeal to the Board of Trade, who may, if they think fit, remit the case to a special examiner or body of examiners for final decision.

9. The expenses of candidates who are examined by the special examiners, *and are reported by them to have passed the three tests*, will, under certain circumstances, be paid by the Board of Trade, at a rate which will be notified to the candidate, but no payment whatever will be made towards the expenses of candidates who upon their own application are examined by the special examiners and are reported by them to have failed. The special examinations will be held in London only.

10. When a candidate fails to pass the Color Test the Examiner will point out to him the conditions under which he can appeal. Appeals are to be made through the Examiner, and forwarded to the Board of Trade with the Examiner's remarks.

11. The holder of a Certificate which bears on it a statement of failure in the first test (Form Vision) or in the third test (Color Ignorance) can have the statement removed by passing after the prescribed interval the test to which it refers, but the instruction in the last paragraph of Rule 2 must be followed.

12. The fee paid for examination for a Certificate of Competency includes the fee of One Shilling for examination in Form Vision, Color Vision, and Color Ignorance, and if the candidate fails to pass those tests, will, with the exception of One Shilling, be returned to him.

13. Only Examiners who have themselves passed the Color Test are to undertake these examinations.

Form Vision Test.—The tests to be used are Snellen's Letter Test for candidates who can read, and the dot tests for those who cannot read. The sets of tests which have been supplied to the examiners consist respectively of eight sheets of Snellen's letters and two sheets of dots.

Candidates may use both eyes or either eye when being tested, but they must not be allowed to use spectacles or glasses of any kind. If the candidate can read correctly, at a distance of sixteen feet, three of the five letters in the fifth line from the top, or four of the letters in either of the two lines below, he may be considered to have passed the test. If he cannot do so, he should be treated as having failed.

If the candidate cannot read he must be tested with the sheet of dots. For this test he is to be placed at a distance of precisely eight feet from the test sheets, and exactly opposite them. One of the sheets of dots is then to be exposed, and the candidate should be asked to name the number of dots in one or two of the lines or groups. Lines and groups of dots can be formed by holding a piece of white paper over part of the sheet, but care must be taken that when this is being done the candidate's view is not obstructed or the light on the test sheet in any way obscured.


If the candidate answer the questions put to him by the examiner with complete or very nearly complete accuracy, he should be treated as having passed. If he does not answer with very nearly complete accuracy, he should be treated as having failed.

Color Vision Test.—The Color Vision of candidates is to be tested by means of Holmgren's wools.

Color Ignorance Test.—The object of this test is simply to ascertain whether the candidate knows the names of the three colors—red, green, and white—which it is important for every seaman to be acquainted with, and the test is to be confined to naming those colors.

One or two of the purest red and green skeins should be selected from the set of wools, and the candidate should be required to name their colors. He should also be required to name the color of any white object, such as a piece of white paper.

If he answer correctly, he should be considered to have passed the test. If he make any mistake, he should be tried with the lantern which was formerly used for color tests, the plain glass and the standard red and green glasses being employed for the purpose. If he does not name these glasses correctly, he should be reported as having failed to pass the test.



Snellen's Sunrise Figure for Testing for Astigmatism (*See p.* 55.)

INDEX.

Abscission for corneal staphyloma, 170.
Abscess, cerebral, 425; corneal, 140, 161; of eyelid, 187.
Accommodation, 18, 19, 60; amplitude of, 21, 63, 65, in myopia, 44, in hypermetropia, 39; anomalies of, 64; cramp of, 48; definition of, 19; mechanism of, 19; muscle of, 49; paralysis of, 40; pupillary contraction in, 20; relative, 22; relaxation of, 73; spasm of, 40, 41, 42, 69; Tscherning's theory of, 20.
Accommodative asthenopia, 40, 41, 69, 93.
Acromegaly, 529.
Actinomyces, 234.
Acuteness of vision, 28, 32.
Adenoma of the eyelids, 197.
Agraphia, 534.
Albinismus, 91, 298, 499.
Albuminuria of pregnancy, 403, 405.
Albuminuric retinitis (*see* Retinitis).
Alexia, 533; cortical, 534; subcortical, 534.
Amaurosis, 136, 432, 434, 438; from blepharospasm, 136, 503; quinin, 413; reflex, 503; spinal, 434, 435; supra-orbital, 545.
Amaurosis, pretended, 556.
Amaurotic family idiocy, 538.
Amblyopia, 63, 413, 415, 422, 429, 433, 438, 477; alcoholic, 311; central, 408, 439; congenital, 553; ex anopsia, 476; from direct sunlight, 414; from hemorrhages, 437; from wounds of the supra-orbital nerve, 554; glycosuric, 439; in hysteria, 551; in neurasthenia, 549; in traumatic neurosis, 551; nervous, 547; of the squinting eye, 476; reflex, 553; tobacco or toxic central, 310, 429, 439; transitory, 542; uremic, 555; various forms of, 552; with eccentric fixation, 477.
Ametropia, 36, 474, 477.
Amnesia, optic, 535.
Amnesic color-blindness, 535.
Amyloid degeneration of the conjunctiva, 120.
Anemia, 519; progressive pernicious, 410.
Anesthesia of the conjunctiva and cornea, 529.
Anesthetics, local, in ophthalmology, 319.
Aneurysm, 403; arterio-venous, 509; of carotid, innominate, or aorta, 311, 411; of the internal carotid, 509; of the ophthalmic artery, 509; per anastomosis, 509; traumatic, 509.
Angle alpha, 24; gamma, 24, in hypermetropia, 39, in myopia, 45; of convergence, 24; the meter, 24; the visual, 29.
Aniridia (or Irideremia), congenital, 297; traumatic, 287.
Anisometropia, 64.
Ankyloblepharon, 229.
Anosmia, 529.
Anterior chamber, hemorrhage into, 368; late appearance of, after cataract extraction, 370.
Anterior corpus quadrigeminum, lesions of the, 531.
Antrum, maxillary, tumor in the, 511; of Highmore, tumor in the, 511; sphenoid, tumor in the, 511.
Aphakia, 369, 381, 382.
Aphasia, 553; motor, 530; visual, 534.
Apoplexy, cerebral, 471, 519.
Arcus senilis, 185.
Arecoline, hydrobromate, 318.
Argyll Robertson's operation for ectropion, 225; pupil, 538, 544.
Argyrosis, 96.
Arlt-Jaesche operation for trichiasis, 214.
Arlt's operation for cicatricial ectropion, 228.
Army, vision required for commissions in the, 562.

Asthenopia, accommodative, 40, 41; conjunctival, 93; muscular, 494; nervous, 547; retinal, 477.
Asthma, 106.
Astigmatism, 36, 50, 51, 52, 85, 383; after cataract operations, 383; corneal, 64; determination of, by ophthalmoscope, 79, by retinoscopy, 60; estimation of degree of, 57, by astigmometer, 60; headache due to, 56; hypermetropic, compound, 54, 59, simple, 54, 58, 192; irregular, 64, 174; lental, 63, dynamic or active, 63, static or passive, 63; mixed, 54, 60, 85; myopic compound, 54, 59; simple, 54, 58; regular, 52, 54; spectacles in, 57; symptoms of, 55.
Astigmometer, the, 58, 61.
Ataxic paraplegia, 499, footnote.
Ataxy, hereditary, 545; locomotor, 434, 462, 463; of the orbital muscles, 543.
Atheroma, 403, 411, 412.
Athetosis, 531.
Atrophy of the optic nerve (*see* Optic nerve).
Atropin, 40, 60, 63, 68, 69, 73, 84, 102, 117, 138, 144, 149, 150, 152, 155, 158, 162, 165, 166, 168, 176, 180, 182, 187, 251, 256, 268, 277, 284, 286, 287, 289, 309, 316, 335, 338, 342, 347, 354, 369, 372, 375, 377, 378, 396, 485, 554; danger of glaucoma from use of, 257; poisoning by, 257
Axis, optic, 24.

Bader's operation for conical cornea, 175.
Base of skull, fracture of, 127.
Basedow's disease, 517.
Bergeon's treatment of rodent ulcer of the eyelid, 191.
Berlin's operation for entropion, 220.
Binocular vision, 64.
Birnbacher's operation for ptosis, 201.
Bisulphid of carbon, poisoning by, 431.
Black spot in myopia, 47.
Blennorrhea, acute, of conjunctiva, 97, 126, 141; chronic of conjunctiva, 98; neonatorum, 97; of the lacrimal sac, 237.
Blepharitis, 94, 356; chronic, 211; intermarginal, 94; marginal, 191; squamosa, 191; ulcerosa, 191.
Blepharophimosis, 210.
Blepharoptosis, 198.
Blepharospasm, 134, 135, 141, 197.
Blind spot, the, 34, 46.
Blows on the eye, 180.
Bowman's operation for conical cornea, 175.
Brain, basic lesions of the, 447; diagnosis of, 468; focal disease of the, 522, pontine lesions of the, 447.
Bright's Disease, 355, 403, 405, 411, 413, 428.
Broca's lobe, lesion of, 535.
Bronchitis, 106, 264.
Bulbar paralysis, 69.
Bulla of the cornea, 123, 180.
Buphthalmos, 340.

Canaliculus, obstruction of the, 234.
Canthoplastic operation, 211.
Capsule, lesion of the internal, 530.
Capsulotomy, 379; Bowman's method of, 380; Knapp's method of, 381; Noyes' method of, 380.
Carcinoma, 508; of chorioid, 294; of ciliary body, 292; of orbit, 508.
Caries of nasal bones, 241; of the orbit, 503; of the teeth, 553.
Caruncle, sinking of the, 490, 493.
Cataract, 42, 419; membranous, 344; black, 345, 349; accreta, 352; artificial ripening of, 348; calcareous, 352; capsular, 353; central capsular, 104, 353; central lental, 350; complete, 342; complete congenital, 349; complete, of young people, 349; congenital, 262; diabetic, 349; discission, or dilaceration of, 376; fusiform, or spindle-shaped, 351; glaucoma after extraction of, 375; linear extraction for, 357; mental derangement after extraction of, 375; modified peripheral linear extraction of, 360; Morgagnian, 344; myopia in incipient, 45; combined, or three-millimeter flap, operation for, 362; or extraction of, with iridectomy, 362; operation for simple, without iridectomy, 373; pathogenesis of, 342; polar, anterior, or pyramidal, 104, 351; posterior, 352; ripeness of, 343; secondary, and its operation, 379; total, 352; senile, 342; spectacles after extraction of, 382, in in-

cipient, 347; spontaneous cure of, 344; suction operation for, 378; symptoms of, 346; traumatic, 353; von Graefe's operation for, 360; zonular, or lamellar, 350.
Cautery, use of the actual, 145, 151, 153, 155, 176.
Cavernous sinus, thrombosis of the, 312, 469, 501.
Cellulitis, 412, 501, 505.
Cerebellum, tumor of the, 470.
Cerebral abscess, 425; cysts, 425; embolism, 310; hemorrhage, 530; hemispheres, tumors of the, 469; localization, 470; peduncle, lesions in the, 470.
Cerebral tumors, 310, 312, 429; ophthalmoscopic appearances in, 424; the pupil in, 310.
Cerebritis, 426.
Chalazion, 194.
Chemosis, 93, 98, 173, 469.
Chiasma, lesion of the, 522, 528.
Chlorosis, 427, 519.
Choked disc, 424.
Cholera, Asiatic, 504.
Cholesterin in the vitreous humor, 387.
Chorea, 411, 542.
Chorioid, central senile areolar atrophy of the, 296; coloboma of the, 499; detachment of the, 295, 373; extravasation of the blood in the, 288; foreign bodies in the, 288; inflammation of the, 261; injuries of the, 288; miliary tubercles in the, 540; rupture of the, 288; sarcoma of the, 292; staphyloma of the, 295.
Chorioidal atrophy, general, 47, 409; carcinoma, 294; degeneration, near macula lutea, 46; exudation, near macula lutea, 47; ring, 88; osteosarcoma, 294; sarcoma carcinomatosum, 294; tubercle, 294.
Chorioiditis, 189, 289; central, 263, senile guttate, 262, 356; disseminated or exudative, 261, 352, 355, 401; embolic, 264; posterior sclero-, 295; purulent, 263, 328; syphilitic, 262.
Chorioido-retinitis, chronic, 295.
Chromidrosis, palpebral, 195.
Cicatrix, cystoid, 285, 333, 372.
Ciliary body, carcinoma of the, 292; inflammation of the, 259; injuries of the, 287; myosarcoma of the, 292; sarcoma of the, 292.
Ciliary muscle, 19, 66; cramp of the, 38, 39, 40.
Ciliary neuralgia, 112, 293, 329.
Circle of diffusion, 19.
Circumlental space, 330.
Civil Service, vision required for the, 562.
Climacteric period, 410.
Cocain, 309, 319.
Coloboma of the chorioid, 298, 499; of the eyelid, 232; of the iris, 297.
Color-blindness, 26, 27, 28, 435, 439, 535; amnestic, 535.
Color-sense, 25; theories of the, 25, 26, 27.
Commotio retinæ, 422.
Concave lens, 170.
Concussion, injury to optic nerve by, 503.
Condyloma of the iris, 254.
Congestion papilla, 424.
Conical cornea, 173.
Conjugate deviation of eyes, 469.
Conjunctiva, amyloid degeneration of the, 121; anesthesia of the, 529; burn of the, 132; cyst of the, 129; cysticercus under the, 130; dermoid of the, 128; diseases of the, 95; ecchymosis under the, 127; epithelioma of the, 129; essential shrinking of the, 124; hemorrhage in the, 120; hyaline degeneration of the, 120; hyperemia of the, 93; injuries of the, 131; lipoma of the, 128; lithiasis of the, 130; lupus of the, 122; lymphoma of the, 117; nevus of the, 127; papilloma of the, 128; pemphigus of the, 122; pinguecula of the, 127; polypus of the, 127; sarcoma of the, 129; syphilitic disease of the, 128; tubercle of the, 121; xerosis of the, 123.
Conjunctival complication of smallpox, 119.
Conjunctivitis, 93, 356, 503, 540, 554; catarrhal or simple acute, 94; croupous, 117, 123; diphtheritic, 116, 118, 123, 141; follicular, 104, 257; gonorrheal, 97; granular, 107; petrificans, 130; phlyctenular, 133; purulent, 111, 150; spring catarrh, 105.

Constabulary, Royal Irish, vision required for the, 563.
Contact glasses, 175.
Contraction, consensual, of pupil, 304.
Convergence, angle of, 24; and accommodation, connection between, 22.
Convex lenses, 17, 52.
Convulsions, 478.
Corectopia, 297.
Cornea, 17; abrasions of the, 179; abscess of the, 141, 161; abcission of the, 170; absorption ulcer of the, 155; anesthesia of the, 326, 518, 529; arcus senilis of the, 185; bullæ of the, 180; burns of the, 181; calcareous film of the, 168; cautery in treatment of ulcers of the, 154; conical, 42, 173; cysts of the, 177; dermoid of the, 177; diseases of the, 140, disjunction of the, 179; ectasies of the, 169; epithelioma of the, 177; faceted ulcer of the, 156; fibroma of the, 177; foreign bodies in the, 177; globosa, 340; herpes of the, 156; infantile ulceration of the, with xerosis of the conjunctiva, 15; inflammations of the, 140; injuries of the, 177; leukoma of the, 183; macula of the, 183; nebula of the, 183; opacities of the, 477; paracentesis of the, 147, 150; perforating injuries of the, 181, 339; perforation of the, 144; phlyctenular disease of the, 133; recurrent abrasion of the, 179; ring ulcer of the, 154; sclerotizing opacity of the, 168, 246; staphyloma of the, 169; tattooing of the, 184; transplantation of the, 184; transverse calcareous film of the, 168; tumors of the, 177; ulcer, absorption of the, 155; ulcer, deep, of the, 150; ulcer, faceted, of the, 155; ulcer, rodent, of the, 153; serpiginous ulcer of the, 151; simple ulcer of the, 149; superficial transparent ulcer of the, 155; ulceration of the, 143, 145, 418, 540.
Corneal deposits, punctate, 253, 271, 272, 273.
Corona radiata, lesions of the, 447.
Corpora quadrigemina, lesions of the, 304.
Cramp, clonic, of orbicularis, 197; tonic, of orbicularis, 197; of ciliary muscle, 39.
Crus cerebri, lesion of the, 467, 468.
Crystalline lens, 17, 20, 51, 64, 148; absence of the, 382; disease of the, 342; dislocation of the, 381.
Cyclitis, 188, 259, 288, 362, 417, 419; plastic, 260; primary idiopathic, 260; purulent, 261.
Cylindrical lenses, 57.
Cyst, cerebral, 425, 529; of cornea, 177; of eyelid, 232; of iris, 289; orbital, 507; simple, of conjunctiva, 129.
Cysticercus, subconjunctival, 130; subretinal, 417.

Dacryoadenitis, 242.
Dacryocystitis, 153, 356; acute, 117, 241; chronic, 152, 237; phlegmonous, 241.
Dacryolith, 234.
Daturin, 309.
Davidson's, Mackenzie, method for employing Röntgen rays, 390.
Decentration of spectacle glasses, 67, 497.
Delirium tremens, 431.
Dermoid tumors of the conjunctiva and cornea, 128, 262; of the orbit, 507.
Diabetes, 69, 256, 270, 274, 355, 431, 439, 461.
Dianoux' operation for trichiasis, 215.
Dinitro-benzol, poisoning by, 431.
Dioptric unit, 17, media, 17, system, 18.
Dioptry, the, 17.
Diphtheria, 68, 119, 461.
Diphtheritic conjunctivitis, 118.
Diplopia, 126, 446, 452, 463, 487, 506; crossed, 449, 452, 454, 456, 476, 555; homonymous, 449, 450, 452, 456, 475; in concomitant convergent strabismus, 476; in insufficiency of the internal recti, 494; monocular, 286; test, 556.
Direct ophthalmoscopic method, 72.
Disc, optic (*see* Optic papilla).
Discission or dilaceration of lens, 376.
Disseminated sclerosis (*see* Sclerosis).
Distichiasis, 111, 211.
Duboisin, 309.
Dyslexia, 534.

Eccentric vision, 30.
Ecchymosis of the conjunctiva, 127.
Eclipses, blinding of the retina in, 414.
Ectropium, 223; cicatricial, 223, 228; muscular or spastic, 223; of the lid, 223, 503; senile, 224; operation for, 224.
Eczema, 136, 138, 187.
Edema of eyelids, 97, 197; of orbital tissues, 505.
Electric light, effect of, on the eyes, 415.
Electrolysis, 127, 212, 237, 422; for nevi of the eyelids, 195; for trichiasis, 212; in detachment of retina, 422.
Electro-magnets, 394.
Elephantiasis lymphangioides of lids, 197.
Embolism, cerebral, 527, 542; of retinal vessels, 400, 410, 411.
Emmetropia, 18.
Encephalocele, 507, 512.
Encephalopathia saturnia, 541.
Endarteritis, of retinal vessels, 413.
Endocarditis, 104, 264, 411.
Enophthalmos, 504.
Entropion, 111, 123, 211, 218; senile, 218, 221; spastic, 218.
Enucleation, 122, 171, 278, 513.
Ephedrin, 317.
Epicanthus, 231.
Epilepsy, 56, 127, 310, 312, 433, 494, 541.
Epiphora, 233, 238, 545.
Episcleritis, 244, 245.
Epithelioma of the conjunctiva, 129; of the cornea, 177; of the eyelid, 197.
Erysipelas of the eyelids, etc., 187, 197, 241, 412, 426, 469, 501, 503.
Erythema, 187.
Erythropsia, 383, 557.
Eserin (or Physostigmin), 137, 309, 318, 337, 362.
Ethmoid cells, enchondromata in, 511; fibromata in, 511; tumors of, 510, 511.
Eucain B, 319.
Euphthalmin, 317.
Everbusch's operation for congenital ptosis, 204.
Evisceration of the eyeball, 171, 275, 278.
Excision of the eyeball (*see* Enucleation).
Exophthalmic goiter, 515, 517; Dalrymple's sign in, 518; Stellwag's sign in, 518; von Graefe's sign in, 517.
Exophthalmos, 245, 469, 501, 502, 503, 504, 507, 512; caused by bleeding into the capsule of Tenon, 493; monolateral, 505; pulsating, 509.
Exostosis of the orbit, 508, 510, 529; in the sphenoid antrum, 511.
External geniculate body, lesion of the, 531; recti, insufficiency of the, 474, 489; rectus, advancement of the, 481, 488, 491.
Eyeballs, motions of, and their derangement, 441.
Eyelashes, lice on the, 195.
Eyelid, adenoma of the, 197; chromidrosis of the, 195; congenital coloboma of the, 232; cramp of the, 197; diseases of the, 187; ecchymosis of the, 231; ectropion of the, 223, eczema of the, 187; elephantiasis of the, 197; emphysema of the, 231; entropion of the, 218; epithelioma of the, 197; herpes zoster of the, 187; injuries of the, 230; inversion of the, 218; lupus of the, 197; milium of the, 194; molluscum of the, 195; nevi of the, 195; restoration of the, 229; rodent ulcer of the, 190; sarcoma of the, 197; stye on the, 193; syphilitic sores of the 189; vaccine vesicles on the, 190.

Far point, 18, 20.
Fascicular paralysis, 463.
Fatigue field, 548, 550; in traumatic neurosis, 552.
Field of vision, 30; binocular, 34; dimensions of, 30; for colors, 34; in glaucoma, 324, 325; in toxic amblyopia, 429; in optic atrophy, 435; in functional neurosis, 547.
Fifth nerve, influence of the, on the pupil, 307; paralysis of the, 471.
Fixation, field of, 446; line of, 24; object of, 30; point of, 24.
Fluorescin, 142, 147.
Focal disease of brain, 522; illumination, 87, 141; interval, 53; length of trial lenses, 17.

Focus, conjugate, 19; principal, of the eye, 18.
Foramen, fracture of the margin of the optic, 503.
Foreign bodies in the cornea, 177; in the interior of the eye, 388; in the iris, 285.
Form sense, 28.
Fourth nerve, paralysis of the, 469.
Friedreich's disease, 545.
Frontal sinus, osteoma of the, 510; tumors of the, 510.
Fundus oculi, the, 88, 90.

Gaillard's sutures, 221.
Galvano-cautery, 137, 372, 435, 457.
Gamma, angle, 24.
Ganglia, primary optic, lesions of the, 531.
Ganglion, ciliary, 306; Gasserian 307, 308; hebenulæ, 305.
General paralysis of the insane, 538, 434.
Geniculate body, lesion of the external, 305.
Glaucoma, 257, 289, 320, 403, 410, 548; absolute, 159, 352; acute, 163, 329; etiology of, 329; fulminans, 328; hemorrhagic, 340, 410; pathology of, 330; primary, 320, 321, 326; secondary, 255, 285, 290, 320, 339, 354, 378, 417; treatment of, by iridectomy, 332, by myotics, 338, by posterior sclerotomy, 333, by sclerotomy, 336, by sympathectomy, 337.
Glaucomatous attack, 327; cup, 322; degeneration, 328; ring, 325.
Glioma of the retina, 416; pseudo-, 263, 384, 417.
Globulin, 353.
Goiter, exophthalmic, 325, 463, 517.
Gonorrheal arthritis, 104; ophthalmia, 97.
Gout, 245, 249, 270.
von Graefe's operation for conical cornea, 175, for senile entropion, 222; sign in exophthalmic goiter, 517; test for insufficiency of the internal recti, 495.
Granular conjunctivitis (*see* Trachoma).
Granuloma of the iris, 194, 289.
Graves' Disease, 505, 517.
Green's operation for entropium, 219.

Hallucinations, visual, 535, 550; in connection with glaucoma, 535.
Harlan's operation for symblepharon, 209.
Hay fever, 106.
Helmholtz's ophthalmometer, 60; ophthalmoscope, 71.
Hematemesis, 410, 438.
Hemiachromatopsia, 525, 529, 530.
Hemianesthesia, 468.
Hemianopic prism phenomenon, 532; pupil, 532.
Hemianopsia, 425, 439, 462, 466, 469, 522; absolute, 523; altitudinal, 523, 529, 553; bitemporal, 511, 528, 529; complete, 523; cortical, 530; homonymous, 304, 523, 529; in complete or partial, 523; localization of the lesion, in cases of, 528; nasal, 523, 529; relative, 523; superior and inferior, 523; temporal, 523; transitory, 552.
Hemiatrophy, facial, 504.
Hemiplegia, 462, 468, 469, 470, 471; crossed, 468.
Hering's drop experiment, 486, footnote; theory of the color-sense, 25.
Hernia cerebri, 512.
Herpes corneæ febrilis, 156; herpes zoster ophthalmicus, 156, 187.
Heterophthalmos, 297.
Hippus, 305, 308, 313.
Holmgren's method for testing the color-sense, 559.
Holocain, 319.
Homatropin, 317.
Hemophthalmos, 248.
Hemorrhage, cerebral, 530; from stomach, bowels, or uterus, 437, 519; in anterior chamber, 181; in retina, 51, 121, 400, 633; at yellow spot, 48; in vitreous humor, 386, 398.
Hooping-cough, 127, 158, 264.
Hordeolum, 193.
Hot eye, 244.
Hotz's operation for senile entropium, 222.
Hyaline degeneration of the conjunctiva, 121.
Hyaloid artery, persistent, 397.
Hydrocephalus, 426, 433, 434, 462, 540.
Hydrophthalmos, congenital, 340.
Hyoscyamin, 309.
Hyperemia of the papilla, 88.

Hyperesthesia of the retina, 553.
Hyperbolic lenses, 174.
Hypermetropia, 36, 52, 192, 475; accommodative asthenopia in, 40; amplitude of accommodation in, 39; angle gamma in, 39; axial, 36; estimation of, by ophthalmoscope, 77; by retinoscopy, 80; cause of internal strabismus in, 474; convergent concomitant strabismus in, 41; correction of, 37; cramp of ciliary muscle in, 39; curvature, 36; latent, 40; manifest, 38, 40; spectacles in, 41; the pupil in, 303.
Hyperostosis of the orbit, 529; in the sphenoid antrum, 511.
Hyphema, 248, 284.
Hypopyon, 135, 138, 143, 145, 259, 393.
Hysteria, 519, 547.

Idiocy, amaurotic family, 538.
Illaqueatio, 212.
Image, inverted ophthalmoscopic, 73; upright opthalmoscopic, 72.
Influenza, 68, 429, 432, 461, 501, 545.
Insomnia, 550.
Insufficiency of convergence, 494.
Intermittent fever, 158.
Internal capsule, lesion of the, 471, 530, 531; recti, insufficiency of the, 48, 67; rectus, tenotomy of the, 481, 488.
Intra-ocular media, 18.
Iodoform poisoning, 431.
Iridectomy, 300, 376; for glaucoma, 332, 338; in cataract operations, 361, in sympathetic ophthalmitis, 282; instruments for, 300; optical, 175.
Irideremia, 297.
Iridis, rupture of the sphincter, 286.
Iridochorioiditis, 120, 255, 289; plastic, 540.
Iridocyclitis, 120, 151, 159, 250, 255, 293, 330, 340, 373, 379, 398; chronic, 418.
Iridodialysis, 285.
Iridodonesis, 381.
Iridoplegia, 312.
Iridotomy, 302, 380.
Iris, absence of the, 287; coloboma of the, 297; cysts of the, 289; dehiscence of the, 286; foreign bodies in the, 285; granuloma of the, 289; injuries of the, 284; malformations of the, 297; operations on the, 300; persistent pupillary membrane of the, 297; prolapse of the, 134, 137, 148, 182; retroflexion of the, 286; rupture of sphincter of, 286; sarcoma of, 291; tubercle of the, 289.
Iritis, 94, 95, 102, 112, 120, 135, 138, 144, 151, 189, 372, 378, 405, 417, 419; gonorrheal, 253; purulent, 259; quiet, 252; rheumatic, 253; serous, 253; symptoms of, 251; syphilitic, 253, 254, 259; tubercular, 289.
Irritation, sympathetic, 265.

Jacob's ulcer, 190.
Japanese muff-warmer, 145, 165.
Javal's orthoptic treatment for strabismus, 485.
Jequirity, 115.

Keratitis, 120; aspergillina, 161; bullous, 159; dendriform, 160; diffuse interstitial, or parenchymatous, 162, 540; fascicular, 134; filamentary, 159; grating-like, 166; guttate, 167; neuroparalytic, 155; phlyctenular, 126, 135, 141; profunda, 165; punctata, 167, 253; purulent, 264; riband-like, 168; striped, 370.
Keratoconus, 173.
Klebs-Löffler bacillus, 118, 119.
Knapp's operation for tumor of optic nerve, 437.
Koch-Weeks' bacillus, 95, 98.
Krönlein's operation, 515.
Kuhnt's operation for extirpation of lacrimal sac, 239; for ectropion, 224.

Lacrimal apparatus, diseases of the, 233; canaliculus, obstruction of the, 234; duct, stricture of the, 236; fistula, 239.
Lacrimal gland, extirpation of the, 243; hypertrophy of the, 242; inflammation of the, 242; removal of the, 241, 243; tumors of the, 512.
Lacrimal punctum, inversion of the, 233; stenosis and complete occlusion of the, 233.
Lacrimal sac, lupus of the, 122; blennorrhea of the, or chronic dacry-

ocystitis, 237; extirpation of, 239; obliteration of, 239.
Lagophthalmos, 207, 471.
Lamellar, or zonular, cataract, 350.
Lamina cribrosa, 89, 323, 437.
Landry's disease, 546.
Lateral sclerosis, 434.
Lead poisoning, 428, 461.
Left ventricle, hypertrophy of the, 410.
Lens, crystalline, discission of, 51, 282, 349, 350; dislocation of, 339, 352; extraction of, 282; injury of, 340; lateral displacement of, 339; sclerosis of, 52, 255.
Lenses, concave, 17; convex, 17; cylindrical, 57; numbering of trial and spectacle, 17.
Lental astigmatism, 63.
Lenticular nucleus, lesions of the, 533.
Lenticonus, 382.
Leptothrix in the canaliculus, 234.
Leukemia, 405, 509.
Leukoma of the cornea, 144, 183; adherent, 103, 150, 151; tattooing for, 184.
Lice on the eyelashes, 193.
Light difference, 25; minimum, 25; sense, 25.
Lipoma of the conjunctiva, 128.
Lithiasis of the conjunctiva, 130.
Locomotor ataxy (*see* Tabes dorsalis).
Lupus of the conjunctiva, 122; of the eyelids, 197; of the lacrimal sac, 122.
Lymphadenoma of the orbit, 505, 508, 509.

Macula Cornea, 144, 183, 449, 452, 475.
Macula lutea, 18, 48, 76, 84, 89, 401, 402, 406, 408, 410, 411, 475; exudation at the, 263.
Macula, Trousseau's cerebral, 519.
Macular center in cortex, theory of arrangement of, 527.
Madarosis, 190.
Maddox's rod-test, 496.
Magnet, the, for removing foreign bodies, 394.
Malaria, 245, 400, 429.
Malingering, tests for, 555.
Mania, acute, 311, 312.
Marasmus, 539.
Massage of eyeball, 412.
Maxwell's operation for shrunken sockets, 513.
Measles, 95, 134, 162, 264, 429, 478.
Media, dioptic, intra-ocular, or refracting, 18.
Megalopsia, 262.
Meibomian cyst, 194; glands, 94, 194.
Melancholia, 312.
Meningitis, 104, 279, 425, 433, 457, 469; acute, 539; acute tubercular, 540; basal, 529; cerebro-spinal, 264, 425, 540; purulent, 501; spinal, 312; syphilitic, 533; traumatic, 540; tubercular, 425.
Meningocele, 513.
Menstruation, suppression or irregularity of, 410, 426.
Mental derangement after cataract extraction, 375.
Mercantile marine, vision required for, 563.
Metamorphopsia, 262, 401, 415.
Meter angle, 24; lens, 17.
Metria, 407, 501.
Metrical system for spectacle lenses, 17.
Meynert's fibers, 532.
Microphthalmos, 499.
Micropsia, 68, 262, 401.
Migraine, 532, 542.
van Millingen's operation for trichiasis, 217.
Milium, 194.
Mind-blindness, 531, 534, 535, 538.
Mitral disease, 411.
Molluscum, 195.
Morax, diplobacillus of, 95.
Morton's operation for conical cornea, 176.
Morvan's disease, 546.
Mouches volantes, 386.
Mucocele of the lacrimal sac, 238, 510.
Mules' operation for corneal staphyloma, etc., 172, 278, 339; for ptosis, 200.
Muscæ volitantes, 386.
Muscarin, 309.
Muscles of the orbit and their derangements, 441; cerebral, paralysis of the, 464; conjugate lateral paralysis of the, 464; paralysis of the, 447; normal action of the, 441.
Muscular atrophy, progressive, 462.

Myasthenia gravis, 463.
Mydriasis, 67, 68, 311, 455; irritation, 311; paralytic, 311; traumatic, 286, 287.
Mydriatics, action of the, 309.
Mydrin, 317.
Myelitis, acute, 428, 545; cervical, 311; pupillary symptoms in, 546.
Myodesopsia, 386.
Myopia, 36, 42; amplitude of accommodation in, 44; angle gamma in, 44; apparent, 39; axial, 42; cause of, 45; complicated with organic disease, 46, 419; curvature, 42; detachment of the retina in, 48, 419; determination of degree of, 44; functional anomalies of, 48; management of, 48; operative cure of, 51; pernicious, 46; prescribing of glasses in, 48; progressive, 46.
Myopic astigmatism, 58; crescent, 46.
Myosarcoma of ciliary body, 292.
Myosis, paralytic, 310; traumatic, 287.
Myotics, action of the, 309.
Myotonia congenita (Thomson's), 546.
Myxo-sarcoma of the optic nerve, 435.

Nevus of the conjunctiva, 127; of the eyelid, 195.
Nasal catarrh, 136, 235, 237; duct, stricture of the, 235.
Navy, vision required for the, 562.
Near objects, 18, 19; point, 21, 64, 66.
Nebula of cornea, 144, 183.
Neisser, gonococcus of, 97.
Nephritis, acute, 403, 405; chronic, 403; of pregnancy, 555.
Nerve fibers, opaque, 89.
Neurasthenia, 313; nervous amblyopia in, 549.
Neurectomy, optico-ciliary, 275, 278, 280, 338.
Neuritis, optic, 501, 529, 541, with persistent dropping from nostril, 432; peripheral, 428, 463, 544; retrobulbar, 427, 432.
Neuro-paralytic keratitis, 155.
Neuro-retinitis, 401.
Neurosis, traumatic, 546, 547; the fatigue field in, 552.
Neurotomy, optic-ciliary, 278, 281.
Night-blindness, 125, 262, 401, 554.
Noyes' operation for secondary cataract, 380.
Nuclear paralysis, 460; acute, 461; chronic, 461.
Nuclear pupil-contracting center, paralysis of, 309.
Nucleus, lenticular, tumor of, 533.
Nystagmus, 298, 349, 499, 537.

Oblique illumination, 87.
Occipital lobe, lesion of, 530, 534.
Ocular diseases and symptoms caused by focal brain disease, 522.
O'Farrell's enucleation, 278.
Ointment, yellow oxid of mercury, 137.
Omphalophlebitis, 264.
Opacities in refracting media, 87.
Opaque nerve fibers, 89.
Opthalmia, chronic, 109, 155, 211, 218; diphtheritic, 124; Egyptian, 109; electric, 415; gonorrheal, 97; granular, 107; migratoria, 265; military, 107; neonatorum, 97; neuro-paralytic, 471; nodosa, 291; phlyctenular, 133, 155; purulent, 97; strumous, 133.
Ophthalmitis, sympathetic, 248, 252, 260, 267, 288, 289, 372, 393; prophylactic operations in, 278, 415; tarsi, 191.
Ophthalmometer, 60.
Ophthalmoplegia, acute hemorrhagic, 461; externa, 460, 463; interna, 67, 460; universa, 460.
Ophthalmoplegic migraine, 459.
Ophthalmoscope, 57, 58, 70, 482; Helmholtz's, 71; modern, 71; direct method, 72; indirect method, 73.
Optic amnesia, 535; atrophy, 251, 312, 355, 405, 413, 414, 433, 435, 438, 439, 506, 538, 548; axis, 24; center, 24; chiasma, 303.
Optic axis, 18, 484; center, 18, 613.
Optic disc (or papilla), 34, 46, 75, 88; atrophy of the, 426, 428, 431; hyaline or colloid outgrowths from the, 437; glaucomatous cupping the, 89, 322; physiological cupping of the, 89, 322.
Optic ganglia, lesions of the primary, 531.
Optic nerve, 75; atrophy of the, 28, 501; atrophy of the, in tabes dor-

salis, 543; diseases of, 424; injuries of, 437; tumors of, 436.
Optic neuritis, 89, 251, 265, 271, 424, 432, 433, 506, 529, 540; retrobulbar, 432; with persistent dropping from nostril, 432.
Optic radiations, lesions of the, 530, 531; tract, disease of, 531, 533.
Orbicular sign, 471.
Orbicularis palpebrarum, paralysis of the, 462.
Orbit, aneurysm of the, 509; carcinoma and sarcoma of the, 508; caries of the, 503; cellulitis of the, 501; cysts, hydatid, of the, 507; derangement of vision in tumors of the, 506; diseases of the, 501; diseases of neighboring cavities, 509; exostosis of the, 508; injuries of the, 503; lymphadenoma of the, 508, 509; palpation of the, 505; perforating injuries of the, 504; periostatis of the, 504; shrinking of cavity of the, after enucleation of eyeball, 513; symmetrical tumors of the, 509; temporary resection of outer wall of the (Krönlein's operation), 515; tumors of the, 426.
Orbital cysts, 507; cellutis, 493, 501,
Orbital muscles, ataxy of, 543, coordinating centers for, 499; relationship of epilepsy to want of power of the, 494; paralysis of, 502, 538, 543, 637.
Orbital nerves, paralysis of, 529; periostitis, 503; tissues, edema of, 505.
Orbital tumors. 504; diagnosis of the nature of, 507.
Orbital wall, fracture of, 504.
Orientation, 551.
Osteoma of the frontal sinus, 510; of the sphenoidal antrum, 511.
Otitis, purulent, 469.
Ozena, 152, 238.

Pagenstecher's operation for ptosis, 200.
Panas' operation for ptosis, 202.
Pannus, 111, 121, 129, 518.
Panophthalmitis, 120, 151, 156, 182, 248, 263, 268, 371, 384, 632.
Papillitis, 400, 404, 424; sympathetic, 274.
Papilloma of conjunctiva, 128.
Paracentesis, 258, 411, 412; of corneal ulcer, 102, 137, 145, 147, 150.
Parallax, 89.
Parallax rays, 18.
Paralysis, acute ascending (Landry's disease), 546; agitans, 541; bulbar, 311, 461, 462; chronic, 462; conjugate lateral, 462; crossed, 470; fascicular, 463; general, of the insane, 311, 312, 435, 463, 538; infantile, 633; nuclear, 460; acute nuclear, 461; chronic nuclear, 461; accommodation, 40, 67, 455, 454; of cervical sympathetic, 504; of external rectus, 68, 450; of the fifth nerve, 471; of fourth nerve, 470; of levator palpebræ (*see* Ptosis); of Müller's muscle, 504; of orbital muscles or nerves, 502, 529, 538, 543, 551; of seventh nerve, 471; of sixth nerve, 470; of sphincter of iris, 68, 545; of superior oblique, 451; of third nerve, 68, 454, 466; of third nerve, intermittent, of one eye, 459; passive orthoptic treatment of orbital, 458; rheumatic, 457.
Paralytic mydriasis, 311; myosis, 309.
Parasitic disease of the retina, 417.
Paresis, latent, of an ocular muscle, method for discovering, 597.
Parkinson's disease, 541.
Pemphigus of the conjunctiva, 122, 124; vulgaris, 123.
Pericarditis, 104.
Perimeter, 30, 446, 483, 484, 548, 552.
Periostitis, 426, 508, 529; chronic, in persons of a rheumatic tendency, 502; gummatous,, 501; of the orbit, 502, 503; syphilitic, 68.
Peritomy, 117.
Perivasculitis, 413.
Petrous portion of temporal bone, fracture of apex of, 470.
Phlegmonous inflammation of the eyelids, 187.
Phlyctenula, 120, 129, 134.
Phlyctenulæ, multiple or miliary, 134.
Phlyctenular, or strumous conjunctivitis, or keratitis, 133.
Phosphenes, 419.
Photometer, 25.
Photophobia, 93, 106, 108, 112, 123, 134, 135, 139, 540.

Photopsia, 550.
Phtheiriasis ciliorum, 193.
Phthisis bulbi, 248, 255, 293, 328, 418, 419, 508.
Physostigmin (*see* Eserin).
Pilocarpin, 309, 318, 337.
Pineal gland, tumor of the, 469.
Pinguecula, 126, 127.
Pituitary body, tumor of the, 529.
Pleuritis, 104.
Pneumococcus, 95.
Pneumonia, 158, 256.
Poliencephalitis, chronic superior, 461.
Poliomyelitis, acute anterior, 461.
Polycoria, 297.
Polyopia, monocular, 346.
Polypi in the sphenoidal antrum, 511.
Polypus of the conjunctiva, 127.
Polyuria, 511.
Pons, disease of the, 470, 471.
Posterior staphyloma, 46.
Pregnancy, 411.
Presbyopia, 64.
Prisms, 458.
Progressive muscular atrophy, 311.
Proptosis, 436, 502, 509, 518, 529; paralytic, 505.
Pseudo-cornea, 170; -glioma, 384, 417; -leukemia, 509.
Pterygium, 125.
Ptosis, 109, 198, 231, 454, 457, 460, 468; cerebral, 466; congenital, 203, 460; monolateral, 467; operations for, 200, 201; pseudo-, 467; sympathetic, 545; with associated movements of eyelid, 206.
Pulvinar, 531, lesion of, 530.
Punctum lacrimale, malposition of the, 233; stenosis of the, 233.
Punctum proximum, 20, 45, 65; remotum, 20, 43, 45.
Pupil, the, 20, 552; action of mydriatics on the, 309; action of myotics on the, 309; Argyll Robertson, 310, 538, 544; artificial, 175, 184, 381; change of, in accommodation, 20, 303; contraction of the, 303; dilatation of the, 306; exclusion of the, 251; hemianopic, the, 532; hippus, or unrest, of the, 308; in chloroform narcosis, 309; inequality of the, 546; malposition of the, 297; motions of the, in health and disease, 303; occlusion of the, 251; reflex mobility of the, 305; size of the, in disease, 309; supernumerary, 297; symptom, paradoxical, 538.
Pupillary alterations in tabes dorsalis, 544; light reflex, 542; membrane, persistent, 397; symptoms in myelitis, 546.
Pyemia, 264.

Quinin amaurosis, 413.

Railway spine (*see* Traumatic neurosis).
Recti, insufficiency of the internal, 494.
Recurrent fever, 256.
Red vision, 551.
Refracting surfaces of the eye, 17.
Refraction and accommodation of the eye, abnormal, 36; normal, 17.
Refraction, estimation of, by ophthalmoscope, 76; by retinoscopy, 80.
Regurgitation, aortic, 325.
Retina, alterations in vascularity of the, 400; anemia of the, 400; aneurysm of central artery of the, 413; apoplexy of the, 386, 409, 410, 413; atrophy of the, 403, 407; blinding of the, by direct sunlight, 414; by electric light, 415; cysticercus under the, 417; detachment of the, 48, 51, 255, 288, 352, 355, 369, 386, 395, 397, 404, 410, 418; development of connective tissue in the, 406; diseases of the, 400; embolism of the central artery of the, 410, 434; exudations in the, 400; glioma of the, 416; gyrate atrophy of the, 409; hemorrhages of the, 400, 406; hyperemia of the, 400; hyperesthesia of the, 553; inflammations of the, 398; injury of the, 414; ischemia of the, 414; normal, 89; parasitic disease of the, 417; sclerosis of vessels of the, 413; spasm of vessels of the, 400; thrombosis of artery of the, 400; thrombosis of vein of the, 400; traumatic anesthesia of the, 422; traumatic edema of the, 422; tumor of the, 416.
Retinal affections in diabetes, 405; artery, thrombosis of the, 412, 413; asthenopia, 477; detachment, causes of, 419; hemorrhages, 541; ische-

mia, 414; vein, thrombosis of the, 412; vessels, 91; vessels, sclerosis of, 413.
Retinitis, 251, 400, 540; albuminuric, 400, 402, 403, 424, 555; chorioido-, 401, 434; circinata, 406; hemorrhagic, 402, 405; leukemic, 405; neuro-, 401; ophthalmoscopic appearances in, 404; pigmentosa, 352, 406, 407, 434, 554; proliferans, 406; punctata albescens, 262, 406; purulent, 407; specific, 386; syphilitic, 403, 434; syphilitic chorioido-, 401.
Retinoscopy, 80; with concave mirror, 80; with plane mirror, 86.
Retrobulbar neuritis, 432.
Rheumatism, 168, 245, 247, 256, 270, 274, 427, 432, 503, 545.
Robertson, Argyll, operation for ectropium, 225.
Rodent ulcer of the cornea, 190.
Rods and cones, layer of, 18, 402.
Röntgen rays, 390; used for diagnosis of retrobulbar growths, 506.

Saemisch's ulcer, 151.
Saline injection, subconjunctival, 47, 153.
Sarcoma of chorioid, 292; of ciliary body, 292; of conjunctiva, 129; of eyelid, 197; of iris, 291; of orbit, 508; of sclerotic, 249.
Scarlatina, 95, 134, 162, 429, 501, 555.
Scleritis, 168, 244.
Sclero-chorioiditis, 381.
Sclerosis, diffuse, of brain, 537; disseminated, 313, 428, 434, 462, 463, 499, 536.
Sclerotic, diseases of the, 244; injuries of the, 247; pigment spots of the, 249; ring, 88; rupture of the, 247; tumors of the, 249.
Sclerotomy, 336, 377; posterior, 333.
Scopolamin, 309, 316.
Scorbutus, 555.
Scotoma, 288, 410, 542; central, 401, 407, 429, 432, 435, 555; central color, 536; paracentric, 430; pericentric, 430; peripheral 401; positive, 261, 386, 414; ring, 401, 408; scintillating, 552.
Scotomata, symmetrical, 552.
Scott's operation for ectropion, 227.
Scrofula, 108.
Septicemia, 256, 407.
Seventh nerve, paralysis of, 471.
Shadow test, the, 80 (*see* Retinoscopy).
Short-sight (*see* Myopia).
Shot-silk retina, 89.
Sideroscope, 354, 392.
Sight, sense of, 25.
Sixth nerve, paralysis of the, 470.
Skin grafts, 513; Thiersch's, 127, 230; Wolfe and Lefort's, 230.
Smallpox, 95, 162, 256; conjunctival complication of, 119.
Smith, Priestley, method for measuring strabismus, 482; fusion tubes of, 487.
Snellen's operation for ectropion, 224; for entropion, 219; sunrise, 55, 58, 59; sutures, 224; test-types 59.
Snow-blindness, 415.
Spectacles, after extraction of cataract, 382; in accommodative asthenopia, 41; in albinismus, 299; in anisometropia, 64; in aphakia, 382; in astigmatism, 57; in conical cornea, 174; in convergent strabismus, 41, 485; in cramp of accommodation, 39; in hypermetropia, 41; in incipient cataract, 347; in irideremia, 298; in myopia, 48; in nebulous cornea, 183; in paralysis of accommodation, 69; in presbyopia, 66; numbering of lenses in, 17.
Sphenoid antrum, exostosis in the, 511; hyperostosis in the, 511; osteoma in the, 511; polypi in the, 511; sarcomata in the, 511; tumors in the, 511; fracture of the body of the, 529.
Sphenoidal fissure, lesion at the, 447; periostitis at the, 68.
Spinal amaurosis, 434.
Spinal cord, optic atrophy in disease of the, 434: pupil in diseases of the, 310; injuries of the, 546.
Spring catarrh, 105.
Squint, 472.
Squinting eye, amblyopia of the, 476.
Staphyloma, 120, 150, 151. 207, 246.
Staphyloma, anterior, 398; abscission, for, 170; evisceration for, 171; Mules' operation for, 172.
Staphyloma, corneal, 148, 182; of the eyeball, 246; partial, of the

cornea, 103, 169; posterior, 46; total of cornea, 103, 112, 170.
Stellwag's sign in exophthalmic goiter, 518.
Stenopeic apparatus, 174; spectacles, 183, 298, 299, 347.
Stenosis, lacrimal, 93.
Stereoscope, 479; for treatment of strabismus, 486.
Stomach, hemorrhages from the, 438.
Strabismus, 446; apparent, 446; apparent convergent, 45; apparent divergent, 39.
Strabismus, alternating, 477, 479; concomitant convergent, 41, 471, 472; dangers of operation for, 493; hypermetropia in, 41; measurement of, 480; by angular method, 483; by Hirschberg's method, 480; by linear method, 480; by method by tangents, 484; mobility of eye in, 484; monolateral, 476, 479; operative treatment of, 487; orthoptic treatment of, 485; periodic, 474, 479, 485; single vision in, 475; stereoscope for treatment of, 486; treatment by spectacles, 485.
Strabismus, concomitant divergent, 494; absolute, 498; dangers of operation for, 498; decentration of glasses in, 498.
Strabismus, paralytic, 472; tangents for estimating paralysis of orbital muscles, 484.
Strabometer, 480.
Strabotomy, 477.
Streatfield's operation for entropion, 218.
Streptothrix in the canalicus, 234.
Strumous ophthalmia, 133.
Stye, 193.
Superior oblique, paralysis of the, 451.
Sylvius, aqueduct of, 305, 307.
Symblepharon, 119, 132, 208, 229.
Sympathectomy, 337.
Sympathetic irritation, 265, 554.
Sympathetic ophthalmitis, 248, 252, 260, 266, 288, 289, 372, 393.
Synchysis, 386; scintillans, 387.
Synechia, anterior, 103, 147; circular, 251; complete, 251, 339; posterior, 251, 255, 305, 339; ring, 251, 339.
Syphilis, 68, 108, 168, 235, 270, 274, 427, 457, 470, 539, 545; congenital, 164, 247, 256, 264, 413; of conjunctiva, 128; primary, 189; secondary, 190, 403, 502; tertiary, 190.
Syphilitic chorioido-retinitis, 401; iritis, 253, 254, 259; sores of the eyelids, 189.
Syphiloma, 425.
Syringomyelia, 546.

Tabes dorsalis, 311, 312, 428, 434, 543.
Tenia solium, 417.
Tarsal tumor, 194.
Tarsorrhaphy, 207, 224, 521.
Tarsus, 110.
Tattooing for leukoma of cornea, 177, 184.
Teale's operation for symblepharon, 209.
Teeth, disease of the, 501, 553.
Teleangiectic tumors of the eyelids, 195, 509.
Temporo-sphenoidal lobe, tumor of the, 533.
Tenotomy for strabismus, 459, 481, 488, 489.
Tension of the eyeball, high or glaucomatous, 52, 182, 321; normal, 321.
Test-types, 29.
Thermo-cautère, 145, 176.
Third nerve, paralysis of, 68, 454, 459, 466.
Thomson's disease, 546.
Thrombosis, cerebral, 527; of retinal vessels, 412.
Tonometers, 321.
Toxic amblyopia, 28, 429; paralysis, 68.
Trachoma (or granular ophthalmia), 107, 121, 122, 124, 356; acute, 108; chronic, 110; forceps for treatment of, 115.
Trial lenses, numbering of, 17.
Trichiasis, 111, 192, 211.
Tropa-cocain, 319.
Trousseau's cerebral macula, 519.
Tubercle of the brain, 425, 529, 533; of the chorioid, 294; of the conjunctiva, 121; of the iris, 289.
Tubercular meningitis, 425, 632.
Tuberculosis, attenuated, of the iris, 290.
Tumor, cerebral, 425; intracranial, 312, 511; intra-ocular, 312, 340,

352, 398; in the antrum of Highmore, 511; in the antrum of the sphenoid, 511; in the body of the sphenoid bone, 511; in the ethmoid cells, 510; in the frontal sinus, 510; in the maxillary antrum, 511; of the lacrimal gland, 512; of the lenticular nucleus, 533; of the optic nerve, 436; of the optic thalamus, 523; of the pituitary body, 511, 529; of the temporo-sphenoidal lobe, 533.
Typhoid fever, 158, 256, 429, 501.

Ulcer of the cornea, absorption, 155; deep, 150; faceted, 155; rodent, 153; in purulent conjunctivitis, 98; ring, 135, 154; serpiginous, 151; simple, 149; superficial transparent, 156.
Uremia, 404.
Uric acid deposits in the conjunctiva, 131.
Uterine hemorrhage, 437.
Uveal tract, diseases of the, 250.
Uveitis, plastic, 354, 372, 393; purulent, 393.

Vaccine, vesicles, 190.
Valve of Vieussens, lesion of, 470.
Van Millingen's operation for trichiasis, 217.
Vertigo, 550.
Vienna operation for enucleation, 278.
Vision, acuteness of, 28, 32, 52; binocular field of, 32; central, 30, 401; center of, lesion at the, 522; color, 434; contraction of field of, 125; direct, 30; eccentric, 30; field of, 30, 324, 356; indirect, 30; monocular, double, 381; nulle, 530, 531.
Visual angle, 29; aphasia, 534; axis, 484; center, situation of, 524; hallucinations, 535, 550; line, 22, 24; memory, 535, 536.
Vitreous humor, 17, 51; artificial, 172; blood-vessels in the, 397; cholesterin in the, 337; cysticercus in the, 396; detachment of the from the retina, 397; diseases of the, 384; fluidity of the, 386; foreign bodies in the, 386; hemorrhages into the, 386, 398, 405, 406; inflammatory affections of the, 384; opacities in the, 47, 120, 246, 251, 260, 384, 385, 417, 418; persistent hyaloid artery in the, 397; prolapse of the, 362, 368; purulent inflammation of the, 384.
Vossius' operation for trichiasis, 216.

de Wecker's operation for staphyloma of the cornea, 170.
Wernicke's pupil-symptom, 532.
Wharton-Jones' operation for cicatricial ectropion, 228.
Whooping-cough, 127, 158, 264, 478.
Wolff's operation for congenital ptosis, 205.
Word-blindness, 531, 533.

Xanthelasma of the eyelids, 195.
Xerophthalmos, or xerosis of the conjunctiva, 119, 123, 208, 512, 554.

Yellow oxid of mercury ointment, 137.
Young-Helmholtz theory of the color-sense, 25.

Zonular cataract, 350.

MEDICAL BOOKS

Note Subject Index, Page 6



There have been sold more than
145,000 copies of Gould's Dictionaries
See Pages 12 and 13

P. BLAKISTON'S SON & COMPANY
PUBLISHERS OF MEDICAL AND SCIENTIFIC BOOKS
1012 WALNUT STREET, PHILADELPHIA

Montgomery's Gynecology

New Edition—Just Ready

A modern comprehensive Text-Book. By EDWARD E. MONTGOMERY, M.D., Professor of Gynecology in Jefferson Medical College, Philadelphia; Gynecologist to the Jefferson and St. Joseph's Hospitals, etc. Second Edition, Revised and Enlarged. 570 Illustrations, many of which are from original sources. Octavo. Cloth, $5.00; Leather, $6.00

*** This is a systematic modern treatise on Diseases of Women. The author's aim has been to produce a book that will be thorough and practical in every particular. The illustrations, nearly all of which are from original sources, have for the most part been drawn by special artists who, for a number of months, devoted their sole attention to this work. The present edition has been thoroughly revised.

"The book is one that can be recommended to the student, to the general practitioner—who must sometimes be a gynecologist to a certain extent whether he will or not—and to the specialist, as an ideal and in every way complete work on the gynecology of to-day—a practical work for practical workers."—*The Journal of the American Medical Association.*

Byford's Gynecology

Third Revised Edition

A MANUAL FOR STUDENTS AND PHYSICIANS

By HENRY T. BYFORD, M.D., Professor of Gynecology and Clinical Gynecology in the College of Physicians and Surgeons of Chicago; Professor of Clinical Gynecology, Women's Medical School of Northwestern University, and in Post-Graduate Medical School, etc. Third Edition, Enlarged. 363 Illustrations, many of which are from original drawings and several of which are Colored. 12mo.
Cloth, $3.00

"As a book to help the student to quickly review what ought to be gotten up, so as to be prepared for the early examination, it is of great service. Such a book would also make a most excellent text-book for the college class room."—*Virginia Medical Semi-Monthly*, Richmond.

By JAMES TYSON, M. D.,

Professor of Medicine, University of Pennsylvania, Physician to the Philadelphia Hospital, etc.

The Practice of Medicine. Third Edition.

A Text-Book for Physicians and Students, with Special Reference to Diagnosis and Treatment. With Colored Plates and many other Illustrations. Third Edition, Revised and Enlarged. 124 Illustrations. 8vo.

Cloth, $5.50; Leather, $6.50; Half Russia, $7.50

*** This edition has been entirely reset from new type. The author has revised it carefully and thoroughly, and added much new material and new illustrations.

"We are firmly convinced that at the present time Dr. Tyson's book on Practice can be most heartily commended to both the practitioner and student as a safe, reliable, and thoroughly up-to-date guide in the practice of medicine."—*The Therapeutic Gazette.*

"The clinical descriptions are clear and full, and the methods of treatment described are those generally recognized as being the most modern and satisfactory."—*The London Lancet.*

Guide to the Examination of Urine. Tenth Edition.

For the Use of Physicians and Students. With Colored Plate and Numerous Illustrations Engraved on Wood. Tenth Edition, Revised, Enlarged, and in many parts entirely rewritten. Cloth, $1.50

*** *A French translation of this book has been published in Paris.*

"The book is probably more widely and generally known and appreciated than any of its similars in subject and scope."—*New York Medical Journal.*

"The book is a reliable one, and should find a place in the library of every practitioner and student of medicine."—*Boston Medical and Surgical Journal.*

Handbook of Physical Diagnosis. Fourth Edition.

Revised and Enlarged. With two Colored Plates and 55 other Illustrations. 298 pages. 12mo. Cloth, $1.50

"Like everything else emanating from this distinguished author this little book is replete with practical information from beginning to end."—*The Chicago Medical Recorder.*

"The author approaches his subject from a practical point of view and the little work will prove a good friend to the student."—*The American Journal of the Medical Sciences.*

NEW THIRD EDITION—NOW READY

MORRIS' ANATOMY

Rewritten—Revised—Improved

WITH MANY NEW ILLUSTRATIONS

Out of 102 of the leading medical schools 60 recommend "Morris." It contains many features of special advantage to students. It is modern, up-to-date in every respect. It has been carefully revised, the articles on Osteology and Nervous System having been rewritten. **Each copy contains the colored illustrations and a Thumb Index.**

Octavo. With 846 Illustrations, of which 267 are printed in colors.

CLOTH, $6.00; LEATHER, $7.00

"The ever-growing popularity of the book with teachers and students is an index of its value, and it may safely be recommended to all interested."—From *The Medical Record*, New York.

"Of all the text-books of moderate size on human anatomy in the English language, Morris is undoubtedly the most up-to-date and accurate."—From *The Philadelphia Medical Journal.*

McMurrich—Embryology

THE DEVELOPMENT OF THE HUMAN BODY

With 270 Illustrations

A Text-Book for Medical Students. By J. PLAYFAIR MCMURRICH, Professor of Anatomy, Medical Department, University of Michigan. 527 pages. Cloth, $3.00

NINTH EDITION

POTTER'S MATERIA MEDICA, PHARMACY, AND THERAPEUTICS

An Exhaustive Handbook

Including the Action of Medicines, Special Therapeutics of Disease, Official and Practical Pharmacy, and Minute Directions for Prescription Writing, etc. Including over 650 Prescriptions and Formulæ. By SAMUEL O. L. POTTER, M.A., M.D., M.R.C.P. (Lond.), formerly Professor of the Principles and Practice of Medicine, Cooper Medical College, San Francisco; Major and Brigade Surgeon, U. S. Vol. Ninth Edition, Revised and Enlarged. 8vo.

With Thumb Index in each copy.

Cloth, $5.00; Leather, $6.00

*** This is the most complete and trustworthy book for the use of students and physicians. Students who purchase it will find it to contain a vast deal of information not in the usual text-books arranged in the most practical manner for facilitating study and reference. It cannot be surpassed as a physician's working book.

WHITE AND WILCOX. Materia Medica, Pharmacy, Pharmacology, and Therapeutics. Fifth Edition.

A Handbook for Students. By W. HALE WHITE, M.D., F.R.C.P., etc., Physician to, and Lecturer on Materia Medica and Therapeutics at, Guy's Hospital, etc. Fifth American Edition, Revised by REYNOLD W. WILCOX, M A, M.D., LL.D., Professor of Clinical Medicine and Therapeutics at the New York Post-Graduate Medical School and Hospital; Visiting Physician, St. Mark's Hospital; Assistant Visiting Physician, Bellevue Hospital. 12mo.

Cloth, $3.00; Leather, $3.50

SUBJECT INDEX.

Gould's Medical Dictionaries, - Pages 12, 13
Morris' Anatomy, New Edition, - - Page 4
Compends for Students, - - - - Page 27

SUBJECT.	PAGE
Alimentary Canal (see Surgery)	24
Anatomy	7
Anesthetics	18, 19
Autopsies (see Pathology)	20
Bacteriology	8
Bandaging (see Surgery)	24
Blood, Examination of	8
Brain	8
Chemistry. Physics	9
Children, Diseases of	11
Climatology	19
Clinical Charts	25
Compends	27
Consumption (see Lungs)	16
Cyclopedia of Medicine	13
Dentistry	11
Diabetes (see Urin. Organs)	25
Diagnosis	11
Diagrams (see Anatomy)	8
Dictionaries, Cyclopedias	12
Diet and Food	13
Disinfection	16
Dissectors	7
Ear	14
Electricity	14
Embryology	7
Emergencies	24
Eye	14
Fevers	15
Food	13
Formularies	21
Gynecology	15
Hay Fever	25
Heart	15
Histology	15
Hydrotherapy	19
Hygiene	16
Hypnotism	8
Insanity	8
Intestines	23
Latin, Medical (see Pharmacy)	21
Life Insurance	19
Lungs	16
Massage	17
Materia Medica	17
Mechanotherapy	17
Medical Jurisprudence	18
Mental Therapeutics	8
Microscopy	18
Milk	8, 10
Miscellaneous	18
Nervous Diseases	19
Nose	25
Nursing	20
Obstetrics	20
Ophthalmology	14
Organotherapy	18
Osteology (see Anatomy)	7
Pathology	20
Pharmacy	21
Physical Diagnosis	11
Physical Training	17
Physiology	22
Pneumotherapy	19
Poisons (see Toxicology)	18
Practice of Medicine	22
Prescription Books (Pharmacy)	21
Refraction (see Eye)	14
Rest	19
Sanitary Science	16
Serum-Therapy	17
Skin	23
Spectacles (see Eye)	14
Spine (see Nervous Diseases)	19
Stomach	23
Students' Compends	27
Surgery and Surgical Diseases	24
Technological Books	9
Temperature Charts	25
Therapeutics	17
Throat	25
Toxicology	18
Tumors (see Surgery)	24
U. S. Pharmacopœia	22
Urinary Organs	25
Urine	25
Venereal Diseases	26
Veterinary Medicine	26
Visiting Lists, Physicians'. (*Send for Special Circular.*)	
Water Analysis	16
Women, Diseases of	15

Self-Examination for Medical Students. 3500 Questions on Medical Subjects, with References to Standard Works in which the correct replies will be found. Together with Questions from State Examining Boards. 3d Ed. Paper Cover, 10 cts.

SPECIAL NOTE.—The prices given in this catalogue are net; no discount can be allowed retail purchasers under any consideration. This rule has been established in order that everyone will be treated alike, a general reduction in former prices having been made to meet previous retail discounts. Upon receipt of the advertised price any book will be forwarded by mail or express, all charges prepaid.

ANATOMY. EMBRYOLOGY.

MORRIS. Text-Book of Anatomy. Third Revised and Enlarged Edition. 846 Illustrations, 267 of which are printed in colors. *Thumb Index in Each Copy.* Cloth, $6.00; Leather, $7.00
"The ever-growing popularity of the book with teachers and students is an index of its value."—*Medical Record, New York.*

BROOMELL. Anatomy and Histology of the Human Mouth and Teeth. 2d Edition, Enlarged. 337 Illus. Cloth, $4.50

DAVISSON. Mammalian Anatomy. With Special Reference to the Cat. 110 Illustrations. *In Press.*

DEAVER. Surgical Anatomy. A Treatise on Anatomy in its Application to Medicine and Surgery. With 499 very handsome full-page Illustrations Engraved from Original Drawings made by special Artists from dissections prepared for the purpose. Three vols. By *Subscription only.*
Half Morocco or Sheep, $24.00; Half Russia, $27.00

GORDINIER. Anatomy of the Central Nervous System. With 271 Illustrations, many of which are original. Cloth, $6.00

HEATH. Practical Anatomy. 9th Edition. 321 Illus. $4.25

HOLDEN. Anatomy. A Manual of Dissections. Revised by A. HEWSON, M.D., Demonstrator of Anatomy, Jefferson Medical College, Philadelphia. 320 handsome Illustrations. 7th Ed. In two compact 12mo volumes. 850 pages. Large New Type.
Vol. I. Scalp—Face—Orbit—Neck—Throat—Thorax—Upper Extremity. $1.50
Vol. II. Abdomen—Perineum—Lower Extremity—Brain—Eye—Ear—Mammary Gland—Scrotum—Testes. $1.50

HOLDEN. Human Osteology. Comprising a Description of the Bones, with Colored Delineations of the Attachments of the Muscles. The General and Microscopical Structure of Bone and its Development. With Lithographic Plates and numerous Illustrations. 8th Edition. $5.25

HOLDEN. Landmarks, Medical and Surgical. 4th Ed. .75

HUGHES AND KEITH. Dissections. With 527 Colored Plates and other Illustrations. In three parts.
I, Upper and Lower Extremity. $3.00
II, Abdomen—Thorax. $3.00
III, Head—Neck—Central Nervous System. $3.00

LAZARUS-BARLOW. Pathological Anatomy. 21 Plates and 171 other Illustrations. *Just Ready.* $6.50

McMURRICH. Embryology. The Development of the Human Body. 276 Illustrations. $3.00

MARSHALL. Physiological Diagrams. Eleven Life-Size Colored Diagrams (each seven feet by three feet seven inches). Designed for Demonstration before the Class.
In Sheets, Unmounted, $40.00; Backed with Muslin and Mounted on Rollers, $60.00; Ditto, Spring Rollers, in handsome Walnut Wall Map Case, $100.00; Single Plates—Sheets, $5.00; Mounted, $7.50. Explanatory Key, .50. *Purchaser must pay freight charges.*

MINOT. Laboratory Text-Book of Embryology. 218 Illustrations. *Just Ready.* $4.50

POTTER. Compend of Anatomy, Including Visceral Anatomy. 7th Edition, Revised and Enlarged. 16 Plates and 138 other Illustrations. *Just Ready.* .80; Interleaved, $1.00

WILSON. Anatomy. 11th Edition. 429 Illus., 26 Plates. $5.00

YUTZY. Guide to the Dissection of the Human Body. Based on Morris' Anatomy. Paper Cover, .25

BACTERIOLOGY.

CONN. Agricultural Bacteriology. Including the Study of Bacteria as relating to Agriculture, Soil, Dairy and Food Products, Sewage, Domestic Animals, etc. Illustrated. $2.50

CONN. Bacteria in Milk and Its Products. Designed for Students of Dairying, Boards of Health, Bacteriologists, etc. Illustrated. $1.25

EMERY. Bacteriological Diagnosis. 2 Colored Plates and 32 other Illustrations. $1.50

HEWLETT. Manual of Bacteriology. 75 Illustrations. Second Edition, Revised and Enlarged. $4.00

SMITH. Laboratory Exercises in Bacteriology. A Handbook for Students. Illustrated. $1.50

WILLIAMS. Bacteriology. A Manual for Students. 88 Illustrations. 3d Edition, Revised. *Just Ready.* $1.50

BLOOD, Examination of.

DA COSTA. Clinical Hematology. A Practical Guide to the Examination of the Blood, with Reference to Diagnosis. Six Colored Plates and 48 other Illus. Cloth, $5.00; Sheep, $6.00

BRAIN AND INSANITY (see also Nervous Diseases.)

BLACKBURN. A Manual of Autopsies. Designed for the Use of Hospitals for the Insane and other Public Institutions. Ten full-page Plates and other Illustrations. $1.25

CHASE. General Paresis. Illustrated. $1.75

DERCUM. Mental Therapeutics, Rest, Suggestion. *See Cohen, Physiologic Therapeutics, page* 17.

GORDINIER. The Gross and Minute Anatomy of the Central Nervous System. With full-page and other Illus. $6.00

IRELAND. The Mental Affections of Children. 2d Ed. $4.00

LEWIS (BEVAN). Mental Diseases. A Text-Book having Special Reference to the Pathological Aspects of Insanity. 26 Lithographic Plates and other Illustrations. 2d Ed. $7.00

MANN. Manual of Psychological Medicine. $3.00

PERSHING. Diagnosis of Nervous and Mental Disease. Illustrated. $1.25

REGIS. Mental Medicine. Authorized Translation by H. M. BANNISTER, M.D. $2.00

STEARNS. Mental Diseases. With a Digest of Laws Relating to Care of Insane. Illustrated. Cloth, $2.75; Sheep, $3.25

TUKE. Dictionary of Psychological Medicine. Giving the Definition, Etymology, and Symptoms of the Terms used in Medical Psychology, with the Symptoms, Pathology, and Treatment of the Recognized Forms of Mental Disorders. Two volumes. $10.00

WOOD, H. C. Brain and Overwork. .40

CHEMISTRY AND TECHNOLOGY.

Special Catalogue of Chemical Books sent free upon application.

ALLEN. Commercial Organic Analysis. A Treatise on the Modes of Assaying the Various Organic Chemicals and Products Employed in the Arts, Manufactures, Medicine, etc., with Concise Methods for the Detection of Impurities, Adulterations, etc. 8vo.

Vol. I. Alcohols, Neutral Alcoholic Derivatives, etc., Ethers, Vegetable Acids, Starch, Sugars, etc. 3d Edition. $4.50

Vol. II, Part I. Fixed Oils and Fats, Glycerol, Explosives, etc. 3d Edition. $3.50

Vol. II, Part II. Hydrocarbons, Mineral Oils, Lubricants, Benzenes, Naphthalenes and Derivatives, Creosote, Phenols, etc. 3d Edition. $3.50

Vol. II, Part III. Terpenes, Essential Oils, Resins, Camphors, etc. 3d Edition. *Preparing.*

Vol. III, Part I. Tannins, Dyes, and Coloring Matters. 3d Edition, Enlarged and Rewritten. Illustrated. $4.50

Vol. III, Part II. The Amines, Hydrazines and Derivatives, Pyridine Bases. The Antipyretics, etc. Vegetable Alkaloids, Tea, Coffee, Cocoa, etc. 8vo. 2d Edition. $4.50

Vol. III, Part III. Vegetable Alkaloids, Non-Basic Vegetable Bitter Principles. Animal Bases, Animal Acids, Cyanogen Compounds, etc. 2d Edition, 8vo. $4.50

Vol. IV. The Proteids and Albuminous Principles. 2d Edition. $4.50

BAILEY AND CADY. Qualitative Chemical Analysis. $1.25

BARTLEY. Medical and Pharmaceutical Chemistry. A Text-Book for Medical, Dental, and Pharmaceutical Students. With Illustrations, Glossary, and Complete Index. 5th Ed. $3.00

BARTLEY. Clinical Chemistry. The Examination of Feces, Saliva, Gastric Juice, Milk, and Urine. $1.00

BLOXAM. Chemistry, Inorganic and Organic. With Experiments. 9th Ed., Revised. 281 Engravings. *Preparing.*

BUNGE. Physiologic and Pathologic Chemistry. From the Fourth German Enlarged Edition. $3.00

CALDWELL. Elements of Qualitative and Quantitative Chemical Analysis. 3d Edition, Revised. $1.00

CAMERON. Soap and Candles. 54 Illustrations. $2.00

CLOWES AND COLEMAN. Quantitative Analysis. 5th Edition. 122 Illustrations. $3.50

COBLENTZ. Volumetric Analysis. Illustrated. $1.25

CONGDON. Laboratory Instructions in Chemistry. With Numerous Tables and 56 Illustrations. $1.00

GARDNER. The Brewer, Distiller, and Wine Manufacturer. Illustrated. $1.50

GRAY. Physics. Volume I. Dynamics and Properties of Matter. 350 Illustrations. $4.50

GROVES AND THORP. Chemical Technology. The Application of Chemistry to the Arts and Manufactures.

Vol. I. Fuel and its Applications. 607 Illustrations and 4 Plates. Cloth, $5.00; ½ Mor., $6.50

Vol. II. Lighting. Illustrated. Cloth, $4.00; ½ Mor., $5.50

Vol. III. Gas Lighting. Cloth, $3.50; ½ Mor., $4.50

Vol. IV. Electric Lighting. Photometry. Cloth, $3.50; ½ Mor., $4.50

HEUSLER. The Chemistry of the Terpenes. $4.00

HOLLAND. The Urine, the Gastric Contents, the Common Poisons, and the Milk. Memoranda, Chemical and Microscopical, for Laboratory Use. 6th Ed. Illustrated. $1.00

LEFFMANN. Compend of Medical Chemistry, Inorganic and Organic. 4th Edition, Revised. .80; Interleaved, $1.00

LEFFMANN. Analysis of Milk and Milk Products. 2d Edition, Enlarged. Illustrated. $1.25

LEFFMANN. Water Analysis. For Sanitary and Technic Purposes. Illustrated. 4th Edition. $1.25

LEFFMANN. Structural Formulæ. Including 180 Structural and Stereo-Chemical Formulæ. 12mo. Interleaved. $1.00

LEFFMANN AND BEAM. Select Methods in Food Analysis. Illustrated. $2.50

MUTER. Practical and Analytical Chemistry. 3d American from the Ninth English Edition. Revised to meet the requirements of American Students. 56 Illustrations. $1.25

OETTEL. Exercises in Electro-Chemistry. Illustrated. .75

OETTEL. Electro-Chemical Experiments. Illustrated. .75

RICHTER. Inorganic Chemistry. 5th American from 10th German Edition. Authorized translation by EDGAR F. SMITH, M.A., PH.D. 89 Illustrations and a Colored Plate. $1.75

RICHTER. Organic Chemistry. 3d American Edition, translated from the 8th German by EDGAR F. SMITH. Illus. 2 vols.

Vol. I. Aliphatic Series. 625 pages. $3.00

Vol. II. Carbocyclic Series. 671 pages. $3.00

ROCKWOOD. Chemical Analysis for Students of Medicine, Dentistry, and Pharmacy. Illustrated. $1.50

SMITH. Electro-Chemical Analysis. 3d Ed. 39 Illus. $1.50

SMITH AND KELLER. Experiments. Arranged for Students in General Chemistry. 4th Edition. Illustrated. .60

SUTTON. Volumetric Analysis. A Systematic Handbook for the Quantitative Estimation of Chemical Substances by Measure, Applied to Liquids, Solids, and Gases. 8th Edition, Revised. 112 Illustrations. $5.00

SYMONDS. Manual of Chemistry. 2d Edition. $2.00

TRAUBE. Physico-Chemical Methods. 97 Illustrations. $1.50

THRESH. Water and Water Supplies. 3d Edition. $2.00

ULZER AND FRAENKEL. Chemical Technical Analysis. Translated by Fleck. Illustrated. $1.25

WOODY. Essentials of Chemistry and Urinalysis. 4th Edition. Illustrated. $1.50

*** *Special Catalogue of Books on Chemistry free upon application.*

CHILDREN.

HATFIELD. Compend of Diseases of Children. With a Colored Plate. 3d Ed. *Just Ready.* .80; Interleaved, $1.00

IRELAND. The Mental Affections of Children. Idiocy, Imbecility, Insanity, etc. 2d Edition. $4.00

POWER. Surgical Diseases of Children and their Treatment by Modern Methods. Illustrated. $2.50

STARR. The Digestive Organs in Childhood. The Diseases of the Digestive Organs in Infancy and Childhood. 3d Edition, Rewritten and Enlarged. Illustrated. $3.00

STARR. Hygiene of the Nursery. Including the General Regimen and Feeding of Infants and Children, and the Domestic Management of the Ordinary Emergencies of Early Life, Massage, etc. 6th Edition. 25 Illustrations. $1.00

SMITH. Wasting Diseases of Children. 6th Edition. $2.00

TAYLOR AND WELLS. The Diseases of Children. 2d Edition, Revised and Enlarged. Illustrated. 8vo. $4.50

"It is well worthy the careful study of both student and practitioner, and can not fail to prove of great value to both. We do not hesitate to recommend it."—*Boston Medical and Surgical Journal.*

DIAGNOSIS.

BROWN. Medical Diagnosis. A Manual of Clinical Methods. 4th Edition. 112 Illustrations. Cloth, $2.25

DA COSTA. Clinical Hematology. A Practical Guide to Examination of Blood, with Reference to Diagnosis. 6 Colored Plates, 48 other Illustrations. Cloth, $5.00; Sheep, $6.00

DOUGLAS. Surgical Diseases of Abdomen, with Reference to Diagnosis. 20 Full-Page Plates. *Just Ready.* Cloth, $7.00; Sheep, $8.00

EMERY. Bacteriological Diagnosis. 2 Colored Plates and 32 other Illustrations. $1.50

MEMMINGER. Diagnosis by the Urine. 2d Ed. 24 Illus. $1.00

PERSHING. Diagnosis of Nervous and Mental Diseases. Illustrated. $1.25

STEELL. Physical Signs of Pulmonary Disease. $1.25

TYSON. Handbook of Physical Diagnosis. For Students and Physicians. By the Professor of Clinical Medicine in the University of Pennsylvania. Illus. 4th Ed., Improved and Enlarged. With 2 Colored and 55 other Illustrations. $1.50

DENTISTRY.

Special Catalogue of Dental Books sent free upon application.

BARRETT. Dental Surgery for General Practitioners and Students of Medicine and Dentistry. Extraction of Teeth, etc. 3d Edition. Illustrated. $1.00

BROOMELL. **Anatomy and Histology of the Human Mouth and Teeth.** Second Edition, Revised and Enlarged. 337 handsome Illustrations. Cloth, $4.50; Leather, $5.50

FILLEBROWN. **Operative Dentistry.** Illustrated. $2.25

GORGAS. **Dental Medicine. A Manual of Materia Medica and Therapeutics.** 7th Edition. Cloth, $4.00; Sheep, $5.00

GORGAS. **Questions and Answers for the Dental Student.** Embracing all the subjects in the Curriculum of the Dental Student. Octavo. $6.00

HARRIS. **Principles and Practice of Dentistry.** Including Anatomy, Physiology, Pathology, Therapeutics, Dental Surgery, and Mechanism. 13th Edition. Revised by F. J. S. GORGAS, M.D., D.D.S. 1250 Illus. Cloth, $6.00; Leather, $7.00

HARRIS. **Dictionary of Dentistry.** Including Definitions of Such Words and Phrases of the Collateral Sciences as Pertain to the Art and Practice of Dentistry. 6th Edition, Revised and Enlarged by FERDINAND J. S. GORGAS, M.D., D.D.S. Cloth, $5.00; Leather, $6.00

RICHARDSON. **Mechanical Dentistry.** 7th Edition. Thoroughly Revised and Enlarged by DR. GEO. W. WARREN. 691 Illustrations. Cloth, $5.00; Leather, $6.00

SMITH. **Dental Metallurgy.** 2d Edition. Illustrated. $2.00

TAFT. **Index of Dental Periodical Literature.** $2.00

TOMES. **Dental Anatomy.** 263 Illustrations. 5th Ed. $4.00

TOMES. **Dental Surgery.** 4th Edition. 289 Illus. $4.00

WARREN. **Compend of Dental Pathology and Dental Medicine.** With a Chapter on Emergencies. 3d Edition. Illustrated. .80; Interleaved, $1.00

WARREN. **Dental Prosthesis and Metallurgy.** 129 Illus. $1.25

WHITE. **The Mouth and Teeth.** Illustrated. .40

DICTIONARIES. CYCLOPEDIAS.

GOULD. **The Illustrated Dictionary of Medicine, Biology, and Allied Sciences.** Being an Exhaustive Lexicon of Medicine and those Sciences Collateral to it: Biology (Zoology and Botany), Chemistry, Dentistry, Pharmacology, Microscopy, etc., with many useful Tables and numerous fine Illustrations. 1633 pages. Fifth Edition.
Sheep or Half Morocco. $10.00; with Thumb Index, $11.00
Half Russia, Thumb Index, $12.00

GOULD. **The Medical Student's Dictionary.** 11th Edition. Illustrated. Including those Words and Phrases generally used in Medicine, with their Proper Pronunciation and Definition, Based on Recent Medical Literature. With Table of Eponymic Terms and Tests and Tables of the Bacilli, Micrococci, Mineral Springs, etc., of the Arteries, Muscles, Nerves, Ganglia, Plexuses, etc. Eleventh Edition. Enlarged and illustrated with a large number of Engravings. 840 pages.
Half Morocco, $2.50; with Thumb Index, $3.00
Flexible Leather, Burnished Edges, Thumb Index, 3.50

GOULD. The Pocket Pronouncing Medical Lexicon. 4th Edition. (30,000 Medical Words Pronounced and Defined.) Containing all the Words, their Definition and Pronunciation, that the Medical, Dental, or Pharmaceutical Student Generally Comes in Contact with; also Elaborate Tables of Eponymic Terms, Arteries, Muscles, Nerves, Bacilli, etc., etc., a Dose List in both English and Metric Systems, etc., Arranged in a Most Convenient Form for Reference and Memorizing. Fourth Edition, Revised and Enlarged. 838 pages.
Full Limp Leather, Gilt Edges, $1.00; Thumb Index, $1.25
145,000 Copies of Gould's Dictionaries have been sold.

GOULD AND PYLE. Cyclopedia of Practical Medicine and Surgery. Seventy-two Special Contributors. Illustrated. One Volume. A Concise Reference Handbook of Medicine, Surgery, Obstetrics, Materia Medica, Therapeutics, and the Various Specialties, with Particular Reference to Diagnosis and Treatment. Compiled under the Editorial Supervision of GEORGE M. GOULD, M.D., Author of "An Illustrated Dictionary of Medicine," etc.; and WALTER L. PYLE, M.D., Assistant Surgeon Wills Eye Hospital; formerly Editor "International Medical Magazine," etc., and Seventy-two Special Contributors. With many Illustrations. Large Square 8vo, to correspond with Gould's "Illustrated Dictionary."
Full Sheep or Half Mor., $10.00; with Thumb Index, $11.00
Half Russia, Thumb Index, $12.00 net.

GOULD AND PYLE. Pocket Cyclopedia of Medicine and Surgery. Based upon above book and uniform in size with "Gould's Pocket Dictionary."
Full Limp Leather, Gilt Edges, $1.00
With Thumb Index, $1.25

HARRIS. Dictionary of Dentistry. Including Definitions of Such Words and Phrases of the Collateral Sciences as Pertain to the Art and Practice of Dentistry. 6th Edition, Revised and Enlarged by FERDINAND J. S. GORGAS, M.D., D.D.S.
Cloth, $5.00; Leather, $6.00

LONGLEY. Pocket Medical Dictionary. Cloth, .75

TREVES AND LANG. German-English Medical Dictionary.
Half Calf, $3.25

DIET AND FOOD.

ALLEN. Proteids and Albuminous Principles. An analytical Study of Food Products. 2d Edition. $4.50

BURNETT. Foods and Dietaries. A Manual of Clinical Dietetics, with Diet Lists for Various Diseases, etc. 2d Ed. $1.50

DAVIS. Dietotherapy. Food in Health and Disease. With Tables of Dietaries, Relative Value of Foods, etc. *See Cohen, Physiologic Therapeutics, page* 17.

GREENISH. Microscopical Examination of Foods and Drugs. Illustrated. *Just Ready.* $3.50

HAIG. Diet and Food. Considered in Relation to Strength and Power of Endurance. 4th Edition. $1.00

LEFFMANN. Select Methods in Food Analysis. Illus. $2.50

EAR (see also Throat and Nose).

BURNETT. Hearing and How to Keep It. Illustrated. .40

HOVELL. Diseases of the Ear and Naso-Pharynx. Including Anatomy and Physiology of the Organ, together with the Treatment of the Affections of the Nose and Pharynx which Conduce to Aural Disease. 128 Illustrations. 2d Ed. $5.50

PRITCHARD. Diseases of the Ear. 4th Edition, Enlarged. Many Illustrations and Formulæ. *In Press.*

ELECTRICITY.

BIGELOW. Plain Talks on Medical Electricity and Batteries. With a Therapeutic Index and a Glossary. 43 Illustrations. 2d Edition. $1.00

HEDLEY. Therapeutic Electricity and Practical Muscle Testing. 99 Illustrations. $2.50

JACOBY. Electrotherapy. 2 volumes. Illustrated. *See Cohen, Physiologic Therapeutics, page* 17.

JONES. Medical Electricity. 3d Edition. 117 Illus. $3.00

EYE.

A Special Circular of Books on the Eye sent free upon application.

DARIER. Ocular Therapeutics. *Just Ready.* $3.00

DONDERS. The Nature and Consequences of Anomalies of Refraction. With Portrait and Illus. Half Morocco, $1.25

FICK. Diseases of the Eye and Ophthalmoscopy. Translated by A. B. HALE, M.D. 157 Illus. Cloth, $4.50; Sheep, $5.50

GOULD AND PYLE. Compend of Diseases of the Eye and Refraction. Including Treatment and Operations, and a Section on Local Therapeutics. With Formulæ, Useful Tables, a Glossary, and 111 Illus., several of which are in colors. 2d Edition, Revised. Cloth, .80; Interleaved, $1.00

GREEFF. The Microscopic Examination of the Eye. Illustrated. $1.25

HARLAN. Eyesight, and How to Care for It. Illus. .40

HARTRIDGE. On the Ophthalmoscope. 4th Edition. With 4 Colored Plates and 68 Wood-cuts. $1.50

HARTRIDGE. Refraction. 104 Illustrations and Test Types. 12th Edition, Enlarged. $1.50

HANSELL AND SWEET. Treatise on Diseases of the Eye. With many Illus. drawn by special artists, etc. *In Press.*

HANSELL AND REBER. Muscular Anomalies of the Eye. Illustrated. $1.50

HANSELL AND BELL. Clinical Ophthalmology. Colored Plate of Normal Fundus and 120 Illustrations. $1.50

HENDERSON. Notes on the Eye. 3d Ed. *Just Ready.* $1.50

JENNINGS. Manual of Ophthalmoscopy. 95 Illustrations and 1 Colored Plate. $1.50

MORTON. Refraction of the Eye. Its Diagnosis and the Correction of its Errors. 6th Edition. $1.00

OHLEMANN. Ocular Therapeutics. Authorized Translation, and Edited by DR. CHARLES A. OLIVER. $1.75

PARSONS. Elementary Ophthalmic Optics. With Diagrammatic Illustrations. $2.00

PHILLIPS. Spectacles and Eyeglasses. Their Prescription and Adjustment. 3d Edition. 52 Illustrations. $1.00

SWANZY. Diseases of the Eye and Their Treatment. 8th Edition, Revised and Enlarged. 167 Illustrations, 1 Plain Plate, and a Zephyr Test Card. *Just Ready.* $2.50

From The Medical News.

"Swanzy has succeeded in producing the most intellectually conceived and thoroughly executed résumé of the science within the limits he has assigned himself. As a 'student's handbook,' small in size and of moderate price, it can hardly be equaled."

THORINGTON. Retinoscopy. 4th Edition, Carefully Revised. Illustrated. $1.00

THORINGTON. Refraction and How to Refract. 200 Illustrations, 13 of which are colored. 2d Edition. $1.50

WALKER. Student's Aid in Ophthalmology. Colored Plate and 40 other Illustrations and a Glossary. $1.50

WORTH. Squint: Its Causes, Pathology, Treatment. $2.00

WRIGHT. Ophthalmology. 2d Edition, Revised and Enlarged. 117 Illustrations and a Glossary. $3.00

FEVERS.

GOODALL AND WASHBOURN. Fevers and Their Treatment. Illustrated. $3.00

GYNECOLOGY.

BISHOP. Uterine Fibromyomata. Their Pathology, Diagnosis, and Treatment. Illustrated. Cloth, $3.50

BYFORD (H. T.). Manual of Gynecology. 3d Edition, Revised and Enlarged. 363 Illustrations. $3.00; Sheep, $3.50

DÜHRSSEN. A Manual of Gynecological Practice. 105 Illustrations. $1.50

FULLERTON. Surgical Nursing. 3d Edition, Revised and Enlarged. 69 Illustrations. $1.00

LEWERS. Diseases of Women. 146 Illus. 5th Ed. $2.50

LEWERS. Cancer of the Uterus. *Just Ready.* $3.00

MONTGOMERY. Practical Gynecology. A Complete Systematic Text-Book. 2d Edition, Revised and Enlarged. With 570 Illus. *Just Ready.* Cloth, $5.00; Leather, $6.00

ROBERTS. Gynecological Pathology. With 127 Full-page Plates containing 151 Figures. $6.00

WELLS. Compend of Gynecology. Illustrated. 3d Edition, Revised and Enlarged. *Just Ready.* .80; Interleaved, $1.00

HEART.

THORNE. The Schott Methods of the Treatment of Chronic Heart Disease. Fourth Edition. Illustrated. $2.00

HISTOLOGY.

CUSHING. Compend of Histology. By H. H. Cushing, M.D., Demonstrator of Histology, Jefferson Medical College, Philadelphia. Illus. *Nearly Ready.* .80; Interleaved, $1.00

LAZARUS-BARLOW. Pathological Anatomy and Histology. Illustrated. *Just Ready.* $6.50

STIRLING. Outlines of Practical Histology. 368 Illustrations. 2d Edition, Revised and Enlarged. With new Illus. $2.00

STÖHR. Histology and Microscopical Anatomy. Edited by A. SCHAPER, M.D., University of Breslau, formerly Demonstrator of Histology, Harvard Medical School. Fifth American from 10th German Edition, Revised and Enlarged. 353 Illustrations. $3.00

HYGIENE.

Special Catalogue of Books on Hygiene sent free upon application.

CANFIELD. Hygiene of the Sick-Room. A Book for Nurses and Others. Being a Brief Consideration of Asepsis, Antisepsis, Disinfection, Bacteriology, Immunity, Heating, Ventilation, etc. $1.25

CONN. Agricultural Bacteriology. Illustrated. $2.50

CONN. Bacteriology of Milk and Milk Products. Illus. $1.25

COPLIN. Practical Hygiene. A Complete American Text-Book. 138 Illustrations. New Edition. *Preparing.*

HARTSHORNE. Our Homes. Illustrated. .40

KENWOOD. Public Health Laboratory Work. 116 Illustrations and 3 Plates. $2.00

LEFFMANN. Select Methods in Food Analysis. 53 Illustrations and 4 Plates. $2.50

LEFFMANN. Examination of Water for Sanitary and Technical Purposes. 4th Edition. Illustrated. $1.25

LEFFMANN. Analysis of Milk and Milk Products. Illustrated. Second Edition. $1.25

LINCOLN. School and Industrial Hygiene. .40

McFARLAND. Prophylaxis and Personal Hygiene. Care of the Sick. *See Cohen, Physiologic Therapeutics, page* 17.

NOTTER. The Theory and Practice of Hygiene. 15 Plates and 138 other Illustrations. 8vo. 2d Edition. $7.00

PARKES AND KENWOOD. Hygiene and Public Health. 2d Edition, Enlarged. Illustrated. $3.00

ROSENAU. Disinfection and Disinfectants. Illus. $2.00

STARR. The Hygiene of the Nursery. Including the General Regimen and Feeding of Infants and Children, and the Domestic Management of the Ordinary Emergencies of Early Life, Massage, etc. 6th Edition. 25 Illustrations. $1.00

STEVENSON AND MURPHY. A Treatise on Hygiene. By Various Authors. In three octavo volumes. Illustrated. Vol. I, $6.00; Vol. II, $6.00; Vol. III, $5.00

THRESH. Water and Water Supplies. 3d Edition. $2.00

WILSON. Handbook of Hygiene and Sanitary Science. With Illustrations. 8th Edition. $3.00

WEYL. Sanitary Relations of the Coal-Tar Colors. Authorized Translation by HENRY LEFFMANN, M.D., PH.D. $1.25

LUNGS AND PLEURÆ.

KNOPF. Pulmonary Tuberculosis. Its Modern Prophylaxis and Treatment in Special Institutions and at Home. Illus. $3.00

STEELL. Physical Signs of Pulmonary Disease. Illus. $1.25

MASSAGE. PHYSICAL EXERCISE.

OSTROM. Massage and the Original Swedish Movements. Their Application to Various Diseases of the Body. A Manual for Students, Nurses, and Physicians. Fifth Edition, Enlarged. 115 Illustrations, many of which are original. $1.00

MITCHELL AND GULICK. Mechanotherapy. Exercise, Orthopedics, Massage, Ocular Corrections, etc. Illustrated. *See Cohen, Physiologic Therapeutics, below.*

TREVES. Physical Education. Its Value, Methods, etc. .75

MATERIA MEDICA AND THERAPEUTICS.

BRACKEN. Outlines of Materia Medica and Pharmacology. $2.75

COBLENTZ. The Newer Remedies. Including their Synonyms, Sources, Methods of Preparation, Tests, Solubilities, Doses, etc. 3d Edition, Enlarged and Revised. $1.00

COHEN. Physiologic Therapeutics. Methods other than Drug-Giving useful in the Prevention of Disease and in the Treatment of the Sick. Mechanotherapy, Mental Therapeutics, Suggestion, Electrotherapy, Climatology, Hydrotherapy, Pneumatotherapy, Prophylaxis, Dietetics, Organotherapy, Phototherapy, Mineral Waters, Baths, etc. 11 volumes, 8vo. Illustrated. (*Subscription.*) Cloth, $27.50; ½ Mor., $38.50

Special Descriptive Circular will be sent upon application.

GORGAS. Dental Medicine. A Manual of Materia Medica and Therapeutics. 7th Edition, Revised. $4.00

GROFF. Materia Medica for Nurses, with Questions for Self-Examination. 2d Edition, Revised and Improved. $1.25

HELLER. Essentials of Materia Medica, Pharmacy, and Prescription Writing. $1.50

HEWLETT. Serum-Therapy. *In Press.*

MAYS. Theine in the Treatment of Neuralgia. ½ bound. .50

POTTER. Handbook of Materia Medica, Pharmacy, and Therapeutics, including the Action of Medicines, Special Therapeutics, Pharmacology, etc., including over 600 Prescriptions and Formulæ. 9th Edition, Revised and Enlarged. With Thumb Index in each copy. *Just Ready.* Cloth, $5.00; Sheep, $6.00

"In conclusion we may add that Dr. Potter's Therapeutics covers a wider field than many books which bear this title. He discusses a good many drugs which are rarely employed, and therefore the book is as useful to one who wishes to look for unusual information as it is to him who wishes a handbook for ready reference in the treatment of disease as he meets it from day to day."—*Therapeutic Gazette.*

POTTER. Compend of Materia Medica, Therapeutics, and Prescription Writing, with Special Reference to the Physiological Action of Drugs. 6th Edition. .80; Interleaved, $1.00

MURRAY. Rough Notes on Remedies. 4th Edition. $1.25

SAYRE. Organic Materia Medica and Pharmacognosy. An Introduction to the Study of the Vegetable Kingdom and the Vegetable and Animal Drugs. Comprising the Botanical and Physical Characteristics, Source, Constituents, and Pharmacopeial Preparations, Insects Injurious to Drugs, and Pharmacal Botany. With sections on Histology and Microtechnique, by W. C. Stevens. 374 Illustrations, many of which are original. 2d Edition. Cloth, $4.50

TAVERA. Medicinal Plants of the Philippines. $2.00

WHITE AND WILCOX. Materia Medica, Pharmacy, Pharmacology, and Therapeutics. 5th American Edition, Revised by REYNOLD W. WILCOX, M.A., M.D., LL.D., Professor of Clinical Medicine and Therapeutics at the New York Post-Graduate Medical School. Cloth, $3.00; Leather, $3.50

"The care with which Dr. Wilcox has performed his work is conspicuous on every page, and it is evident that no recent drug possessing any merit has escaped his eye. We believe, on the whole, this is the best book on Materia Medica and Therapeutics to place in the hands of students, and the practitioner will find it a most satisfactory work for daily use."—*The Cleveland Medical Gazette.*

MEDICAL JURISPRUDENCE AND TOXICOLOGY.

REESE. Medical Jurisprudence and Toxicology. A Text-Book for Medical and Legal Practitioners and Students. 6th Edition. Revised by HENRY LEFFMANN. M.D. Cloth, $3.00; Leather, $3.50

"To the student of medical jurisprudence and toxicology it is invaluable, as it is concise, clear, and thorough in every respect." —*The American Journal of the Medical Sciences.*

MANN. Forensic Medicine and Toxicology. Illus. $6.50

TANNER. Memoranda of Poisons. Their Antidotes and Tests. 9th Edition, by DR. HENRY LEFFMANN. *Just Ready.* .75

MICROSCOPY.

CARPENTER. The Microscope and Its Revelations. 8th Edition, Revised and Enlarged. 817 Illustrations and 23 Plates. Cloth, $8.00; Half Morocco, $9.00

GREENISH. Microscopical Examination of Foods and Drugs. Illustrated. *Just Ready.* $3.50

LEE. The Microtomist's Vade Mecum. A Handbook of Methods of Microscopical Anatomy. 887 Articles. 5th Edition, Enlarged. $4.00

OERTEL. Medical Microscopy. A Guide to Diagnosis, Elementary Laboratory Methods and Microscopic Technic. 131 Illustrations. *Just Ready.* $2.00

REEVES. Medical Microscopy, including Chapters on Bacteriology, Neoplasms, Urinary Examination, etc. Numerous Illustrations, some of which are printed in colors. $2.50

WETHERED. Medical Microscopy. A Guide to the Use of the Microscope in Practical Medicine. 100 Illustrations. $2.00

MISCELLANEOUS.

BERRY. Diseases of Thyroid Gland. Illustrated. $4.00

BUXTON. Anesthetics. Illustrated. 3d Edition. $1.50

COHEN. Organotherapy. *See Cohen, Physiologic Therapeutics, page 17.*

FRENKEL. Tabetic Ataxia. Illustrated. $3.00

GOULD. Borderland Studies. Miscellaneous Addresses and Essays. 12mo. $2.00

GOULD. Biographic Clinics. The Origin of the Ill-Health of DeQuincy, Carlyle, Darwin, Huxley, and Browning. *Just Ready.* $1.00

GREENE. Medical Examination for Life Insurance. Illus. With colored and other Engravings. 2d Edition. *In Press.*

HAIG. Causation of Disease by Uric Acid. The Pathology of High Arterial Tension, Headache, Epilepsy, Gout, Rheumatism, Diabetes, Bright's Disease, etc. 6th Edition. $3.50

HENRY. A Practical Treatise on Anemia. Half Cloth, .50

NEW SYDENHAM SOCIETY'S PUBLICATIONS. Circulars upon application. Per Annum, $8.00

OSGOOD. The Winter and Its Dangers. .40

PACKARD. Sea Air and Sea Bathing. .40

RICHARDSON. Long Life and How to Reach It. .40

SCHEUBE. Diseases of Warm Countries. Illustrated. *Just Ready.* $8.00

TISSIER. Pneumotherapy, Aerotherapy, Inhalation Methods. *See Cohen, Physiologic Therapeutics, page* 17.

TURNBULL. Artificial Anesthesia. 4th Ed. Illus. $2.50

WARDEN. The Paris Medical School. Paper, .75

WEBER AND HINSDALE. Climatology and Health Resorts. Including Mineral Springs. 2 vols. Illustrated with Colored Maps. *See Cohen, Physiologic Therapeutics, page* 17.

WILSON. The Summer and Its Diseases. .40

WINTERNITZ. Hydrotherapy, Thermotherapy, Phototherapy, Mineral Waters, Baths, etc. Illustrated. *See Cohen, Physiologic Therapeutics, page* 17.

NERVOUS DISEASES.

DERCUM. Rest, Suggestion, Mental Therapeutics. *See Cohen, Physiologic Therapeutics, page* 17.

GORDINIER. The Gross and Minute Anatomy of the Central Nervous System. With 271 original colored and other Illustrations. Cloth, $6.00; Sheep, $7.00

GOWERS. Syphilis and the Nervous System. $1.00

GOWERS. Manual of Diseases of the Nervous System. A Complete Text-Book. Revised, Enlarged, and in many parts Rewritten. With many new Illustrations. Two volumes.
Vol. I. Diseases of the Nerves and Spinal Cord. 3d Edition, Enlarged. Cloth, $4.00; Sheep, $5.00
Vol. II. Diseases of the Brain and Cranial Nerves; General and Functional Disease. 2d Ed. Cloth, $4.00; Sheep, $5.00

GOWERS. Epilepsy and Other Chronic Convulsive Diseases. 2d Edition. $3.00

HORSLEY. The Brain and Spinal Cord, the Structure and Functions of. Numerous Illustrations. $2.50

ORMEROD. Diseases of the Nervous System. 66 Wood Engravings. $1.00

PERSHING. Diagnosis of Nervous and Mental Diseases. Illustrated. $1.25

PRESTON. Hysteria and Certain Allied Conditions. Their Nature and Treatment. Illustrated. $2.00

WOOD. Brain Work and Overwork. .40

NURSING (see also Massage).

Special Catalogue of Books for Nurses sent free upon application.

CANFIELD. Hygiene of the Sick-Room. A Book for Nurses and Others. Being a Brief Consideration of Asepsis, Antisepsis, Disinfection, Bacteriology, Immunity, Heating and Ventilation, and Kindred Subjects for the Use of Nurses and Other Intelligent Women. $1.25

CUFF. Lectures to Nurses on Medicine. 4th Edition. $1.25

DAVIS. Bandaging. Its Principles and Practice. 163 Original Illustrations. $1.50

DOMVILLE. Manual for Nurses and Others Engaged in Attending the Sick. 9th Edition. With Recipes for Sick-room Cookery, etc. *In Press.*

FULLERTON. Obstetric Nursing. 5th Ed. 41 Illus. $1.00

FULLERTON. Surgical Nursing. 3d Ed. 69 Illus. $1.00

GROFF. Materia Medica for Nurses. With Questions for Self-Examination. 2d Edition, Revised and Improved. *Just Ready.* $1.25

HADLEY. General, Medical, and Surgical Nursing. A very Complete Manual, Including Sick-room Cookery. $1.25

HUMPHREY. A Manual for Nurses. Including General Anatomy and Physiology, Management of the Sick-room, etc. 24th Edition. 79 Illustrations. $1.00

STARR. The Hygiene of the Nursery. Including the General Regimen and Feeding of Infants and Children, and the Domestic Management of the Ordinary Emergencies of Early Life, Massage, etc. 6th Edition. 25 Illustrations. $1.00

TEMPERATURE AND CLINICAL CHARTS. See page 25.

VOSWINKEL. Surgical Nursing. Second Edition, Enlarged. 112 Illustrations. $1.00

WILCOX. Fever Nursing. *Preparing.*

OBSTETRICS.

CAZEAUX AND TARNIER. Midwifery. With Appendix by MUNDÉ. The Theory and Practice of Obstetrics, including the Diseases of Pregnancy and Parturition, Obstetrical Operations, etc. 8th Edition. Illustrated by colored and other full-page Plates, and numerous Wood Engravings.
Cloth, $4.50; Full Leather, $5.50

EDGAR. Text-Book of Obstetrics. By J. CLIFTON EDGAR, M.D., Professor of Obstetrics and Clinical Midwifery, Medical Department of Cornell University, New York City, etc. 1221 Illustrations. *Just Ready.* Cloth, $6.00; Sheep, $7.00

FULLERTON. Obstetric Nursing. 5th Ed. Illus. $1.00

LANDIS. Compend of Obstetrics. 7th Edition, Revised by WM. H. WELLS, M.D., Demonstrator of Clinical Obstetrics, Jefferson Medical College. 52 Illustrations.
.80; Interleaved, $1.00

WINCKEL. Text-Book of Obstetrics, Including the Pathology and Therapeutics of the Puerperal State. Illustrated. $5.00

PATHOLOGY.

BLACKBURN. Autopsies. A Manual of Autopsies Designed for the Use of Hospitals for the Insane and other Public Institutions. Ten full-page Plates and other Illustrations. $1.25

COPLIN. Manual of Pathology. Including Bacteriology, Technic of Post-Mortems, Methods of Pathologic Research, etc. 330 Illustrations, 7 Colored Plates. 3d Edition. $3.50

DA COSTA. Clinical Hematology. A Practical Guide to the Examination of the Blood. Six Colored Plates and 48 Illustrations. Cloth, $5.00; Sheep, $6.00

LAZARUS-BARLOW. Pathological Anatomy. With 7 Colored Plates and 171 other Illustrations. $6.50

MacLEOD. The Pathology of the Skin. Colored and other Illustrations. *Just Ready.* $5.00

MARTIN. Manual of Pathology. Illustrated. *Nearly Ready.*

ROBERTS. Gynecological Pathology. Illustrated. $6.00

THAYER. Compend of General Pathology. Illustrated. .80; Interleaved, $1.00

THAYER. Compend of Special Pathology. Illustrated. .80; Interleaved, $1.00

THAYER. Manual of General and Special Pathology. Illustrated. *Nearly Ready.*

VIRCHOW. Post-Mortem Examinations. 3d Edition. .75

WHITACRE. Laboratory Text-Book of Pathology. With 121 Illustrations. $1.50

PHARMACY.

Special Catalogue of Books on Pharmacy sent free upon application.

COBLENTZ. Manual of Pharmacy. A Complete Text-Book by the Professor in the New York College of Pharmacy. 2d Ed., Revised and Enlarged. 437 Illus. Cloth, $3.50; Sheep, $4.50

COBLENTZ. Volumetric Analysis. Illustrated. $1.25

BEASLEY. Book of 3100 Prescriptions. Collected from the Practice of the Most Eminent Physicians and Surgeons—English, French, and American. A Compendious History of the Materia Medica, Lists of the Doses of all the Officinal and Established Preparations, an Index of Diseases and their Remedies. 7th Edition. $2.00

BEASLEY. Druggists' General Receipt Book. Comprising a Copious Veterinary Formulary, Recipes in Patent and Proprietary Medicines, Druggists' Nostrums, etc.; Perfumery and Cosmetics, Beverages, Dietetic Articles and Condiments, Trade Chemicals, Scientific Processes, and many Useful Tables. 10th Edition. $2.00

BEASLEY. Pharmaceutical Formulary. A Synopsis of the British, French, German, and United States Pharmacopœias. Comprising Standard and Approved Formulæ for the Preparations and Compounds Employed in Medicine. 12th Ed. $2.00

GREENISH. Microscopical Examination of Foods and Drugs. Illustrated. *Just Ready.* $3.50

ROBINSON. Latin Grammar of Pharmacy and Medicine. 4th Edition. With elaborate Vocabularies. *Just Ready.* $1.50

SAYRE. Organic Materia Medica and Pharmacognosy. An Introduction to the Study of the Vegetable Kingdom and the Vegetable and Animal Drugs. Comprising the Botanical and Physical Characteristics, Source, Constituents, and Pharmacopeial Preparations, Insects Injurious to Drugs, and Pharmacal Botany. With sections on Histology and Microtechnique, by W. C. STEVENS. 374 Illustrations. Second Edition. Cloth, $4.50

SCOVILLE. **The Art of Compounding.** Second Edition, Revised and Enlarged. Cloth, $2.50

STEWART. **Compend of Pharmacy.** Based upon "Remington's Text-Book of Pharmacy." 5th Edition, Revised in Accordance with the U. S. Pharmacopœia, 1890. Complete Tables of Metric and English Weights and Measures. .80; Interleaved, $1.00

TAVERA. **Medicinal Plants of the Philippines.** $2.00

UNITED STATES PHARMACOPŒIA. 7th Decennial Revision. Cloth, $2.50 (postpaid, $2.77); Sheep, $3.00 (postpaid, $3.27); Interleaved, $4.00 (postpaid, $4.50); Printed on one side of page only, unbound, $3.50 (postpaid, $3.90).

Select Tables from the U. S. P. Being Nine of the Most Important and Useful Tables, Printed on Separate Sheets. .25

POTTER. **Handbook of Materia Medica, Pharmacy, and Therapeutics.** 600 Prescriptions. 9th Edition. Cloth, $5.00; Sheep, $6.00

PHYSIOLOGY.

BIRCH. **Practical Physiology.** An Elementary Class Book. 62 Illustrations. $1.75

BRUBAKER. **Text-Book of Physiology.** Illus. *Nearly Ready.*

BRUBAKER. **Compend of Physiology.** 11th Edition, Revised and Enlarged. Illustrated. .80; Interleaved, $1.00

JONES. **Outlines of Physiology.** 96 Illustrations. $1.50

KIRKES. **Handbook of Physiology.** 17th Authorized Edition. Revised, Rearranged, and Enlarged. By Prof. W. D. Halliburton, of Kings College, London. 681 Illustrations, some of which are in colors. Cloth, $3.00; Leather, $3.75

LANDOIS. **A Text-Book of Human Physiology.** Including Histology and Microscopical Anatomy, with Special Reference to the Requirements of Practical Medicine. 5th American, translated and edited from the last German Edition by A. P. Brubaker, M.D., and A. A. Eshner, M.D. *In Press.*

STARLING. **Elements of Human Physiology.** 100 Illus. $1.00

STIRLING. **Outlines of Practical Physiology.** Including Chemical and Experimental Physiology, with Special Reference to Practical Medicine. 3d Edition. 289 Illustrations. $2.00

TYSON. **Cell Doctrine.** Its History and Present State. $1.50

PRACTICE.

BEALE. **On Slight Ailments: their Nature and Treatment.** 2d Edition, Enlarged and Illustrated. $1.25

COHEN. **Physiologic Therapeutics.** The Treatment of Disease by Methods other than Drug-giving. *See page* 17.

FAGGE. **Practice of Medicine.** 4th Edition, by P. H. Pye-Smith, M.D. 2 volumes. Vol. I, $6.00; Vol. II, $6.00

FOWLER. **Dictionary of Practical Medicine.** By various writers. An Encyclopædia of Medicine. Cloth, $3.00; Half Morocco, $4.00

GOULD AND PYLE. Cyclopedia of Practical Medicine and Surgery. A Concise Reference Handbook, with particular Reference to Diagnosis and Treatment. Edited by DRS. GOULD and PYLE, Assisted by 72 Special Contributors. Illustrated, one volume. Large Square Octavo, Uniform with "Gould's Illustrated Dictionary."

Sheep or Half Mor., $10.00; with Thumb Index, $11.00
Half Russia, Thumb Index, $12.00

☞ *Complete descriptive circular free upon application.*

GOULD AND PYLE'S Pocket Cyclopedia of Medicine and Surgery. Based upon the above and Uniform with "Gould's Pocket Dictionary." Full Limp Leather, Gilt Edges, Round Corners, $1.00; with Thumb Index, $1.25.

HUGHES. Compend of the Practice of Medicine. 6th Edition, Revised and Enlarged.

Part I. Continued, Eruptive, and Periodical Fevers, Disease of the Stomach, Intestines, Peritoneum, Biliary Passages, Liver, Kidneys, etc., and General Diseases, etc.

Part II. Diseases of the Respiratory System, Circulatory System, and Nervous System; Diseases of the Blood, etc.

Price of each part, .80; Interleaved, $1.00

Physician's Edition. In one volume, including the above two parts, a Section on Skin Diseases, and an Index. 6th Revised Edition. 625 pp. Full Morocco, Gilt Edge, $2.25

TAYLOR. Practice of Medicine. 6th Edition. $4.00

TYSON. The Practice of Medicine. By JAMES TYSON, M.D. Professor of Medicine in the University of Pennsylvania. Complete Systematic Text-book, with Special Reference to Diagnosis and Treatment. 3d Edition, Enlarged and Revised. Colored Plates and 124 other Illustrations.

Cloth, $5.50; Leather, $6.50

STOMACH. INTESTINES.

FENWICK. Cancer of the Stomach. *Just Ready.* $3.00

HEMMETER. Diseases of the Stomach. Their Special Pathology, Diagnosis, and Treatment. With Sections on Anatomy, Analysis of Stomach Contents, Dietetics, Surgery of the Stomach, etc. 3d Edition, Revised. With 15 Plates and 41 other Illustrations, a number of which are in colors.

Cloth, $6.00; Sheep, $7.00

HEMMETER. Diseases of the Intestines. Their Special Pathology, Diagnosis, and Treatment. With Sections on Anatomy and Physiology, Microscopic and Chemic Examination of Intestinal Contents, Secretions, Feces and Urine, Intestinal Bacteria and Parasites, Surgery of the Intestines, Dietetics, Diseases of the Rectum, etc. With Full-page Colored Plates and many other Original Illustrations. 2 volumes. Octavo.

Price of each volume, Cloth, $5.00; Sheep, $6.00

SKIN.

BULKLEY. The Skin in Health and Disease. Illustrated. .40

CROCKER. Diseases of the Skin. Their Description, Pathology, Diagnosis, and Treatment, with Special Reference to the Skin Eruptions of Children. 3d Edition, Thoroughly Revised. With New Illus. *Just Ready.* Cloth, $5.00; Sheep, $6.00

MacLEOD. The Pathology of the Skin. Colored and other Illustrations. *Just Ready.* $5.00

SCHAMBERG. Diseases of the Skin. 3d Edition, Revised and Enlarged. 106 Illustrations. Being No. 16 ? Quiz-Compend? Series. *Just Ready.* Cloth, .80; Interleaved, $1.00

VAN HARLINGEN. On Skin Diseases. A Practical Manual of Diagnosis and Treatment, with Special Reference to Differential Diagnosis. 3d Edition, Revised and Enlarged. With Formulæ and 60 Illustrations, some of which are printed in colors. $2.75

SURGERY AND SURGICAL DISEASES (see also Urinary Organs).

BERRY. Diseases of the Thyroid Gland. Illustrated. $4.00

BUTLIN. Operative Surgery of Malignant Disease. 2d Edition. Illustrated. Octavo. $4.50

CASPER AND RICHTER. Functional Kidney Diagnosis. $1.50

DAVIS. Bandaging. Its Principles and Practice. 163 Original Illustrations. $1.50

DEAVER. Surgical Anatomy. A Treatise on Human Anatomy in its Application to Medicine and Surgery. With about 500 very handsome full-page Illustrations Engraved from Original Drawings made by special Artists from Dissections prepared for the purpose. Three volumes. Royal Square Octavo. By *Subscription only. Now Ready.*
Half Morocco or Sheep, $24.00; Half Russia, $27.00

DEAVER. Appendicitis: its Symptoms, Diagnosis, Pathology, Treatment, and Complications. Elaborately Illustrated with Colored Plates and other Illus. 3d Edition. *Preparing.*

DOUGLAS. Surgical Diseases of the Abdomen. Illustrated by 20 Full-page Plates. *Just Ready.*
Cloth, $7.00; Sheep, $8.00

DULLES. What to do First in Accidents and Poisoning. 5th Edition. New Illustrations. $1.00

FULLERTON. Surgical Nursing. 3d Ed. 69 Illus. $1.00

HAMILTON. Lectures on Tumors. 3d Edition. $1.25

HEATH. Minor Surgery and Bandaging. 12th Edition, Revised and Enlarged. 195 Illus., Formulæ, Diet List, etc. $1.50

HEATH. Clinical Lectures on Surgical Subjects. Second Series. $2.00

HORWITZ. Compend of Surgery and Bandaging. Including Minor Surgery, Amputations, Fractures, Dislocations, Surgical Diseases, etc., with Differential Diagnosis and Treatment. 5th Edition, very much Enlarged and Rearranged. 167 Illus., 98 Formulæ. Cloth, .80; Interleaved, $1.00

JACOBSON. Operations of Surgery. 4th Ed., Enlarged. 550 Illus. Two volumes. Cloth, $10.00; Leather, $12.00

KEAY. Medical Treatment of Gall-Stones. $1.25

KEHR. Gall-stone Disease. Translated by WILLIAM WOTKYNS SEYMOUR, M.D. $2.50

MAKINS. Surgical Experiences in South Africa. 1899–1900. Illustrated. $4.00

MAYLARD. Surgery of the Alimentary Canal. 97 Illustrations. 2d Edition, Revised. $3.00

MOULLIN. Text-Book of Surgery. With Special Reference to Treatment. 3d American Edition. Revised and edited by JOHN B. HAMILTON, M.D., LL.D., Professor of the Principles of Surgery and Clinical Surgery, Rush Medical College, Chicago. 623 Illustrations, many of which are printed in colors. Cloth, $6.00; Leather, $7.00

SMITH. Abdominal Surgery. Being a Systematic Description of all the Principal Operations. 224 Illustrations. 6th Edition. 2 volumes. Cloth, $10.00

VOSWINKEL. Surgical Nursing. Second Edition, Revised and Enlarged. 111 Illustrations. $1.00

WALSHAM. Manual of Practical Surgery. 7th Ed., Revised and Enlarged. 483 Engravings. 950 pages. $3.50

TEMPERATURE CHARTS, ETC.

GRIFFITH. Graphic Clinical Chart for Recording Temperature, Respiration, Pulse, Day of Disease, Date, Age, Sex, Occupation, Name, etc. Printed in three colors. Sample copies free. Put up in loose packages of fifty, 50 cts. Price to Hospitals, 500 copies, $4.00; 1000 copies, $7.50.

KEEN'S Clinical Charts. Seven Outline Drawings of the Body, on which may be marked the Course of Disease, Fractures, Operations, etc. Each Drawing may be had separately, twenty-five to pad, 25 cents.

THROAT AND NOSE (see also Ear).

COHEN. The Throat and Voice. Illustrated. .40

HALL. Diseases of the Nose and Throat. 2d Edition, Enlarged. Two Colored Plates and 80 Illustrations. $2.75

HOLLOPETER. Hay Fever. Its Successful Treatment. $1.00

KNIGHT. Diseases of the Throat. A Manual for Students. Illustrated. *Nearly Ready.*

KYLE (J. J.). Diseases of the Ear, Nose, and Throat. A Compend for Students. Illustrated. .80; Interleaved, $1.00

McBRIDE. Diseases of the Throat, Nose, and Ear. With Colored Illustrations from Original Drawings. 3d Ed. $7.00

POTTER. Speech and its Defects. Considered Physiologically, Pathologically, and Remedially. $1.00

URINE AND URINARY ORGANS.

ACTON. The Functions and Disorders of the Reproductive Organs in Childhood, Youth, Adult Age, and Advanced Life, Considered in their Physiological, Social, and Moral Relations. 8th Edition. $1.75

CASPER AND RICHTER. Functional Kidney Diagnosis. $1.50

HOLLAND. The Urine, the Gastric Contents, the Common Poisons, and the Milk. Memoranda, Chemical and Microscopical, for Laboratory Use. Illustrated and Interleaved. 6th Edition. $1.00

KLEEN. Diabetes and Glycosuria. $2.50

MEMMINGER. Diagnosis by the Urine. 2d Edition. 24 Illustrations. $1.00

MORRIS. Renal Surgery, with Special Reference to Stone in the Kidney and Ureter and to the Surgical Treatment of Calculous Anuria. Illustrated. $2.00

MOULLIN. Enlargement of the Prostate. Its Treatment and Radical Cure. 2d Edition. Illustrated. $1.75

MOULLIN. Inflammation of the Bladder and Urinary Fever. Octavo. $1.50

SCOTT. The Urine. Its Clinical and Microscopical Examination. 41 Lithographic Plates and other Illustrations. Quarto. Cloth, $5.00

TYSON. Guide to Examination of the Urine. For the Use of Physicians and Students. With Colored Plate and Numerous Illustrations engraved on wood. 10th Edition, Revised, Enlarged, and partly Rewritten. With New Illustrations. *Just Ready.* $1.50

VAN NUYS. Chemical Analysis of Urine. 39 Illus. $1.00

VENEREAL DISEASES.

GOWERS. Syphilis and the Nervous System. $1.00

STURGIS AND CABOT. Student's Manual of Venereal Diseases. 7th Revised and Enlarged Edition. 12mo. $1.25

VETERINARY.

BALLOU. Equine Anatomy and Physiology. 29 Graphic Illustrations. .80; Interleaved, $1.00

JACOBSON. The Operations of Surgery. By W. H. A. JACOBSON, F.R.C.S., Surgeon to Guy's Hospital; Consulting Surgeon Royal Hospital for Children and Women; and F. J. STEWARD, F.R.C.S., Assistant Surgeon Guy's Hospital. Fourth Edition—Revised, Enlarged, and Improved. 550 Illustrations. Two Volumes, Octavo, 1524 pages.

Cloth, $10.00; Sheep, $12.00

"The important anatomical points are clearly set forth, the conditions indicating or contraindicating operative interference are given, the details of the operations themselves are brought forward prominently, and frequently the after-treatment is considered. Herein is one of the strong points of the book."—*New York Medical Journal.*

"We know of no series of books issued by any house that so fully meets our approval as these ? Quiz-Compends?. They are well arranged, full and concise, and are really the best line of text-books that could be found for either student or practitioner."
—*Southern Clinic.*

BLAKISTON'S ?QUIZ-COMPENDS?

The Best Series of Manuals for the Use of Students.

Price of each, Cloth, .80. Interleaved, for taking Notes, $1.00

These Compends are based on the most popular text-books and the lectures of prominent professors, and are kept constantly revised, so that they may thoroughly represent the present state of the subjects upon which they treat. The authors have had large experience as Quiz-Masters and attaches of colleges, and are well acquainted with the wants of students. They are arranged in the most approved form, thorough and concise, containing nearly 1000 illustrations and lithograph plates, inserted wherever they could be used to advantage. Can be used by students of *any* college. They contain information nowhere else collected in such a condensed, practical shape.

No. 1. POTTER. HUMAN ANATOMY. Seventh Edition. 138 Illustrations and 16 Plates of Nerves and Arteries.

No. 2. HUGHES. PRACTICE OF MEDICINE. Part I. Sixth Edition, Enlarged and Improved.

No. 3. HUGHES. PRACTICE OF MEDICINE. Part II. Sixth Edition, Revised and Improved.

No. 4. BRUBAKER. PHYSIOLOGY. Eleventh Edition. Illus.

No. 5. LANDIS. OBSTETRICS. Seventh Edition. 52 Illus.

No. 6. POTTER. MATERIA MEDICA, THERAPEUTICS, AND PRESCRIPTION WRITING. Sixth Revised Edition.

No. 7. WELLS. GYNECOLOGY. Third Edition. 140 Illus.

No. 8. GOULD AND PYLE. DISEASES OF THE EYE. Second Edition. Refraction, Treatment, Surgery, etc. 109 Illus.

No. 9. HORWITZ. SURGERY. Including Minor Surgery, Bandaging, Surgical Diseases, Differential Diagnosis and Treatment. Fifth Edition. With 98 Formulæ and 71 Illustrations.

No. 10. LEFFMANN. MEDICAL CHEMISTRY. Fourth Edition. Including Urinalysis, Animal Chemistry, Chemistry of Milk, Blood, Tissues, the Secretions, etc.

No. 11. STEWART. PHARMACY. Fifth Edition. Based upon Prof. Remington's Text-Book of Pharmacy.

No. 12. BALLOU. EQUINE ANATOMY AND PHYSIOLOGY. 29 graphic Illustrations.

No. 13. WARREN. DENTAL PATHOLOGY AND DENTAL MEDICINE. Third Edition, Illustrated.

No. 14. HATFIELD. DISEASES OF CHILDREN. 3d Edition.

No. 15. THAYER. GENERAL PATHOLOGY. 78 Illus.

No. 16. SCHAMBERG. DISEASES OF THE SKIN. Third Edition, Revised and Enlarged. 106 Illustrations.

No. 17. CUSHING. HISTOLOGY. Illustrated. *In Press.*

No. 18. THAYER. SPECIAL PATHOLOGY. 34 Illustrations.

No. 19. KYLE. DISEASES OF THE EAR, NOSE, AND THROAT. Illustrated.

DA COSTA

Clinical Hematology

A Practical Guide to the Examination of the Blood by Clinical Methods. With Reference to the Diagnosis of Disease. With Colored Illustrations. Cloth, $5.00

*** A new, thorough, systematic, and comprehensive work, its purpose being, first, to show how to examine the blood, and second, how to diagnose from such examination diseases of the blood itself and general diseases. The author's aim has been to cover not alone the field of original research, but to supply a book for the student, the hospital physician and the general practitioner. It will be found wanting in none of these respects.

OERTEL

Medical Microscopy

JUST READY

A GUIDE TO DIAGNOSIS, ELEMENTARY LABORATORY METHODS, AND MICROSCOPIC TECHNIC

By T. E. Oertel, M.D.,

Professor of Pathology and Clinical Microscopy, Medical Department, University of Georgia.

WITH 131 ILLUSTRATIONS. 12mo. Cloth, $2.00

The Pocket Cyclopedia of Medicine and Surgery

Full Limp Leather, Round Corners, Gilt Edges, $1.00
With Thumb Index, $1.25

Uniform with "Gould's Pocket Dictionary"

A concise practical volume of nearly 600 pages, containing a vast amount of information on all medical subjects, including Diagnosis and Treatment of Disease, with Formulas and Prescriptions, Emergencies, Poisons, Drugs and Their Uses, Nursing, Surgical Procedures, Dose List in both English and Metric Systems, etc.

By Drs. Gould and Pyle

Based upon their large "Cyclopedia of Medicine and Surgery."

*** This is a new book which will prove of the greatest value to students. It is to the broad field of general medical information what "Gould's Pocket Dictionary" is to the more special one of definition and pronunciation of words. The articles are concise but thorough, and arranged in shape for quick reference. In no other book can be found so much exact detailed knowledge so conveniently classified, so evenly distributed, so methodically grouped. It is *Multum in Parvo*. **Sample Pages Free.**

A NEW EDITION

CROCKER ON THE SKIN

The Diseases of the Skin. Their Description, Pathology, Diagnosis, and Treatment, with Special Reference to the Skin Eruptions of Children. By H. RADCLIFFE CROCKER, M.D., Physician to the Department of Skin Diseases, University College Hospital, London. With new Illustrations.

Third Edition, Rewritten and Enlarged

OCTAVO. JUST READY; CLOTH, $5.00

*** This new edition will easily hold the high position given the previous printings. The author is a member of American, English, French, German, and Italian Dermatological Societies, and a recognized authority the world over.

STURGIS—MANUAL OF VENEREAL DISEASES

By F. R. STURGIS, M.D., Sometime Clinical Professor of Venereal Diseases in the Medical Department of the University of the City of New York. Seventh Edition, Revised and in Part Rewritten by the Author and FOLLEN CABOT, M.D., Instructor in Genito-Urinary and Venereal Diseases in the Cornell University Medical College. 12mo. 216 pages. Cloth, $1.25

*** This manual was originally written for students' use, and is as concise and as practical as possible. It presents a careful, condensed description of the commoner forms of venereal diseases which occur in the practice of the general physician, together with the most approved remedies.

FOR THE DISSECTING ROOM

Holden's Anatomy—Seventh Edition
320 Illustrations

A Manual of the Dissections of the Human Body. By JOHN LANGTON, F.R.C.S. Carefully Revised by A. HEWSON, M.D., Demonstrator of Anatomy, Jefferson Medical College, Philadelphia, etc. 320 Illustrations. Two small compact volumes. 12mo.

Vol. I. Scalp, Face, Orbit, Neck, Throat, Thorax, Upper Extremity. 435 pages. 153 Illustrations. Oil Cloth, $1.50

Vol. II. Abdomen, Perineum, Lower Extremity, Brain, Eye, Ear, Mammary Gland, Scrotum, Testes. 445 pages. 167 Illustrations. Oil Cloth, $1.50

Each volume sold separately.

Hughes and Keith—Dissections Illustrated

A Manual of Dissections by ALFRED W. HUGHES, M.B., M.R.C.S. (Edin.), late Professor of Anatomy and Dean of Medical Faculty, King's College, London, etc., and ARTHUR KEITH, M.D., Joint Lecturer on Anatomy, London Hospital Medical College, etc. In three parts. With 527 Colored and other Illustrations.

I. Upper and Lower Extremity. 38 Plates, 116 other Illustrations. Cloth, $3.00

II. Abdomen. Thorax. 4 Plates, 149 other Illustrations. Cloth, $3.00

III. Head, Neck, and Central Nervous System. 16 Plates, 204 other Illustrations. Cloth, $3.00

Each volume sold separately.

*** The student will find it of great advantage to have a "Dissector" to supplement his regular text-book on anatomy. These books meet all requirements, and as they can be purchased in parts as wanted, the outlay is small.

EDGAR'S

OBSTETRICS

A NEW TEXT-BOOK

1221 Illustrations

Edgar's Obstetrics excels all other works on this subject in completeness, in uniformity and consistency in arrangement, thoroughness and clearness in handling details, and in the number and usefulness of its illustrations.

OCTAVO. CLOTH, $6.00; SHEEP, $7.00

AUG 25 1903

CARD EXPLANATORY OF HOLMGREN'S TESTS FOR COLOUR BLINDNESS.

Test I.



Confusion Colours.

1 2 3 4 5



Test II. *a*.



Confusion Colours. Red Blindness.

6 7



Confusion Colours. Green Blindness.

8 9



Test II. *b*.



Confusion Colours. Red Blindness.

10 11



Confusion Colours. Green Blindness.

12 18



N.B.—*This Card is merely intended to illustrate the text (The Colour Sense, Chap. I. and Appendix I.), and is not itself for use as a test.*

LIBRARY OF CONGRESS
0 021 062 797 2